

CONTENTS

	Page
บทคัดย่อ	iii
ABSTRACT	iv
ACKNOWLEDGEMENT	v
CONTENTS	vi
LIST OF TABLES	ix
LIST OF FIGURES	xi
CHAPTER	
1 INTRODUCTION	
1.1 General introduction	1
1.2 Objectives	3
2 LITERATURE REVIEW	
2.1 Free radicals and skin aging	4
2.2 Defense mechanism of skin against oxidative damage	6
2.3 Pomegranate	7
2.3.1 Chemistry	9
2.3.1.1 Seed	14
2.3.1.2 Juice	14
2.3.1.3 Pericarp (peel, rind, hull are synonymns)	14
2.3.1.4 Leaf	14
2.3.1.5 Flower	15
2.3.1.6 Tree bark and roots	15
2.3.2 Medicinal properties	15
2.3.3 Biological activities of pomegranate	16
3 MATERIALS AND METHODS	
3.1 Materials	
3.1.1 Plant materials	20
3.1.2 Chemical	20

CONTENTS (Continued)

	Page
4.2 Determination of suitable solvent for fractionation	30
4.3 Determination of extraction method	31
4.4 Preparation of antioxidant active extract of pomegranate fruit peels	32
4.5 Quantitative determination of ellagic acid in the antioxidant active extract of pomegranate fruit peels	35
4.6 Solubility of the antioxidant active extract of pomegranate fruit peels	38
4.7 Stability of the antioxidant active extract of pomegranate fruit peels	
4.7.1 Effect of light	38
4.7.2 Effect of temperature	44
4.7.3 The effect of humidity	50
4.8 Preliminary formulation study of pomegranate-extract cream	57
5 CONCLUSIONS	65
BIBLIOGRAPHY	67
VITAE	78

LIST OF TABLES

Table	Page
2.1 Reactive oxygen species (ROS)	5
2.2 Chemical constituents of pomegranate	9
3.1 The ingredients of the cream bases	28
4.1 Antioxidant activity of the pomegranate fruit peel extracts and quercetin evaluated by DPPH radical scavenging assay	30
4.2 Antioxidant activity of the fractions obtained from liquid-liquid extraction evaluated by DPPH radical scavenging assay	31
4.3 Comparison of extraction time and antioxidant activity of the extracts obtained from two extraction methods	32
4.4 Antioxidant activity of the EtOAc fraction evaluated by DPPH radical scavenging assay and β -carotene bleaching assay	33
4.5 Solubility of pomegranate fruit peel extract with various cosmetic solvents	38
4.6 Ellagic acid content in the antioxidant active extract of pomegranate fruit peels stored at $45 \pm 2^{\circ}\text{C}$, under exposed to light and protected from light conditions	40
4.7 Antioxidant activity of the antioxidant active extract of pomegranate fruit peels stored at $45 \pm 2^{\circ}\text{C}$, under exposed to light and protected from light conditions	42
4.8 Ellagic acid content in the antioxidant active extract of pomegranate fruit peels stored at $25 \pm 2^{\circ}\text{C}$ and $45 \pm 2^{\circ}\text{C}$, protected from light	45
4.9 Antioxidant activity of the antioxidant active extract of pomegranate fruit peels stored at $25 \pm 2^{\circ}\text{C}$ and $45 \pm 2^{\circ}\text{C}$, protected from light	47
4.10 Ellagic acid content in the antioxidant active extract of pomegranate fruit peels stored under 75% relative humidity and protected from humidity conditions, at $25 \pm 2^{\circ}\text{C}$	51

LIST OF TABLES (Continued)

Table	Page
4.11 Antioxidant activity of the antioxidant active extract of pomegranate fruit peels stored under 75% relative humidity and protected from humidity conditions, at 25 ± 2 °C	53
4.12 Physical properties of cream bases before and after heating-cooling test (8 cycles)	58
4.13 Physical properties of pomegranate-extract cream containing 0.5% (Rx 5-1) and 1% (Rx 5-2) the antioxidant active extract of pomegranate fruit peels before and after heating and cooling test (8 cycles)	60
4.14 Ellagic acid content in pomegranate extract creams, before and after heating-cooling test (8 cycles)	62
4.15 Antioxidant activity of pomegranate extract creams evaluated by DPPH assay	63
4.16 Antioxidant activity of pomegranate extract creams evaluated by β -carotene bleaching assay	63

LIST OF FIGURES

Figure	Page
2.1 Skin exposed to ultraviolet (UV) light	4
2.2 Skin aging	6
2.3 Pomegranate fruit	8
2.4 Structure of some interested polyphenolic compounds	12
4.1 Antioxidant active extract of pomegranate fruit peels	32
4.2 Correlation between concentration of EtOAc fraction ($\mu\text{g/ml}$) determined by DPPH radical scavenging assay and its inhibition (%)	33
4.3 Correlation between concentration of ellagic acid ($\mu\text{g/ml}$) determined by DPPH radical scavenging assay and its inhibition (%)	34
4.4 Correlation between concentration of quercetin ($\mu\text{g/ml}$) determined by DPPH radical scavenging assay and its inhibition (%)	34
4.5 HPLC-chromatogram of the authentic ellagic acid (A) and the antioxidant active extract of pomegranate fruit peels (B)	36
4.6 Absorption spectra of the peak at retention time 6.28 minutes and the authentic ellagic acid	37
4.7 Calibration curve of ellagic acid	37
4.8 Color of the antioxidant active extract of pomegranate fruit peels stored under protected from light (A) and stored under light condition (B) after storage at $45 \pm 2^\circ\text{C}$	39
4.9 Ellagic acid content in the antioxidant active extract of pomegranate fruit peels stored at $45 \pm 2^\circ\text{C}$, under exposed to light and protected from light conditions (*Significance at $P < 0.05$ when compared with the content at initial time)	41

LIST OF FIGURES (Continued)

Figure	Page
4.10 Antioxidant activity evaluated by DPPH assay of antioxidant active extract pomegranate fruit peel stored at 45 ± 2 °C, under exposed to light and protected from light conditions (*Significance at $P < 0.05$ when compared with the content at initial time)	43
4.11 Antioxidant activity evaluated by β -carotene bleaching assay of antioxidant active extract pomegranate fruit peel stored at 45 ± 2 °C, under exposed to light and protected from light conditions (*Significance at $P < 0.05$ when compared with the content at initial time)	43
4.12 Ellagic acid content in the antioxidant active extract of pomegranate fruit peels stored at 25 ± 2 °C and 45 ± 2 °C, protected from light (*Significance at $P < 0.05$ when compared with the content at initial time)	46
4.13 Antioxidant activity evaluated by DPPH assay of the antioxidant active extract of pomegranate fruit peels stored at 25 ± 2 °C and 45 ± 2 °C, protected from light (*Significance at $P < 0.05$ when compared with the content at initial time)	48
4.14 Antioxidant activity evaluated by β -carotene bleaching assay of the antioxidant active extract of pomegranate fruit peels stored at 25 ± 2 °C and 45 ± 2 °C, protected from light (*Significance at $P < 0.05$ when compared with the content at initial time)	49
4.15 Color of the antioxidant active extract of pomegranate fruit peels stored under protected from humidity conditions (A) and stored under 75% relative humidity (B)	51

LIST OF FIGURES (Continued)

Figure	Page
4.16 Ellagic acid content in the antioxidant active extract of pomegranate fruit peels stored under 75% relative humidity and protected from humidity conditions, at 25 ± 2 °C (*Significance at $P < 0.05$ when compared with the content at initial time)	52
4.17 Antioxidant activity evaluated by DPPH assay of the antioxidant active extract of pomegranate fruit peels stored 75%RH and protected from humidity conditions, at 25 ± 2 °C (*Significance at $P < 0.05$ when compared with the content at initial time)	54
4.18 Antioxidant activity evaluated by β -carotene bleaching assay of the antioxidant active extract of pomegranate fruit peels stored 75%RH and protected from humidity conditions, at 25 ± 2 °C	55
4.19 HPLC-chromatogram of the extracts at initial time (A) and after stability test (B)	56
4.20 Hydrolysis of ellagic acid	57
4.21 Physical appearances of cream bases before (A) and after (B) heating and cooling test (8 cycles): Rx1 (1), Rx 2 (2), Rx 3 (3) Rx 4 (4), Rx 5 (5)	59
4.22 Pomegranate extract creams, Rx 5-1 (1), Rx 5-2 (2) before (A) and after (B) heating and cooling test	61
4.23 HPLC-chromatogram of the preparation extracts at initial time (A) and after heating-cooling test (B)	62