CONTENT

| | Page |
|---|------|
| Contents | |
| (ix) | |
| List of Tables | |
| (xii) | |
| List of Figures | |
| (xiv) | |
| 1. Introduction | 1 |
| Literature Review | 3 |
| 1. Edible films and biodegradable films | 3 |
| 2. Protein films | 4 |
| 3. Protein film formation and properties | 5 |
| 4. Effect of pH on the properties of protein-based film | 9 |
| 5. Effect of plasticizers on the properties of protein-based film | 12 |
| 6. Effect of lipids on the properties of protein-based film | 15 |
| 7. Protein-based film from different sources | 17 |
| 8. Improvement of protein films properties | 21 |
| 9. Stability of protein-based films during storage | 26 |
| 10.Applications of protein based-films | 27 |
| Objectives | 30 |
| 2. Material and methods | 31 |
| 1. Materials | 31 |

| 2. Instruments | 32 |
|--------------------------------|----|
| 3. Chemical analyses of surimi | 32 |
| 4. Preparation of surimi film | 33 |

CONTENT (CONTINUED)

| 5. Determination of film properties | 33 |
|---|----|
| 6. Effects of pH on the compositional changes of film-forming solution | 36 |
| 7. Effect of lipids on the film properties | 36 |
| 8. Effect of crosslinker on surimi film properties | 37 |
| 9. Characterization of surimi film | 38 |
| 10.Changes of films during storage | 40 |
| 11.Uses of surimi film to extend the shelf-life of dried fish powder | 40 |
| 12.Statistical analysis | 41 |
| 3. Results and Discussion | 42 |
| 1. Chemical composition of surimi from tropical fish | 42 |
| 2. Effect of pH on the properties of film from surimi | 44 |
| 3. Compositional changes of surimi film-forming solution as affected | 55 |
| by acidic and alkaline pHs | |
| 4. Effect of lipid types and concentrations on the properties of bigeye | 60 |
| snapper surimi film | |

Page

| 5. Effect of crosslinker on the properties of bigeye snapper surimi film | 76 |
|--|-----|
| 6. Characterization of bigeye snapper surimi film | 93 |
| 7. Changes of films during storage | 104 |

CONTENT (CONTINUED)

| | Page |
|---|------|
| 8. Uses of bigeye snapper surimi film to extend the shelf-life of dried | 110 |
| fish powder | |
| 4. Conclusion | 116 |
| References | 118 |
| Appendix | 138 |
| Vitae | 151 |

LIST OF TABLES

| Т | Table | |
|----|--|------|
| 1 | Water vapor permeability of various protein films. | 7 |
| 2 | Tensile strength and elongation at break of various films. | 8 |
| 3 | Comparison of specific mechanical and barrier properties of plastic wrap and | 18 |
| | spray-dried (SD) and flash-dried (FD) wheat gluten films. | |
| 4 | Applications of protein based-films in foods. | 28 |
| 5 | Proximate compositions of surimi from different tropical fish. | 43 |
| 6 | Physical and mechanical properties of surimi film. | 46 |
| 7 | L*, a* and b*-values of surimi film. | 50 |
| 8 | Film solubility and protein solubility of surimi film. | 54 |
| 9 | Tensile strength (TS), elongation at break (EAB), water vapor permeability | 63 |
| | (WVP) and thickness of bigeye snapper surimi film as affected by types | |
| | and amounts of lipids. | |
| 1(| O Color parameters and transparency of bigeye snapper surimi film as affected | 67 |
| | by types and amounts of lipids. | |
| 11 | l Film solubility and protein solubility of bigeye snapper surimi film as affected | d 69 |
| | by types and amounts of lipids. | |
| 12 | 2 Tensile strength (TS), elongation at break (EAB), water vapor permeability | 79 |

(xii)

(WVP) and thickness of bigeye snapper surimi film incorporated with palm oil as affected by types and amounts of aldehydes.

13 Color parameters and transparency of bigeye snapper surimi film incorporated81 with palm oil as affected by types and amounts of aldehydes.

LIST OF TABLES (CONTINUED)

| Table | Page |
|--|------|
| 14 Film solubility and protein solubility of bigeye snapper surimi film | 83 |
| incorporated with palm oil as affected by types and amounts of aldehydes. | |
| 15 Tensile strength (TS), elongation at break (EAB), water vapor permeability | 88 |
| (WVP) and thickness of bigeye snapper surimi film incorporated with palm | |
| oil as affected by MTGase. | |
| 16 Color parameters and transparency of bigeye snapper surimi film incorporated | l 89 |
| with palm oil as affected by MTGase levels. | |
| 17 Film solubility and protein solubility of bigeye snapper surimi film | 90 |
| incorporated with palm oil as affected by MTGase levels. | |
| 18 Degree of hydrolysis (%) of bigeye snapper surimi films incorporated with oil | 96 |
| and/or formaldehyde. | |
| 19 Protein solubility (%) of bigeye snapper surimi films in various solvents. | 98 |
| 20 Glass transition temperature (T_g) of the control films, films incorporated | 103 |
| with palm oil at 75 % glycerol substitution, films incorporated with palm oil | |
| at 75 % glycerol and 5 mM formaldehyde. | |
| 21 Degradation temperature (T_d) of the control films, films incorporated with | 104 |
| palm oil at 75 % glycerol substitution, films incorporated with palm oil | |

(xiii)

at 75 % glycerol and 5 mM formaldehyde and surimi (unplasticized, without cryoprotactant).

LIST OF FIGURES

| Figure | | Page |
|--------|---|------|
| 1 | Mechanism of film formation. | 6 |
| 2 | Mechanism of protein cross-linking by formaldehyde, glyoxal or | 22 |
| | glutaraldehyde. | |
| 3 | Transglutaminase-catalyzed cross-linking reaction in proteins. | 25 |
| 4 | Protein patterns of surimi from different tropical fish. | 44 |
| 5 | Protein patterns of surimi film from bigeye snapper, goat fish and | 52 |
| | threadfin bream. | |
| 6 | Reducing sugar content of acidic and alkaline film-forming solutions | 56 |
| | of surimi from different tropical fish with different exposure times. | |
| 7 | Protein patterns of acidic and alkaline film-forming solutions from | 59 |
| | bigeye snapper surimi, goat fish surimi and threadfin bream surimi | |
| | with different exposure times. | |
| 8 | Protein patterns of bigeye snapper surimi films incorporated with | 70 |
| | palm oil, butter and shortening. | |
| 9 | SEM micrographs of bigeye snapper surimi films incorporated | 73 |
| | without and with palm oils at different levels. | |

10 SEM micrographs of bigeye snapper surimi films incorporated with 74 palm oil, butter and shortening at 75% glycerol substitution.

LIST OF FIGURES (CONTINUED)

| Figure | Page |
|---|------|
| 11 Moisture sorption isotherms of bigeye snapper surimi film added | 76 |
| with palm oil at 75% glycerol substitution at 4°C and room temperature. | |
| 12 Protein patterns of bigeye snapper surimi films containing palm oil | 85 |
| and incorporated with formaldehyde, glutaraldehyde and glyoxal. | |
| 13 Protein patterns of bigeye snapper surimi films containing palm oil | 91 |
| and added with MTGase at different levels. | |
| 14 Moisture sorption isotherms of the film incorporated with palm oil and | 92 |
| added with 5 mM formaldehyde at 4°C and room temperature. | |
| 15 SEM micrographs of bigeye snapper surimi films. Control film (without | 94 |
| lipid and formaldehyde), film incorporated with palm oil at 75% glycerol | |
| substitution and film incorporated with palm oil at 75% glycerol | |
| substitution and 5 mM formaldehyde. | |
| 16 First scan of DSC thermogram of the control film, film incorporated | 101 |

(xv)

with palm oil at 75 % glycerol substitution and film incorporated with palm oil at 75 % glycerol and 5 mM formaldehyde.

17 Second scan of DSC thermogram of the control film, film incorporated
101
with palm oil at 75 % glycerol substitution and film incorporated with palm
oil at 75 % glycerol and 5 mM formaldehyde.

LIST OF FIGURES (CONTINUED)

| Figure | Page |
|--|-------------|
| 18 TGA thermogram of the control film, film incorporated with palm oil | 102 |
| at 75 % glycerol substitution and film incorporated with palm oil at 75 $\%$ | ó |
| glycerol and 5 mM formaldehyde. | |
| 19 Changes in tensile strength and elongation at break of bigeye snapper sur | imi 106 |
| films containing palm at 75% glycerol substitution and 5 mM formaldehy | <i>y</i> de |
| during storage at different temperatures and RHs. | |
| 20 Changes in water vapor permeability of bigeye snapper surimi films | 107 |
| containing palm at 75% glycerol substitution and 5 mM formaldehyde | |
| during storage at different temperatures and RHs. | |
| 21 Changes in film solubility of bigeye snapper surimi films containing | 107 |
| palm at 75% glycerol substitution and 5 mM formaldehyde during | |
| storage at different temperatures and RHs. | |
| 22 Changes in color of bigeye snapper surimi films containing palm at 75% | 109 |
| glycerol substitution and 5 mM formaldehyde during storage at different | |
| temperatures and RHs. | |
| 23 Changes in transparency of bigeye snapper surimi films containing palm | 110 |

at 75% glycerol substitution and 5 mM formaldehyde during storage at different temperatures and RHs.

LIST OF FIGURES (CONTINUED)

Figure

Page

- 24 Changes in moisture content of dried fish powder packaged using bigeye 111
 snapper surimi films incorporated with palm oil at 75% glycerol substitution
 (SO), bigeye snapper surimi film incorporated with palm oil at 75% glycerol substitution and 5 mM formaldehyde (SOF) or polyethylene film (PE) during storage at 30-32°C.
- 25 Changes in color of dried fish powder packaged using bigeye snapper surimi 113 films incorporated with palm oil at 75% glycerol substitution (SO), bigeye snapper surimi film incorporated with palm oil at 75% glycerol substitution and 5 mM formaldehyde (SOF) or polyethylene film (PE) during storage at 30-32°C.
- 26 Changes in TBARS of dried fish powder packaged using bigeye snapper 115
 surimi films incorporated with palm oil at 75% glycerol substitution (SO),
 bigeye snapper surimi film incorporated with palm oil at 75% glycerol
 substitution and 5 mM formaldehyde (SOF) or polyethylene film (PE)
 during storage at 30-32°C.