

References

- Abe, Y. 1994. Quality of kamaboko gel prepared from walleye pollock surimi with an additive containing transglutaminase. *Nippon suisan Gakkaishi*. 60: 381-387.
- Akahane, T., Chihara, S., Yoshida, Y., Tsuchiya, T., Nogchi, S., Ookami, H. and Matsumoto, J.J. 1984. Roles of constituent proteins in gel properties of cooked meat gels. *Bull. Jap. Soc. Sci. Fish.* 50: 1029-1033.
- Akahane, Y. and Shimizu, Y. 1990. Effect of setting incubation on the water-holding capacity of salt-ground fish meat and its heated gel. *Nippon Suisan Gakkaishi*. 56: 139-143.
- Almas, K.A. 1981. Chemistry and microbiology of fish proceeding. Dept. of Biochemistry, Norwegian Inst. of Technol. Univ. of Trondheim. Norway.
- Alvarez, C., Couso, I. and Tejada, M. 1995. Sardine surimi gels as affected by salt concentration, blending heat treatment and moisture. *J. Food Sci.* 60:622-626.
- Alvarez, C. and Tejada, M. 1997. Influence of texture of suwari gels on kamaboko gels made from sardine (*Sardina pilchardus*) surimi. *J. Sci. Food Agric.* 75: 472-480.
- An, H., Peters, M.Y. and Seymour, T.A. 1996. Role of endogenous enzymes in surimi gelation. *Trends Food Sci. Technol.* 7: 320-327.
- Ando, H., Adachi, M., Umead, K., Matsuura, A., Nonaka, M., Uchio, R., Tanaka, H. and Motoki, M. 1989. Purification and characteristics of novel transglutaminase derived from microorganisms. *Agric. Biol. Chem.* 53: 2613-2617.

- AOAC. 1999. Official Method of Analysis, 16th ed. Association of Official Analytical Chemists, Washiton, DC, USA.
- Arai, K., Kawamura, K. and Hayashi, C. 1973. The relative thermostability of actomyosin ATPase from the dorsal muscle of various fish species. Bull. Jap. Soc. Sci. Fish. 39: 1077-1082.
- Araki, H. and Seki, N. 1993. Comparison of reactivity of transglutaminase to various fish actomyosin. Nippon Suisan Gakkaishi. 80: 711-716.
- Asagami, T., Ogiwara, M., Wakameda, A. and Noguchi, S. 1995. Effect of microbial transglutaminase on the quality of frozen surimi made from various kinds of fish species. Fish. Sci. 61: 267-272.
- Ashie, I. N. A. and Lanier, T. C. 2000. Transglutaminase in seafood processing. *In* Seafood enzymes utilization and influence on posthaviest seafood quality. (Haard, N.F. and Simpson, B.K., eds.) p. 147-166. Marcel Dekker. New York.
- Azuma, Y. and Konno, K. 1998. Freeze denaturation of carp myofibril compared with thermal denaturation. Fish. Sci. 64: 287-290.
- Benjakul, S. and Bauer, F. 2000. Physicochemical and enzymatic change of cod muscle protein subjected to freeze-thaw cycle. J. Sci. Food Agric. 80: 1143-1150.
- Benjakul, S., Chantarasuwan, C. and Visessanguan, W. 2003. Effect of medium temperature setting on gelling characteristics of surimi from some tropical fish. Food Chem. 82: 567-574.

- Benjakul, S., Seymour, T.A., Morrissey, M.T. and An, H. 1997. Physicochemical changes in Pacific whiting muscle proteins during ice storage. *J. Food Sci.* 62: 729-733.
- Benjakul, S. and Visessanguan, W. 2003. Transglutaminase-mediated setting in bigeye snapper surimi. *Food Res. Int.* 36: 253-266.
- Benjakul, S., Visessanguan, W. and Chantarasuwan, C. 2004a. Effect of high-temperature setting on gelling characteristic of surimi from some tropical fish. *Int. J. Food Sci. Technol.* 39: 671-680.
- Benjakul, S., Visessanguan, W., Ishizaki, S. and Tanaka, M. 2001. Differences in gelation characteristics of natural actomyosin from two species of bigeye snapper, *Priacanthus tayenus* and *Priacanthus macracanthus*. *J. Food Sci.* 66: 1311-1318.
- Benjakul, S., Visessanguan, W. and Kwalumtharn, Y. 2004b. The effect of whitening agents on the gel-forming ability and whiteness of suwari. *Int. J. Food Sci. Technol.* 39: 773-781.
- Benjakul, S., Visessanguan, W. and Pecharat, S. 2004c. Suwari gel properties as affected by transglutaminase activator and inhibitors. *Food Chem.* 85: 91-99.
- Benjakul, S., Visessanguan, W., Riebroy, S., Ishizaki, S. and Tanaka, M. 2002. Gel-forming properties of surimi produced from bigeye snapper, *Priacanthus tayenus* and *P. macracanthus*, stored in ice. *J. Sci. Food Agric.* 82: 1442-1451.
- Benjakul, S., Visessanguan, W. and Srivilai, C. 2001a. Porcine plasma proteins as gel enhancer in bigeye snapper (*Priacanthus tayenus*) surimi. *J. Food Biochem.* 25: 285-305.

- Benjakul, S., Visessanguan, W. and Srivilai, C. 2001b. Gel properties of bigeye snapper (*Priacanthus tayenus*) surimi as affected by setting and porcine plasma proteins. *J. Food Qual.* 24: 453-471.
- Benjakul, S., Visessanguan, W. and Tueksuban, J. 2003b. Heat-activated proteolysis in lizardfish (*Saurida tumbil*) muscle. *Food Res. Int.* 80: 535–544.
- Benjakul, S., Visessanguan, W. and Tueksuban, J. 2003c. Changes in physicochemical properties and gel-forming ability of lizardfish (*Saurida tumbil*) during post-mortem storage in ice. *Food Chem.* 80: 535-544.
- Chan, J.K., Gill, T.A., and Paulson, A.T. 1992. Cross-linking of myosin heavy chains from cod, herring and silver hake during thermal setting. *J. Food Sci.* 57: 906-912.
- Chang, C.C. and Regenstein, J.M. 1997a. Water uptake, protein solubility, and protein changes of cod mince stored on ice as affected by polyphosphates. *J. Food Sci.* 62: 305-309.
- Chang, C.C. and Regenstein, J.M. 1997b. Textural changes and functional properties of cod mince proteins as affected by kidney tissue and cryoprotectants. *J. Food Sci.* 62: 299-304.
- Chen, H.H. 1995. Thermal stability and gel-forming ability of shark muscle as related to ionic strength. *J. Food Sci.* 60: 1237-1240.
- Dziezak, J.D. 1990. Phosphates improve many food. *Food Technol.* 44(4): 80-92.
- Ebasi, S., Kodama, A. and Ebashi, F. 1968. Troponin I. Preparation and physiological function. *J. Biochem.* 64: 465-477.
- Ellinger, R.H. 1972a. Nomenclature, classification and structure of phosphates used in foods. *In Phosphates as Food Ingredients.* p.3. CRC Press. Boca Raton. Fla.

- Ellinger, R.H. 1972b. The functions and applications of phosphates in food systems. *In Phosphates as Food Ingredients*. p.31. CRC Press. Boca Raton. Fla.
- Ellinger, R.H. 1975. Phosphates in food processing. *In Hand book of food additives*. (Furia, T.E., ed.) p. 645-698. Cleveland CRC Press. California. USA.
- Ellinger, R.H. 1977. Phosphates in food processing. *In CRC Handbook of Food Additives*. (Furia, T.E., ed.) p. 671. CRC Press, Inc., Boca Raton. Fla.
- Fiske, C.H. and Subbarow, Y. 1925. The colorimetric determination of phosphorus. *J. Biol. Chem.* 66: 375-400.
- Foegeding, E., Lanier, T.C. and Hultin, H.O. 1996. Characteristics of edible plant tissues. *In Food Chemistry*. (Fennema, O.R., ed.) p. 94301012. Marcel Dekker. New York.
- Folk, J.E. 1970. Transglutaminase. *Methods Enzymol.* 17: 889-894.
- Folk, J. E. 1980. Transglutaminases. *Annual Rev. Biochem.* 49: 517–531.
- Folk, J. E. 1983. Mechanism and basis for specificity of transglutaminase-catylyzed α -(γ -glutamyl)lysine bond formation. *Adv. Enzymol.* 54: 9-35.
- Folk, J.E., and Chung, S.I. 1973. Molecular and catalytic properties of transglutaminase. *Adv. Enzymol.* 38: 109-191.
- Gerber, U., Jucknischke, U., Putzien. S. and Fuchsbauer, H. L. 1994. A rapid and simple method for the purification of transglutaminase from *Streptovercillium mobaraense*. *J. Biochem.* 299: 825-829.
- Gill, T.A., Chan, J.K., Phonchareon, K.F. and Paulson, A.T. 1992. Effect of salt concentration and temperature heat-induced aggregation and gelation of fish myosin. *Food Res. Int.* 25: 333-341.

- Gill, T.A. and Conway, J.T. 1989. Thermal aggregation of cod (*Gadus morhua*) muscle proteins using lethyl-3-(3-dimethylaminopropyl) carbodiimide as a zero length crosslinker. *Agric. Biol. Chem.* 53: 2553-2562.
- Grabowska, J. and Sikorski, Z.E. 1976. The gel-forming capacity of fish myofibrillar proteins. *Lebensm. Wiss. U. Technol.* 9: 33-37.
- Grantham, G.J. 1981. Minced fish technology: A Review (FAO Fisheries Technical Paper No 216). UK: Melton Mowbray, Leics.
- Greene, D.H. and Babbitt, J. 1990. Control of muscle softening and protease parasite interactions in arrowtooth flounder (*Atheresthes stomias*). *J. Food Sci.* 55: 579-580.
- Hamm, R. 1971. Interactions between phosphates and meat proteins. *In Symposium: Phosphates in Food Processing.* (Deman, J.M. and Melnychyn, P., eds.) p. 65-82. The AVI Publishing Co., Westport. CT.
- Ho, M.L., Leu, S.Z., Hsieh, J.F. and Jiang, S.T. 2000. Technical approach to simplify the purification method and characterization of microbial transglutaminase produced from *Streptoverticillium ladakanum*. *J. Food Sci.* 65: 76-80.
- Hossain, S.M.Z., Ito, T., Kanoh, S. and Niwa, E. 1998. Incorporation of dansyl glutamine into muscle protein during incubation of fish flesh sol at 30°C. *Fish. Sci.* 64: 95-98.
- Huang, Y. P., Seguro, K., Motoki, M. and Tawada, K. 1992. Cross-linking of contractile proteins from skeletal muscle by treatment with microbial transglutaminase. *J. Biochem.* 112: 229-234.
- Huidobro, A. and Tejada, M. 1994. Emulsifying capacity of fish mince from several species during frozen storage. *J. Sci. Food Agric.* 61:333-338.

- Hultin, H.O. 1985. Characteristics of muscle tissue. *In* Food Chemistry, 2nd ed (Fennema, O.R., ed.) p.725. Marcel Dekker. New York.
- Hwang, G.C., Ushio, H., Watabe, S., Iwamoto, M. and Hashimoto, K. 1991. The effect of thermal acclimation on rigor mortis progress of carp stored at different temperatures. *Nippon Suisan Gakkaishi*. 57: 541-548.
- Ishikawa, S. 1978. Fish jelly product (kamaboko) and frozen minced meat (frozen surimi) made of sardine-II. Influence of the temperature during the manufacturing process on the jelly strength of kamaboko. *Bull. Tokai Reg. Fish Res. Lab.* 94: 37-44.
- Itawa, K., Kanna, K. and Okada, M. 1977. Kamaboko formation from mackerel and red sea bream myosins. *Bull. Jap. Soc. Sci. Fish.* 43: 237-240.
- Jiang, S.T., Leu, S.Z. and Tsai, G.J. 1998a. Cross-linking of mackerel surimi actomyosin by microbial transglutaminase and ultraviolet irradiation. *J. Agric. Food Chem.* 46: 5278-5282.
- Jiang, S.H., Ho, M.L., Jiang, S.H., Lo, L. and Chen, H.C. 1998b. Color and quality of mackerel surimi as affected by alkaline washing and ozonation. *J. Food Sci.* 63: 652-655.
- Jiang, S.T., Wang, F.J. and Chen, C.S. 1989. Properties of actin and stability of the actomyosin reconstituted from milkfish (*Chanos chanos*) actin and myosin. *J. Agric. Food Chem.* 37: 1232-1235.
- Jiang, S.T., Hsieh, J.F., Ho, M.L. and Chung, Y.C. 2000a. Microbial transglutaminase affects gel properties of golden threadfin-bream and pollack surimi. *J. Food Sci.* 65: 694-699.

- Jiang, S.T., Hsieh, J.F., Ho, M.L. and Chung, Y.C. 2000b. Combination effects of microbial transglutaminase, reducing agent, and protease inhibitor on the quality of hairtail surimi. *J. Food Sci.* 65: 241-245.
- Kamath, G., Lanier, T.C., Foegeding, E.A. and Hamann, D.D. 1992. Nondisulfide covalent cross-linking of myosin heavy chain in “setting” of Alaska pollock and Atlantic croaker surimi. *J. Food Biochem.* 16: 151-172.
- Kang, T.S. and Lanier, T.C. 2000. Heat-induced softening of surimi gels by proteinases. *In* Surimi and surimi seafood. (Park, J.W., ed.) p. 445-474. Marcel Dekker. New York.
- Kato, T., Tomimatsu, K. and Toba, S. 1991. Manufacture of storage-stable retort mapuo-doufu. *Jap. Kokai Tokkyo Koho JP.* 03168059.
- Kim, C.Y., Choi, Y.J. and Pyeun, J.H. 1982. Changes occurred in protein and amino acid composition during postmortem aging of white and dark muscle of yellowtail at 2°C. *Bull. Jap. Soc. Sci. Fish.* 15: 123-127.
- Kimura, I.M., Sugimoto, M., Toyoda, K., Seki, N., Arai, K. and Fujita, T. 1991. A study on the cross-links reaction of myosin in kamaboko “Suwari” gels. *Nippon Suisan Gakkaishi.* 57: 1386-1396.
- Kishi, H., Nozawa, H. and Seki, N. 1991. Reactivity of muscle transglutaminase on carp myofibrils and myosin B. *Nippon Suisan Gakkaishi.* 57: 1203-1210.
- Kudo, G., Okada, M. and Mivauchi, D. 1973. Gel-forming capacity of washed and unwashed flesh of some Pacific coast species of fish. *Mar. Fish. Rev.* 35: 10-13.
- Kumazawa, Y., Numazawa, T., Seguro, K. and Motoki, M. 1995. Suppression of surimi gel setting by transglutaminase inhibitors. *J. Food Sci.* 60: 715–717.

- Kumazawa, Y., Seguro, K., Takamura, M. and Motoki, M. 1993a. Participation of transglutaminase in the manufacturing process of “kamaboko” Paper no. 124A, presented at 53rd Annual Meeting of the Institute of Food Technologists, Chicago, IL, July, 10-14.
- Lanier, T.C. 2000. Gelation chemistry. *In* Surimi and surimi sea food. (Park, J.W., ed) p. 237-265. Marcel Dekker. New York.
- Laemmli, U.K. 1970. Cleavage of structural protein during the assembly of the head bacteriophage T4. *Nature*. 227: 680-685.
- Lee, C.M. 1984. Surimi process technology. *Food Technol.* 38 (11): 69-80.
- Lee, C.M. 1986. Surimi manufacturing and fabrication of surimi-based products. *Food Technol.* 40(30): 115-124.
- Lee, C.M. 1990. Countercurrent and continuous washing systems. *In* Engineered Seafood Including Surimi. (Martin, R.E. and Collette, R.L., eds.). Park Ridge, NJ: Noyes Data Corp.
- Lee, H.G., Lanier, T.C., Hamann, D.D. and Knopp, J.A. 1997. Transglutaminase effects on low temperature gelation of fish protein sols. *J. Food Sci.* 62: 20-24.
- Lee, N.G. and Park, J.W. 1998. Calcium compounds to improve gel functionality of Pacific whiting and Alaska pollock surimi. *J. Food Sci.* 63: 969-974.
- Lee, N.H., Seki, N., Kato, N., Nakagawa, S., Terui, S. and Arai, K. 1990. Gel forming ability and cross-linking ability of myosin heavy chain in salted meat paste from threadfin bream. *Nippon Suisan Gakkaishi*. 56: 329-336.
- Lewis, D.F., Groves, K.H.M., and Holgate, J.H. 1986. Action of polyphosphates in meat products. *Food Microstructure*. 5: 53-62.

- Lin, T.M. and Park, J.W. 1997. Effective washing conditions reduce water usage for surimi processing. *J. Aquat. Food Prod. Technol.* 6(2): 65-79.
- Liu, Y.M., Lin, T.S. and Lanier, T.C. 1982. Thermal denaturation and aggregation of actomyosin from Atlantic croaker. *J. Food Sci.* 47: 1916-1924.
- Mackie, I.M. 1994. Fish protein. *In* New and developing sources of food proteins (Hubson, B.J.G., ed.) p. 95-137. Chapman & Hall. New York.
- Makinodan, Y., Hirotsuka, M. and Ikeda, S. 1980. Autolysis of carp muscle. *Bull. Jap. Soc. Sci. Fish.* 46: 1507-1511.
- Matsukawa, M., Hirata, F., Kimura, S. and Arai, K. 1995. Effect of sodium pyrophosphate on gelling property and cross-linking of myosin heavy chain in setting-heating gel from six transglutaminase from ordinary muscles of various fishes and marine invertebrates. *Comp. Biochem. Physiol.* 118B: 313-317.
- Matsumoto, J.J. 1979. Denaturation of fish muscle proteins during frozen storage. *In* Protein at low temperatures. (Fennema, O., ed.) p. 206. ACS. Adv. Chem. Series, Washington. DC. USA.
- Matsumoto, J.J. 1980. Chemical deterioration of muscle proteins during frozen storage. *In* Chemical deterioration of protein (Whitaker, J.K. and Fujimaki, J.R. eds.) p. 95. ACS Symp. Series 123. Washington, DC.
- McCormick, R.T. 1994. Structure and properties of tissues. *In* Muscle food. (Kinsman, D.M., Kotula, A.W. and Breidenstein, B.C., eds.) p. 25-62. Chapman & Hall. New York.
- Montero, P. Lopez-Caballero, M.E., Perez-Mateos, M., Solas, M.T. and Gomez-Guillen, M.C. 2005. Transglutaminase activity in pressure-induced gelation assisted by prior setting. *Food Chem.* 90: 751-758.

- Muguruma, M., Sakamoto, K., Numata, M., Yamada, H. and Nakamura, T. 1990. Studies on application of transglutaminase to meat and meat products, II. Effect of microbial transglutaminase on gelation of myosin B, myosin and actin. *Nippon Shokuhin Kogyo Gakkaishi*. 37: 446-453.
- Nakahara, C., Nozawa, H. and Seki, N. 1999. A comparison of cross-linking of fish myofibrillar proteins by endogenous and microbial transglutaminases. *Fish. Sci.* 65: 138-144.
- Ni, S., Nozawa, H. and Seki, N. 1998. Effect of microbial transglutaminase on thermal gelation of carp actomyosin sol. *Fish. Sci.* 64: 434-438.
- Nielsen, R.G. and Pigott, G.M. 1994. Gel strength increased in low-grade heat-set surimi with blended phosphates. *J. Food Sci.* 59: 246-250.
- Ng, C.S. 1987. Measurement of free and expressible drips. *In* Laboratory manual on analytical methods and procedure for fish and fish products. (Hasegawa, H. ed.) p. 1-2. Southeast Asian Fisheries Development Center. Singapore.
- Niwa, E. 1992. Chemistry of surimi gelation. *In* Surimi Technology. (Lanier, T.C. and Lee, C.M., eds.) p. 389-428. Marcel Dekker. New York.
- Noguchi, S. and Matsumoto, J.J. 1971. Studies on the control of the denaturation of the fish muscle proteins during frozen storage-II. Preventive effect of amino acids and related compounds. *Bull. Jap. Soc. Sci. Fish.* 37: 1115-1121.
- Nonaka, M., Tanaka, H., Okiyama, A., Motoki, M., Anso, H., Umeda, K. and Matsukuwa, A. 1989. Polymerization of several proteins by Ca^{2+} -independent transglutaminase derived from microorganisms. *Agric. Biol. Chem.* 36: 1078-1081.

- Nowsad, A.A.K.M., Kanoh, S. and Niwa, E. 1994a. Setting of surimi paste in which transglutaminase is inactivated by *N*-ethylmaleimide. *Fish. Sci.* 60: 189-191.
- Nowsad, A.A.K.M., Kanoh, S. and Niwa, E. 1994b. Setting of transglutaminase free actomyosin paste prepared from Alaska pollack surimi. *Fish. Sci.* 60: 295-297.
- Nowsad, A.A.K.M., Kanoh, S. and Niwa, E. 1995. Contribution of transglutaminase on the setting of various actomyosin pastes. *Fish. Sci.* 61: 79-81.
- Nozawa, H., Mamagoshi, S. and Seki, N. 1997. Partial purification and characterization of walleye pollack surimi. *Nippon Suisan Gakkaishi.* 62: 94-103.
- Ochiai, Y. and Chow, C.J. 2000. Myosin ATPase. *In* Seafood enzymes. (Haard, N.F. and Simpson, B.K., eds.) p. 69-90. Marcel Dekker. New York.
- Offer, G. and Trinick, J. 1983. A unifying hypothesis for the mechanism of changes in the water-holding capacity of meat. *J. Sci. Food Agric.* 34: 1018-1025.
- Ouali, A. and Valin, C. 1981. Effect of muscle lysosomal enzymes and calcium activated neutral proteinase on myofibrillar ATPase activity: relationship with aging changes. *Meat Sci.* 5: 233-245.
- Park, J.W. 1994. Functional protein additives in surimi gels. *J. Food Sci.* 59: 525-527.
- Park, J.W. 2000. Ingredient technology and formulation development. *In* Surimi and Surimi Seafood. (Park, J.W. ed.) p. 343-329. Marcel Dekker. New York. USA.
- Park, J.W. and Lanier, T.C. 1987. Combined effects of phosphates and sugar or polyol on protein stabilization of fish myofibrils. *J. Food Sci.* 52: 1509-1513.

- Park, J.W. and Lanier, T.C. 1989. Scanning calorimetric behavior of tilapia myosin and actin due to processing of muscle and protein denaturation. *J. Food Sci.* 54: 49-51.
- Park, J.W. and Morrissey, M.T. 1994. The need for developing surimi standard. *In* Quality Assurance and Quality Control for Seafood. (Sylvia, G. and Morrissey, M.T. eds). Corvallis, OR: Oregon Sea Grant. USA.
- Park, J.W. and Morrissey, M.T. 2000. Manufacturing of surimi from light muscle fish. *In* Surimi and Surimi Seafood. (Park, J.W., ed.) p. 23-58. Marcel Dekker. New York.
- Pearsons, A.M. and Young, R.B. 1989. Muscle and meat biochemistry. Academic Press, Inc., San Diego. CA.
- Pomeranz, Y. 1991. Proteins: General. *In* Functional properties of food components. (Pomeranz, Y., ed.). p. 147-189. Academic Press, Inc., San diego. CA.
- Ramirez, J.A., Martin-Polo, J.O. and Bandman, E. 2000. Fish myosin aggregation as affected by freezing and initial physical state. *J. Food Sci.* 65: 556-559.
- Ramirez, J., Uresti, R., Tellez, S. and Vazquez, M. 2002. Using salt and microbial transglutaminase as binding agents in restructured fish products resembling hams. *J. Food Sci.* 67: 1778-1784.
- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T.C. 2005. Combination effects of chicken plasma protein and setting phenomenon on gel properties and cross-linking of bigeye snapper muscle proteins. *Lebensmittel-Wissenschaft und-Technologie.* 38: 353-362.

- Robinson, H.W. and Hodgen, C.G. 1940. The biuret reaction in the determination of serum protein. I. A study of condition necessary for the production of the stable color which bears a quantitative relationship to the protein concentration. *J. Biol. Chem.* 135: 707-725.
- Roussel, H. and Cheftel, J.C. 1990. Mechanism of gelation of sardine protein: influence of thermal processing and various additives on the texture and protein solubility and kamaboko gel. *Int. J. Food Sci. Technol.* 25: 260-280.
- Saeki, H., Iseya, Z., Sugiura, S. and Seki, N. 1995. Gel forming characteristics of frozen surimi from chum salmon in the presence of protease inhibitors. *J. Food Sci.* 60: 917-928.
- Sakamoto, H., Kumazawa, Y., Toiguchi, S., Seguro, K., Soeda, T. and Motoki, M. 1995a. Gel strength enhancement by addition of microbial transglutaminase during onshore surimi manufacture. *J. Food Sci.* 60: 300-309.
- Sakamoto, S., Kumazawa, Y., Kawajiri, H. and Motoki, M. 1995. ϵ -(γ -glutamyl) lysine crosslink distribution in foods as determined by improved method. *J. Food Sci.* 60: 416-419.
- Sano, T., Noguchi, S.F., Tsuchiya, T. and Matsumoto, J.J. 1989. Role of F-actin in thermal gelation of fish actomyosin. *J. Food. Sci.* 54: 800-811.
- Sano, T., Noguchi, S.F., Marsumoto, J.J. and Tsuchiya, T. 1990. Thermal gelation characteristics of myosin subfragments. *J. Food Sci.* 55: 55-58.
- Sano, T., Ohno, T., Fuchino, H.O., Matsumoto, J. and Tsuchiya, T. 1994. Carp natural actomyosin: thermal denaturation mechanism. *J. Food Sci.* 59: 1002-1008.
- Seguro, K., Kumazawa, Y., Ohtsuka, T., Toiguchi, S. and Motoki, M. 1995. Microbial transglutaminase and ϵ -(γ -glutamyl) lysine cross-link effects on elastic properties of kamaboko gels. *J. Food Sci.* 60: 305-311.

- Seki, N., Uno, H., Lee, N.H., Kimura, I., Toyoda, K., Fujita, T. and Arai, K. 1990. Transglutaminase activity in Alaska pollack muscle and surimi and its reaction with myosin B. *Nippon Suisan Gakkaishi*. 56: 125–132.
- Seki, N., Nakahara, C., Takeda, H., Maruyama, N. and Nozawa, H. 1998. Dimerization site of carp myosin heavy chains by the endogenous transglutaminase. *Fish. Sci.* 64: 314-319.
- Semejima, K., Ishioroshi, M. and Yasui, T. 1981. Relative roles of the head and tail portions of the molecule in heat induced gelation of myosin. *J. Food Sci.* 46: 1412-1416.
- Shimizu, Y. 1985. Biochemical and functional properties of material fish. *In* Proceedings of the International Symposium on Engineered Seafood Including Surimi. (Seattle, W.A., Martin, R.E. and Collete, R.L. eds.) Nov 19-21. p. 148-167. National Fish. Institute. Washington, DC.
- Shimizu, Y., Nishioka, F., Machida, R. and Shien, C.M. 1983. Gelation characteristics of salt added myosin sol. *Bull. Jap. Soc. Sci. Fish.* 49: 1239-1240.
- Shults, G.W., Russell, D.R. and Wierbicki, E. 1972. Effect of condensed phosphates on pH, swelling and water-holding capacity of beef. *J. Food Sci.* 37: 860-864.
- Sikorski, Z.E., Kolakowska, A. and Pan, B.S. 1990. The nutritive composition of the major groups of marine food organisms. *In* Seafood: Resources, Nutritional Composition and Preservation. (Sikorski, Z.E. ed.) p. 29-54, CRC Press, Boca Raton. Florida.
- Simpson, R., Kolbe, E., MacDonald, G., Lanier, T. and Morrissey, M.T. 1994. Surimi production from partially processed and frozen Pacific whiting (*Merluccius productus*). *J. Food Sci.* 59: 272-276.

- Soeda, T., Toiguchi, S., Mumazawa, T., Sakaguchi, S. and Kuhara, C. 1996. Studies on the functionalities of microbial transglutaminase for food processing. II. Effects of microbial transglutaminase on the texture of gel prepared from Suketoudara surimi. *Bull. Jap. Soc. Food Sci. Technol.* 43: 780-786.
- Steel, R.D.D. and Torrie, J.H. 1980. Principles and procedures of statistics: A biometrical approach (2nd ed.) p. 862. McGraw-Hill. New York.
- Sultanbawa, Y. and Li-Chan, E.C.Y. 2001. Structural changes in natural actomyosin and surimi from ling cod (*Ophiodon elongatus*) during frozen storage in the absence or presence of cryoprotectants. *J. Agric. Food Chem.* 49: 4716-4725.
- Suzuki, T. 1981. Fish and Krill Protein: Processing Technology. Applied Science Publishers. London.
- Sych, J., Lacroix, C., Adambounou, L.T. and Castaigne, F. 1990. Cryoprotective effects of lactitol, palatinit and polydextrose on cod surimi proteins during frozen storage. *J. Food Sci.* 55: 356-360.
- Taguchi, T., Ishizaka, H., Tanaka, M., Nagashima, Y. and Amano, K. 1987. Protein-protein interaction of fish myosin fragments. *J. Food Sci.* 52: 1103-1108.
- Takeda, H. and Seki, N. 1996. Enzyme-catalyzed cross-linking and degradation of myosin heavy chain in walleye pollack surimi paste during setting. *Fish. Sci.* 62: 462-467.
- Torigai, M. and Konno, K. 1996. Pyrophosphate-accelerated actin denaturation mechanism in myofibril. *Fish. Sci.* 62: 307-311.
- Torley, P.J. and Young, O.A. 1995. Rheological changes during isothermal holding of salted beef homogenates. *Meat. Sci.* 39: 23-34.

- Toyohara, H. and Shimizu, Y. 1988. Relation between the modori phenomenon and myosin heavy chain breakdown in the threadfin bream gel. *Agric. Biol. Chem.* 52: 255-257.
- Toyohara, H., Sakata, T., Yamashita, K., Kinishita, M. and Shimizu, Y. 1990. Degradation of ovalfilefish meat gel caused by myofibrillar proteinase(s). *J. Food Sci.* 55: 364-368.
- Tokunaga, T. and Nishioka, F. 1988. The improvement of technology for surimi production from fatty Japanese sardine. *In Upgrading from Feed to Food. Proceedings of a national technical conference, Fatty Fish Utilization.* (Raleigh, N.C. ed). 88-04. UNC. Sea Grant.
- Trout, G.R. and Schmidt, G.R. 1983. Utilization of phosphates in meat products. *Proc. Recipr. Meat Conf.* 36: 24-28.
- Trout, G.R. and Schmidt, G.R. 1984. Effect of phosphate type and concentration, salt level and method of preparation on binding in restructured beef rolls. *J. Food Sci.* 49: 687-694.
- Trout, G.R. and Schmidt, G.R. 1986. Effect of phosphates on the functional properties of restructured beef rolls: the role of pH, ionic strength, and phosphate type. *J. Food Sci.* 51: 1416-1423.
- Trout, G.R. and Schmidt, G.R. 1987. The effect of cooking temperature on the functional properties of beef proteins: The role of ionic strength, pH, and pyrophosphate. *Meat Sci.* 20: 129-135.
- Tsai, S.F., Wang, J.H. and Jiang, S.T. 1989. Stability of muscle proteins from some subtropical fish. *J. Agric. Food Chem.* 37: 749-752.

- Tsai, G.J., Lin, S.M. and Jiang, S.T. 1996. Transglutaminase from *Streptoverticillium ladakanum* and application to minced fish product. *J. Food Sci.* 61: 1234-1238.
- Tsukamasa, Y. and Shimizu, Y. 1990. Setting property of sardine and Pacific mackerel meat. *Nippon Suisan Gakkaishi.* 56: 1105-1112.
- Tsukamasa, Y. and Shimizu, Y. 1991. Factors affecting the transglutaminase-associated setting phenomenon in fish meat sol. *Bull. Jap. Soc. Sci. Fish.* 57: 535-540.
- Tsukawasa, Y., Miyake, Y., Ando, M. and Makinodan, Y. 2002. Total activity of transglutaminase at various temperatures in several fish meats. *Fish. Sci.* 68: 929-933.
- Tsukamasa, Y., Sato, K., Shimizu, Y., Imai, C., Sugiyama, M., Minegishi, Y. and Kaqabata, M. 1993. ϵ -(γ -glutamyl) lysine crosslink formation in sardine myofibril sol during setting at 25°C. *J. Food Sci.* 59: 785-787.
- Turner, S., Senaratna, T., Touchell, D., Bunn, E., Dixon, K. and Tan, B. 2001. Stereochemical arrangement of hydroxyl groups in sugar and polyalcohol molecules as an important factor in effective cryopreservation. *Plant Sci.* 160: 489-497.
- VanWazer, J.R. 1971. Chemistry of the phosphates and condensed phosphates. *In* Symposium: Phosphates in Food Processing. (Deman, J.M. and Malnychyn, P., eds.) p. 1. The AVI Pub. Co., Inc., Westport, Conn.
- Van Wazer, J.R. and Callis, F.F. 1958. Metal complexing by phosphates. *Chem Rev.* 58: 1011-1046.

- Van Wazer, J.R. and Campanella, D.A. 1950. Structure and properties of the condensed phosphates. IV. Complex ion formation in polyphosphate solutions. *J. Am. Chem. Soc.* 72: 655-663.
- Wan, J., Kimura, I., Satake, M. and Seki, N. 1994. Effect of calcium ion concentration on the gelling properties and transglutaminase activity of walleye pollack surimi paste. *Fish. Sci.* 60: 107-114.
- Wan, J., Kimura, I., Satake, M. and Seki, N. 1995. Causes of inferior gel-forming ability of salmon surimi paste. *Fish. Sci.* 61: 711-715.
- Wang, S.L. and Lanier, T.C. 1998. Effects of endogenous fungal and microbial transglutaminase in pollock surimi gelation. Presented at the IFT Annual Meeting, Abstract No. 32-4. Atlanta. G.A.
- Watabe, S., Ushio, H., Iwamoto, M., Yamanaka, H. and Hashimoto, K. 1989. Temperature-dependency of rigor-mortis of fish muscle: myofibrillar Mg^{2+} -ATPase activity and Ca^{2+} uptake by sarcoplasmic reticulum. *J. Food Sci.* 54: 1107-1110.
- Wicker, L., Lanier, T.C., Hamann, D.D. and Akahane, T. 1986. Thermal transitions in myosin-ANS fluorescence and gel rigidity. *J. Food Sci.* 51: 1540-1543.
- Xiong, Y.L. 1997. Structure-function relationships of muscle proteins. *In* Food proteins and their applications. (Damodaran, S. and Paraf, A., eds.) p. 341-392. Marcel Dekker. New York.
- Xiong, Y.L. 2005. Role of myofibrillar proteins in water-binding in brine-enhanced meats. *Food Res. Int.* 38: 281-287.
- Xiong, Y.L. and Brekke, C.J. 1989. Changes in protein solubility and gelation properties of chicken myofibrils during storage. *J. Food Sci.* 54: 1141-1146.

- Xiong, Y.L. and Kupski, D.R. 1999. Time-dependent marinade absorption and retention, cooking yield, and palatability of chicken fillets marinated in various phosphates solutions. *Poultry Sci.* 78: 1048-1052.
- Yagi, H., Sakamoto, M., Wakameda, A. and Arai, K. 1985. Effect of inorganic polyphosphate on thermal denaturation of carp myofibrillar protein at low ionic strength. *Bull. Jap. Soc. Sci. Fish.* 51: 667-673.
- Yamamoto, Y., Okubo, T., Hatayama, S., Naito, M. and Ebisu, T. (1991). Frozen surimi product and process for preparing. US Patent 5,028,444.
- Yamazawa, M. 1990. Effect of high temperature heating on physical properties of kamaboko gel. *Nippon Suisan Gakkaishi.* 56: 497-503.
- Yasunaga, K., Abe, Y., Yamazawa, M. and Arai, K. 1996. Heat-induced change in myosin heavy chains in salt-ground meat with a food additive containing transglutaminase. *Nippon Suisan Gakkaishi.* 62: 659-668.
- Yasuyuki, T., Sato, K., Shimizu, Y., Imai, C., Sugiyama, M., Minegishi, Y. and Kawabata, M. 1993. ϵ -(γ -glutamyl) lysine crosslink formed in sardine myofibril sol during setting at 25°C. *J. Food Sci.* 58: 785-787.
- Yongsawatdigul, J., Park, J.W., Kolbe, E., Abu Dagga, Y. and Morrissey, M.T. 1995. Ohmic heating maximizes gel functionality of Pacific whiting surimi. *J. Food Sci.* 60: 10-14.
- Ziegler, G.R. and Aton, J.C. 1984. Mechanism of gel forming by proteins of muscle tissue. *Food Technol.* 38(5): 77-82.