

## Contents

	Page
Abstract	(5)
Acknowledgment	(7)
Contents	(8)
List of Tables	(10)
List of Figures	(11)
1. Introduction	1
Literature Review	2
1. Chemical composition of fish	2
2. Functional properties of fish protein	6
3. Protein oxidation	8
4. Surimi and surimi production	10
5. Biological (Intrinsic) factors affecting surimi quality	13
6. Chemistry of surimi gelation	15
7. Setting of surimi gel	19
8. Softening of surimi gel	20
9. Ingredient in surimi	21
10. Oxidizing agent	26
Objective	28
2. Materials and methods	29
1. Chemicals	29
2. Instruments	29
3. Fish preparation	30
4. Study on the effect of oxidizing agent in wash water on of washed mince	30

## Contents (continued)

	Page
5. Study on the effect of washing time on gelling properties and physicochemical properties of mince washed with oxidizing agent	35
6. Study on the effect of some chemicals and MTGase on gel properties of surimi prepared by washing with oxidizing agent	36
7. Study on the effect of washing with oxidizing agent on setting of surimi	37
8. Statistical analysis	37
3. Result and Discussion	38
1. Effect of washing with various oxidizing agents on chemical composition, physicochemical properties and gel forming ability of fish mince	38
2. Effect of washing times with oxidizing agents on chemical composition and gel-forming ability of fish mince	73
3. Effect of MTGase and some chemicals on gel improvement of surimi produced by oxidizing agent washing	90
4. Effect of MTGase and chemicals on microstructure of surimi gel	101
5. Effect of washing with oxidizing agent on setting of surimi	105
4. Conclusion	114
Reference	115
Appendix	131
Vitae	143

## List of Tables

Table	Page
1 Whiteness of surimi gels produced from bigeye snapper stored in ice for different times and washed with different washing media	57
2 Whiteness of surimi gels produced from threadfin bream stored in ice for different times and washed with different washing media	57
3 Whiteness of surimi gels produced from starry triggerfish stored in ice for different times and washed with different washing media	58
4 Whiteness of surimi gels produced from bigeye snapper during iced storage and washed with 20 ppm NaOCl for different times.	84
5 Whiteness of surimi gels produced from threadfin bream during iced storage and washed with 20 ppm NaOCl for different times.	85
6 Whiteness of surimi gels produced from starry triggerfish during iced storage and washed with 20 ppm NaOCl for different times.	85
7 Whiteness of surimi gel prepared by 20 ppm NaOCl washing with addition of MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> .	100
8 Solubility of surimi gel produced from bigeye snapper, threadfin bream and starry triggerfish with CaCl <sub>2</sub> and EGTA addition.	113

## List of Figures

Figure	Page
1 Model of myosin molecule	4
2 Structure of actin, troponin and tropomyosin	5
3 Reaction catalyzed by TGase	22
4 Scheme for the superoxide radical-dependent polymerization of MHC.	25
5 Scheme for surimi preparation	32
6 Scheme for surimi gel preparation	33
7 Changes in pH of bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with different washing media.	39
8 Changes in TCA-soluble peptides in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with different washing media.	41
9 Changes in TBARS in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with different washing media.	43
10 Changes of TVB in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with different washing media.	45
11 Changes of TMA in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with different washing media.	46
12 Changes in breaking force of gel of surimi produced from bigeye snapper stored in ice for different times and washed with different washing media.	49

## List of Figures (continued)

Figure	Page
13 Changes in deformation of gel of surimi produced from bigeye snapper stored in ice for different times and washed with different washing media.	50
14 Changes in breaking force of gel of surimi produced from threadfin bream stored in ice for different times and washed with different washing media.	52
15 Changes in deformation of gel of surimi produced from threadfin bream stored in ice for different times and washed with different washing media.	53
16 Changes in breaking force of gel of surimi produced from starry triggerfish stored in ice for different times and washed with different washing media.	54
17 Changes in deformation of gel of surimi produced from starry triggerfish stored in ice for different times and washed with different washing media.	55
18 Surface hydrophobicity of NAM extracted from bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with water or 20 ppm NaOCl.	60
19 Total sulfhydryl group and disulfide bond contents of NAM extracted from bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with water or 20 ppm NaOCl.	63
20 $\text{Ca}^{2+}$ -ATPase, $\text{Mg}^{2+}$ -ATPase, $\text{Mg}^{2+}$ - $\text{Ca}^{2+}$ -ATPase and $\text{Mg}^{2+}$ -EGTA-ATPase activities of NAM extracted from bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with water or 20 ppm NaOCl.	65

### List of Figures (continued)

Figure	Page
21 SDS-PAGE pattern of bigeye snapper, threadfin bream and starry triggerfish stored in ice for different times and washed with water.	68
22 SDS-PAGE pattern of bigeye snapper, threadfin bream and starry triggerfish stored in ice for different times and washed with 20 ppm NaOCl.	69
23 SDS-PAGE pattern of surimi gel of bigeye snapper, threadfin bream and starry triggerfish stored in ice for different times and washed with water.	71
24 SDS-PAGE pattern of surimi gel of bigeye snapper, threadfin bream and starry triggerfish stored in ice for different times and washed with 20 ppm NaOCl.	72
25 Changes in pH of bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with 20 ppm NaOCl for 5, 10 and 15 min.	74
26 Changes in TCA-soluble peptides in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with 20 ppm NaOCl for 5, 10 and 15 min.	75
27 Changes in breaking force of gel of surimi produced from bigeye snapper stored in ice for different times and washed with water or 20 ppm NaOCl for different times.	78
28 Changes in deformation of gel of surimi produced from bigeye snapper stored in ice for different times and washed with water or 20 ppm NaOCl for different times.	79

### List of Figures (continued)

Figure	Page
29 Changes in breaking force of gel of surimi produced from threadfin bream stored in ice for different times and washed with water or 20 ppm NaOCl for different times.	80
30 Changes in deformation of gel of surimi produced from threadfin bream stored in ice for different times and washed with water or 20 ppm NaOCl for different times.	81
31 Changes in breaking force of gel of surimi produced from starry triggerfish stored in ice for different times and washed with water or 20 ppm NaOCl for different times.	82
32 Changes in deformation of gel of surimi produced from starry triggerfish stored in ice for different times and washed with water or 20 ppm NaOCl for different times.	83
33 Changes of TBARS in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with 20 ppm NaOCl for 5, 10 and 15 min.	87
34 Changes of TVB in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with 20 ppm NaOCl for 5 min.	88
35 Changes of TMA in bigeye snapper, threadfin bream and starry triggerfish mince stored in ice for different times and washed with 20 ppm NaOCl for 5 min.	89
36 Changes in breaking force and deformation of the gel of bigeye snapper surimi prepared by 20 ppm NaOCl washing in combination with MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> addition.	93

## List of Figures (continued)

Figure	Page
37 Changes in breaking force and deformation of the gel of threadfin bream surimi prepared by 20 ppm NaOCl washing in combination with MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> addition.	94
38 Changes in breaking force and deformation of the gel of starry triggerfish surimi prepared by 20 ppm NaOCl washing in combination with MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> addition.	95
39 SDS-PAGE pattern of the gel of bigeye snapper surimi prepared by 20 ppm NaOCl washing with addition of MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> .	97
40 SDS-PAGE pattern of the gel of threadfin bream surimi prepared by 20 ppm NaOCl washing with addition of MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> .	98
41 SDS-PAGE pattern of the gel of starry triggerfish surimi prepared by 20 ppm NaOCl washing with addition of MTGase, <i>L</i> -ascorbic acid or CaCl <sub>2</sub> .	99
42 Microstructure of surimi gel prepared from bigeye snapper, washed with water, washed with 20 ppm NaOCl, washed with 20 ppm NaOCl and added with 0.2 % MTGase added, washed with 20 ppm NaOCl and added with 0.1 % <i>L</i> -ascorbic acid washed with 20 ppm NaOCl and added with 50 mmole CaCl <sub>2</sub> /kg.	102
43 Microstructure of surimi gel prepared from threadfin bream, washed with water, washed with 20 ppm NaOCl, washed with 20 ppm NaOCl and added with 0.2 % MTGase.	103
44 Microstructure of surimi gel prepared from starry triggerfish, washed with water, washed with 20 ppm NaOCl, washed with 20 ppm NaOCl and added with 0.2% MTGase.	104



### List of Figures (continued)

Figure	Page
45 Breaking force and deformation of surimi gel prepared from bigeye snapper added with CaCl <sub>2</sub> or EGTA.	107
46 Breaking force and deformation of surimi gel prepared from threadfin bream added with CaCl <sub>2</sub> or EGTA.	108
47 Breaking force and deformation of surimi gel prepared from starry triggerfish added with CaCl <sub>2</sub> or EGTA.	109
48 SDS-PAGE pattern of gel surimi from bigeye snapper, threadfin bream and starry triggerfish.	111