

## Bibliography

- Adams, D.M. and Cornell, J.B. 1967. "Metal-Sulphur Vibrations. Part I. Far-infrared Spectra of Some Complexes of Thiourea and Ethylenethiourea (Imidazoline-2-thione)", J. Chem. Soc. (A). 1967, 884-889.
- Aslanidis, P., Cox, P.J., Divanidis, S. and Tsipis, A.C. 2002. "Copper(I) Halide Complexes with 1,3-Propanebis(diphenylphosphine) and Heterocyclic Thione Ligands: Crystal and Electronic Structures (DFT) of [CuCl(pymtH)(dppp)], [CuBr(pymtH)(dppp)], and [Cu( $\mu$ -I)(dppp)]<sub>2</sub>", Inorg. Chem. 41(2002), 6875-6886.
- Banerjee, S.N. and Sukthankar, A.C. 1962. "Complex Compounds of Substituted Thiourea. Part III. Copper Derivatives of Mono- and Di-*N*-acetylthiourea", J. Indian. Chem. Soc. 39(1962), 197 – 202.
- Banerjee, S.N. and Sukthankar, A. C. 1963. "Complex Compounds of Substituted Thiourea. Part IV. Copper Derivatives of Mono-*N*-phenylthiourea and Di-*N*-phenylthiourea", J. Indian. Chem. Soc. 40(1963), 387-393.
- Battaglia, L.P., Corradi, A.B., Nardelli, M. and Vidoni Tani, M.E. 1976. "X-ray Crystal Structures of Tetrakis(imidazolidine-2-thionato)copper(I) Nitrate and Dichloro- $\mu$ -imidazolidine-2-thionato-tris(imidazolidine-2-thionato)dicopper (I)", J. Chem. Soc. Dalton Tans. (1976), 143-146.

Baumgartner, M., Schmalle, H. and Dubler, E. 1993. "Structural and spectroscopic features of the 'adamantane' type Cu<sub>4</sub>S<sub>6</sub> core found in Cu(I) thiolates with bidentate chelate ligands", Inorg. Chim. Acta. 208(1993), 135-143.

Bott, R.C., Bowmaker, G.A., Davis, C.A., Hope, G.A. and Jones, B.E. 1998. "Crystal Structure of [Cu<sub>4</sub>(tu)<sub>7</sub>](SO<sub>4</sub>)<sub>2</sub>.H<sub>2</sub>O and Vibrational Spectroscopic Studies of Some Copper(I) Thiourea Complexes", Inorg. Chem. 37(1998), 651-657.

Bowmaker, G.A., Pakawatchai, C., Skelton, B.W., Thavornyuthikarn, P., Wattanakanjana, Y. and White, A.H. 1994. "Vibrational Spectra and Crystal Structures of Tris- and Tetrakis-(ethylenethiourea)copper(I) Systems", Aust. J. Chem. 47(1994), 15-24.

Budavari, S., 1989. The Merck Index, 11<sup>th</sup> ed., New Jersey: Merck & co., Inc.

Cernak, J., Chomic, J., Kutschy, P., Svrcinova, D. and Dzurilla, M. 1991. "Preparation and characterization of the complexes Cu(ptu)<sub>2</sub>Cl, Cu(etu)<sub>2</sub>Cl and Cu(btu)<sub>2</sub>Cl (ptu: R = phenyl, etu: R = ethyl, btu: R = benzyl in CH<sub>3</sub>CH=CH-CO-NH-CS-NH-R); crystal structure of the Cu(ptu)<sub>2</sub>Cl complex", Inorg. Chim. Acta. 181(1991), 85-92.

Cingi, M.B., Lanfranchi, M., Marchio, L. and Pellinghelli, M.A. 1971. "The Crystal and Molecular Structure of Tris(thiourea) Copper(I)hydrogen-*o*-phthalate. A monomeric Trigonal Planar Copper(I) Complex", Acta Cryst. B33(1971), 3772-3777.

Cingi, M.B., Bigoli, F., Lanfranchi, M., Leporati, E., Pellinghelli, M.A. and Foglia, C. 1995. "4-Amino-3-methyl-1,2,4- $\Delta^2$ -triazoline-5-thione : an example of thione-thiol tautomerism and stabilization of Cu(I) and Au(I) complexes", Inorg. Chim. Acta. 235(1995), 37-43.

Clegg, W. 1998. Crystal Structure Determination. New York: Oxford University Press.

Cotton, F.A. and Wilkinson, G. 1988. Advance Inorganic Chemistry, 5<sup>th</sup> ed., New York: John Wiley & Sons.

Crumbliss, A.L., *et al.* 1974. "Preparation and X-ray Crystal Structure of Novel TetranuclearCopper(I)Ethylenethiourea Cluster Complex,  $\mu_4$ -thylenethiourea-cyclo-tetrakis- $\mu$ -(ethylenethiourea)tetrakis [ethylenethioureacopper(I)]nitrate Hexahydrate", J. C. S. Chem Comm.(1974), 545-546.

Dubler, E. and Bensce, W. 1986. " $\text{Cu}(N,N'$ -dimethylthiourea)<sub>2</sub> $\text{NO}_3$ , a Copper(I) Complex Exhibiting Infinite Chains of Edge-sharing  $\text{CuS}_4$  Tetrahedral", Inorg. Chim. Acta. 125(1986), 37-42.

Flint, C.D. and Goodgame, M. 1966. "Infrared Spectra (400-135 cm<sup>-1</sup>) of Some Thiourea Co-ordination Complexes", J. Chem. Soc. (A). (1966), 744-747.

Frost, J.G., Lawson, M.E. and McPherson, W.G. 1976. "Nature and Cumulative Stability Constants of Copper(I) Complexes of Some Substituted Thioureas", Inorg. Chem. 15(1976), 940-943.

Fujisawa, K., Imai. S., Suzuki, S., Moro-oka, Y., Miyashita, Y., Yamada, Y. and Okamoto, K. 2000. "M-S vibrational study in three-coordinate thiolato compounds (NEt<sub>4</sub>)<sub>2</sub>[M(SC<sub>6</sub>H<sub>4</sub>-*p*-X)<sub>3</sub>] and (NEt<sub>4</sub>)<sub>2</sub>[M<sub>4</sub>(μ-SC<sub>6</sub>H<sub>4</sub>-*p*-Cl)<sub>6</sub>]: M = Cu(I) and Ag(I), X = Cl and Br", J. Inorg. Biochem. 82(2000), 229-238.

Griffith, H.E., Hunt, G.W. and Amma, E.L. 1976. "The Adamantane Structure in Polynuclear Cu<sub>4</sub>S<sub>6</sub>: the Crystal and Molecular Structures of Cu<sub>4</sub>[SC(NH<sub>2</sub>)<sub>2</sub>]<sub>6</sub>(NO<sub>3</sub>)<sub>4</sub>.4H<sub>2</sub>O and Cu<sub>4</sub>[SC(NH<sub>2</sub>)<sub>2</sub>]<sub>9</sub>(NO<sub>3</sub>)<sub>4</sub>.4H<sub>2</sub>O", J. C. S. Chem. Comm. 1976, 432-433.

Hall, S.R., King, G S D. and Stewart, J.M. 1999. Eds. Xtal3.6System. University of Western Australia. Lamp, Perth.

Hahn, T. 1983. International Tables for Crystallography (Volume A space-group symmetry). Boston: D. Reidel Publishing company.

Hunt, G.W., Terry III, N.W. and Amma, E.L. 1979 "The Structure of Tetrakis (thiourea)Copper(I) Hexafluorosilicate", Acta Cryst. B35(1979), 1235-1236.

Jasim, K.S. and Chieh, C. 1985. "Group VIB Metal Pentacarbonyl Complexes of *N,N'*-dimethylimidazolidine-2-thione", Inorg. Chim. Acta. 99(1985), 25-30.

Kitagawa, S., Nozaka, Y., Munakata, M. and Kawata, S. 1992. "Synthesis and crystal structures of tetra- and hexanuclear copper(I) complexes of pyrimidine derivatives,  $[\text{Cu}_4(\text{C}_4\text{H}_8\text{N}_2\text{S})_4](\text{ClO}_4)_4$  and  $[\text{Cu}_6(\text{C}_5\text{H}_5\text{N}_2\text{S})_6]$ ", Inorg. Chim. Acta. 197(1992), 169-175.

Kunchur, N.R. and Truter, M.R. 1958. "A Detailed Refinement of the Crystal and Molecular Structure of Thiourea", Crystal and Molecular Structure of Thiourea. 1958, 2551-2557.

Lane, T.J., Yamaguchi, A., Quagliano, J.V., Ryan, J.A. and Mizushima, S. 1959. "Infrared Absorption Spectra of Inorganic Coordination Complexes. XXII. Infrared Studies of Methylthiourea and its Metal Complexes", J. Amer. Chem. Soc. 81(1959), 3824-3826.

Malik, A.U. 1968. "Coordination Compounds of Cu(I) with Substituted Thioureas", J. Indian Chem. Soc. 45(2) (1968), 163-166.

Nakamoto, K. 1986. Infrared and Raman Spectra of Inorganic and Coordination Compounds., 4<sup>th</sup> ed., New York: John Wiley & Sons.

Okaya, Y. and Knobler, C. 1964. "Refinement of the crystal structure of tris(thiourea) copper(I)chloride", Acta Cryst. 17(1964), 928-930.

Pakawatchai, C., Thanyasirikul, Y., Saepae, T., Pansook, S., Fun, H and Chinnakali, K. 1998. "Hexakis( $\mu$ -N-ethylthiourea-S)tetrakis[iodocopper(I)] Monohydrate", Acta Cryst. C54(1998), 1750-1752.

Pettinari, C., Pettinari, R., Pellei, M. and Lobbia, G.G. 1999. "Synthesis and characterization of derivatives of copper(I) with N- and S-donor ligands: V. Imidazole and imidazoline-2(3H)-thione derivatives", Polyhedron. 18(1999), 1941-1951.

Radulescu, M., Pirvan, I., Dragomir, M., Popa, L. and Anghel, C. 2001. "The Evaluation of the Efficiency of Some Corrosion Inhibitors used in the Descaling Solutions of the Filmed Carbon Steels", Institute for Nuclear Research, Pitesti, Romania.

Schafer, M. and Curran, C. 1966. "Infrared Spectra of Complexes of Metal Halides with Tetramethylurea and Tetramethylthiourea", Inorg. Chem. 5(1966), 265-268.

Singh, R. and Dikshit, S.K. 1992. "Synthesis and characterization of mixed ligand copper(I) complexes containing halides, triphenylarsine and *N,N*-dimethyl-*N'*-phenylthiourea(dmptH), *N,N*-dibutyl-*N'*-phenylthiourea(dbptH) or 1,3-thiazolidine-2-thione(tzdtH). The X-ray crystal structure of [Cu(PPh<sub>3</sub>)<sub>2</sub>(dmptH)Cl]", Polyhedron. 14(1995), 1799-1807.

Stocker, F.B., Troester, M.A. and Britton, D. 1996. "Crystal Structures of a Family of New Copper(I) Cyanide Complexes of Thiourea and Substituted Thioureas", Inorg. Chem. 35(1996), 3154-3162.

Stocker, F.B., Britton, D. and Young, V.G. 2000. "Crystal Structures of a Family of Copper Cyanide Complexes of Thiourea and Substituted Thioureas", Inorg. Chem. 39(2000), 3479-3484.

Stout, G.H. and Jensen, L.H. 1989. X-ray Structure Determination. 2<sup>nd</sup> ed., New York: John Wiley & Sons.

Swaminathan, K. and Irving, H.M.N.H. 1964. "Infrared Absorption Spectra of Complexes of Thiourea", J. Inorg. Nucl. Chem. 26(1964), 1291-1294.

Taylor, I.F., Weininger, M.S. and Amma, E.L. 1974. "Preparation, Crystal Structure, and Bonding of the Dimers of Tris(thiourea)copper(I)Tetrafluoroborate and Tris(S-dimethylthiourea) copper(I) Tetrafluoroborate", Inorg. Chem. 13 (1974), 2835-2842.

Weininger, M.S., Hunt, G.W. and Amma, E.L. 1972. "Crystal and Molecular Structure of Tris(ethylenethiourea)Copper(I) Sulfate and Tris(tetramethylthiourea)Copper(I) Tetrafluoroborate[Examples of Trigonal Planar Copper(I) Stereochemistry]", J.C.S. Chem. Comm. (1972), 1140-1141.

Yamaguchi, A., Penland, R.B., Mizushima, S., Lane, T.J., Curran, C. and Quagliano, J.V. 1958. "Infrared Absorption Spectra of Inorganic Coordination Complexes. XIV. Infrared Studies of Some Metal Thiourea Complexes", J. Amer. Chem. Soc. 80(1958), 527-529.