

CONTENTS

	Pages
บทคัดย่อ	(3)
ABSTRACT	(5)
ACKNOWLEDGEMENT	(7)
THE RELEVANCE OF THE RESEARCH WORK	(8)
CONTENTS	(9)
LISTS OF TABLES	(14)
LISTS OF FIGURES	(18)
1 INTRODUCTION	1
1.1 Introduction	1
1.2 Background	5
1.2.1 Chemical identification	5
1.2.2 Physical and chemical properties	7
1.3 Literature review	9
1.3.1 Occurrence in food	9
1.3.2 Occurrence in environmental	11
1.3.3 Biological indices of exposure	12
1.3.4 Sample preparation	13
1.3.5 Analysis methods	14
1.4 Objectives	20
2 MATERIALS AND METHODS	21
2.1 Safety	21
2.2 Materials	21
2.2.1 Standard chemicals	21
2.2.2 Derivatization reagent	21
2.2.3 Other chemicals	22
2.2.4 Solid Phase Extraction (SPE)	23

CONTENTS (continued)

	Pages
2.3 Instruments and apparatus	23
2.3.1 Gas Chromatograph-Nitrogen Phosphorus Detector (GC-NPD)	23
2.3.2 Gas Chromatograph-Mass Spectrometer (GC-MS)	23
2.3.3 Apparatus for derivatization	24
2.3.4 Apparatus	24
2.4 Methods	25
2.4.1 Lab-built heating block unit and temperature calibration	25
2.4.2 Preparation of HCAs standard stock solution	26
2.4.3 Preparation of HCAs standard working solutions	26
2.4.4 Derivatization reaction of HCAs	26
2.4.5 Optimization of derivatization reaction for HCAs	27
2.4.5.1 Optimization of the reaction temperature	27
2.4.5.2 Optimization of the reaction time	27
2.4.6 Optimization of the GC-NPD conditions for HCAs analysis	28
2.4.6.1 Optimization of the carrier gas (He) flow rate	28
2.4.6.2 Optimization of the column temperature	28
2.4.6.3 Optimization of the injector temperature	29
2.4.6.4 Optimization of the detector temperature	30
2.4.6.5 Optimization of fuel (H ₂) flow rate	30
2.4.6.6 Optimization of oxidant (air) flow rate	30
2.4.7 Linear dynamic range (LDR)	30
2.4.8 Limit of detection (LOD)	32
2.4.9 Response factor	32

CONTENTS (continued)

	Pages
2.4.10 Confirmation of the HCA derivative structures and comparison of GC-MS and GC-NPD for HCAs analysis	32
2.4.10.1 Linear dynamic range in GC –MS (PE-17 ht column)	33
2.4.10.2 Limit of detection in GC–MS (PE-17 ht column)	33
2.4.11 Comparison of two different capillary columns, PE-17ht and HP-5	34
2.4.11.1 Linear dynamic range in GC-NPD (HP-5 column)	34
2.4.11.2 Limit of detection in GC -NPD (HP-5 column)	35
2.4.12 Optimization of sample preparation	35
2.4.12.1 Extraction time	39
2.4.12.2 Flow rate of eluting solvent	40
2.4.12.3 Type of eluent solvent	40
2.4.12.4 Percentage of toluene	40
2.4.12.5 Volume of eluting solvent	40
2.4.12.6 Volume of 0.5 M ammonium acetate, pH 8.0	41
2.4.12.7 Volume of methanol-ammonia solution (9:1, v/v)	41
2.4.13 Matrix Interference	42
2.4.14 Qualitative and quantitative analysis of heterocyclic amines in food	42
2.4.14.1 Meat sampling	42
2.4.14.2 Qualitative analysis	42
2.4.14.3 Quantification and recovery	42

CONTENTS (continued)

	Pages
3 RESULTS AND DISCUSSIONS	44
3.1 Optimization of the derivatization reaction	48
3.1.1 Calibration of the temperature for the lab-built heating unit	48
3.1.2 The reaction temperature	50
3.1.3 The reaction time	50
3.2 Optimization of the GC-NPD conditions for HCAs analysis	53
3.2.1 The carrier gas flow rate	53
3.2.2 The column temperature programming	58
3.2.3 The injector temperature	68
3.2.4 The detector temperature	70
3.2.5 The fuel gas flow rate (Hydrogen gas)	71
3.2.6 The oxidant gas flow rate (air)	74
3.2.7 Linear dynamic range (LDR)	76
3.2.8 Limit of detection (LOD)	80
3.2.9 Response factor	80
3.3 Confirmation of the HCA derivative structures	81
3.3.1 Linear dynamic range(LDR)	83
3.3.2 Limit of detection (LOD)	87
3.4 Comparison of two different capillary columns	88
3.4.1 Linear dynamic range (LDR) of HP-5 compared with PE-17 ht	89
3.4.2 Limit of detection (LOD)	93
3.5 Sample preparation optimization	94
3.5.1 The extraction time	94
3.5.2 The flow rate of eluting solvent	94

CONTENTS (continued)

	Pages
3.5.3 Selection of eluting solvent	97
3.5.4 Optimization of toluene amount	99
3.5.5 Amount of eluting solvent	99
3.5.6 Amount of 0.5 M ammonium acetate, pH 8.0	102
3.5.7 Amount of methanol: ammonia solution mixture (9:1, v/v)	102
3.6 Matrix interference	107
3.7 Qualitative and quantitative analysis of HCAs in cooked meat samples	112
3.7.1 Qualitative analysis	112
3.7.2 Recovery and Quantitative analysis	112
4 CONCLUSIONS	118
REFERENCES	121
VITAE	129

LISTS OF TABLES

Tables	Pages
1 Carcinogenicities of HCAs in rat and mice	4
2 Cancer potencies of some heterocyclic amines	5
3 Physical properties of IQ, MeIQx and PhIP	8
4 Methods for the identification and quantification of heterocyclic amines (HCAs) in cooked foods	19
5 Optimization of column temperature programming	29
6 The optimum derivatization reaction and GC-NPD conditions for HCA derivatives analysis	31
7 Optimization of sample preparation	39
8 The relationship between the set temperature and the actual temperature of the lab built heating block	48
9 The responses of three HCA derivatives standard solution of reaction temperature on the conversion of HCAs into their	
10 Responses of three N-dimethylaminomethylene HCA derivatives with various reaction times	52
11 Height Equivalent to a Theoretical Plate (HETP) of three HCA mixture derivative at various carrier gas flow rate	57
12 The responses of three HCA derivatives mixture on the isothermalcolumn temperatures	58
13 The responses of three HCA derivatives at various initial temperatures	59
14 The responses of three HCA derivatives at various hold times	60
15 The responses of HCA derivatives at various ramp rates for first stage	62

LISTS OF TABLES (continued)

Tables	Pages
16 The responses of HCA derivatives at various column temperatures at second stage	63
17 The responses of HCA derivatives at various hold times at second stage	64
18 The responses of HCA derivatives at various ramp rates at second stage	65
19 The responses of HCA derivatives at various column temperatures at the final stage	66
20 The responses of HCA derivatives at various hold times at final temperature	67
21 The responses of HCA derivatives at various injector temperatures	69
22 The responses of HCA derivatives at various detector temperatures	70
23 The effect of hydrogen gas flow rate on the responses of HCA derivatives	73
24 The effect of air flow rate on the responses of HCA derivatives	74
25 The optimum conditions of GC-NPD	75
26 The responses of IQ derivative at various concentrations	77
27 The responses of MeIQx derivative at various concentrations	78
28 The responses of PhIP derivative at various concentrations	79
29 The limit of detection of three HCAs with $S/N \geq 3$	80
30 Relative response factors of three HCAs at various concentrations	80
31 The responses of IQ derivative at various concentrations	84
32 The responses of MeIQx derivative at various concentrations	85
33 The responses of PhIP derivative at various concentrations	86
34 The limit of detection of three HCAs with $S/N \geq 3$	87

LISTS OF TABLES (continued)

Tables	Pages
35 The optimum conditions of GC-NPD equipped with HP-5 column	88
36 The linear dynamic range of the response of HP-5 compared with PE-17 ht to IQ at various concentrations	90
37 The linear dynamic range of the response of HP-5 compared with PE-17 ht to MeIQx at various concentrations	91
38 The linear dynamic range of the response of HP-5 compared with PE-17 ht to PhIP at various concentrations	92
39 The limit of detection of three HCAs with $S/N \geq 3$	93
40 The responses of three HCAs at various extraction times	95
41 The responses of HCAs at various flow rates	96
42 The responses of three HCAs eluted from diatomaceous earth with 5%toluene+dichloromethane, dichloromethane, ethyl acetate	98
43 Responses using various amount of toluene in mixed solvents	100
44 Responses using various amount of eluting solvents	101
45 Responses using various amount of 0.5 M ammonium acetate, pH 8.0	103
46 Responses using various amount of methanol: ammonia solution (9:1, v/ v)	104
47 Optimum conditions of sample preparation	105
48 The results of standard addition calibration curves of IQ in each cooked meat sample	109
49 The results of standard addition calibration curves of MeIQx in each cooked meat samples	110

LISTS OF TABLES (continued)

Tables	Pages
50 The result of standard addition calibration curves of PhIP in each cooked meat sample	111
51 The percentage of recoveries of the three HCAs in cooked meat samples	113
52 The results of standard addition calibration curves of IQ in each cooked meat sample	114
53 The results of standard addition calibration curves of MeIQx in each cooked meat samples	115
54 The result of standard addition calibration curves of PhIP in each cooked meat sample	116
55 The HCAs concentration of samples determined by standard addition method	117

LISTS OF FIGURES

Figures	Pages
1 Structure of IQ	6
2 Structure of MeIQx	6
3 Structure of PhIP	7
4 Structure of DMF-DMA	22
5 A Lab built heating block unit	25
6 The glass column packed with diatomaceous earth	36
7 Solid phase extraction of propylsulfonic acid (PRS)	37
8 Solid phase extraction of propylsulfonic acid (PRS) cartridges coupled with C ₁₈ cartridges	37
9 Summary of the sample preparation procedure	38
10 Schiff base reaction of IQ	45
11 Schiff base reaction of MeIQx	11
12 Schiff base reaction of PhIP	47
13 Temperature read from every 5 minutes	49
14 The relationship between the set temperature and the measured temperature	49
15 The responses of three HCA derivatives standard solution of reaction temperature on the conversion of HCAs into their N-dimethylaminomethylene derivatives	51
16 Responses of three N-dimethylaminomethylene HCA derivatives with various reaction times	52
17 Relative magnitudes of the different terms in the van Deemter equation	54
18 Capillary Chromatogram for calculating the theoretical plates	56
19 The van Deemter plot of three HCA derivatives	57

LISTS OF FIGURES (continued)

Figures	Pages
20 The responses of three HCA derivatives at various initial temperatures	59
21 The responses of three HCA derivatives at various hold times	60
22 The responses of HCA derivatives at various ramp rates for first stage	62
23 The responses of HCA derivatives at various column temperatures at second stage	63
24 The responses of HCAs derivatives at various hold times at second stage	64
25 The responses of HCA derivatives at various ramp rates at second stage	65
26 The responses of HCA derivatives at various column temperatures at final stage	66
27 The responses of HCA derivatives at various hold times at final temperature	67
28 The optimum column temperature programming	68
29 The responses of HCA derivatives at various injector temperatures	69
30 The responses of HCA derivatives at various detector temperatures	70
31 Nitrogen Phosphorus Detector (NPD)	72
32 The effect of hydrogen gas flow rate on the responses of HCA derivatives	73
33 The effect of air flow rate on the responses of HCA derivatives	74
34 The chromatograms of three HCAs derivatives at optimum conditions on capillary column (PE-17 ht) with GC-NPD	76
35 The linear dynamic range of IQ derivative	77

LISTS OF FIGURES (continued)

Figures	Pages
36 The linear dynamic range of MeIQx derivative	78
37 The linear dynamic range of PhIP derivative	79
38 Total ion chromatograms of heterocyclic amine derivatives confirmed by GC-MS	81
39 Mass spectrum of the N-dimethylaminomethylene derivative of IQ	82
40 Mass spectrum of the N-dimethylaminomethylene derivative of MeIQx	82
41 Mass spectrum of the N-dimethylaminomethylene derivative of PhIP	83
42 The linear dynamic range of mass selective detector to IQ	84
43 The linear dynamic range of mass selective detector to MeIQx	85
44 The linear dynamic range of mass selective detector to PhIP	86
45 The chromatograms of three HCAs at optimum conditions on HP-5 capillary column	89
46 The linear dynamic range of the response of HP-5 compared with PE-17 ht to IQ at various concentrations	90
47 The linear dynamic range of the response of HP-5 compared with PE-17 ht to MeIQx at various concentrations	91
48 The linear dynamic range of the response of HP-5 compared with PE-17 ht to PhIP at various concentrations	92
49 The responses of three HCAs at various extraction times	95
50 The responses of three HCAs at various flow rates	96

LISTS OF FIGURES (continued)

Figures	Pages
51 The responses of three HCAs eluted from diatomaceous earth with 5%toluene in dichloromethane, dichloromethane, ethyl acetate	98
52 Responses using various amount of toluene in mixed solvents	100
53 Responses using various amount of eluting solvents	101
54 Responses using various amount of 0.5 M ammonium acetate, pH 8.0	103
55 Responses using various amount of methanol: ammonia solution (9:1, v/ v)	104
56 GC-NPD chromatograms HCAs fraction isolated from spiked meat samples at optimum condition of sample preparation	106
57 GC-NPD chromatograms of HCAs of fraction separated from spiked in meat samples	108
58 The results of standard addition calibration curves of IQ in each cooked meat sample	109
59 The results of standard addition calibration curves of MeIQx in each cooked meat sample	110
60 The results of standard addition calibration curves of PhIP in each cooked meat sample	112
61 The percentage of recoveries of the three HCAs in cooked meat samples	113
62 Standard addition calibration curves of IQ in each cooked meat sample	114
63 Standard addition calibration curves of MeIQx in each cooked meat sample	115

LISTS OF FIGURES (continued)

Figures	Pages
64 Standard addition calibration curves of PhIP in each cooked meat sample	116