

Contents

	Page
Abstract	iii
Acknowledgements	vii
Contents	viii
List of Tables	x
List of Figures	xii
Chapter	
1. Introduction	1
1.1 Background and rationale	1
1.2 Review of literature	3
1.3 Objectives for studies	12
1.4 Study site	12
1.5 Road map of the present study	14
2. Methodology	16
2.1 Data managements	16
2.2 Phytoplankton abundance	16
2.2.1 Data source	16
2.2.2 Steps for model development	19
2.2.3 Statistical methods	20
2.3 Polychaeta abundance	24
2.2.1 Data source	24
2.3.2 Path diagram and variables	25

	Page
2.3.3 Statistical methods	26
3. Modeling of phytoplankton and Polychaeta density	28
3.1 Preliminary analysis	28
3.2 The first article	36
3.3 The second article	46
4. Discussion and Conclusions	52
4.1 The statistical modeling of distribution and abundance of phytoplankton	52
4.2 The statistical modeling of Polychaeta density in Na Thap River	57
4.3 limitations and suggestions for further study	58
References	59
Vitae	71

List of Tables

Tables	Page
2.1 Sampling stations, location and description of each station in Na Thap River	18
2.2 Sampling stations, location and description of each station in Na Thap River	24
3.1 Percentages of phytoplankton occurrence of 31 phytoplankton genera in the Na Thap River from June 2005 to December 2007 (n = 144)	29
3.2 The taxa, distribution (station), maximum densities and bimonthly occurrence of phytoplankton in the Na Thap River from June 2005 to December 2007. Data in bold indicates station and month of maximum density.	30
3.2 (Continue)	31
3.3 Variables, their roles and data type	32
The first article	
1 Mean and maximum densities of 31 dominant phytoplankton genera in the upstream freshwater area (sites 1–5) and the downstream estuarine area (sites 6–10), during rainy (November–January) and dry (February–October) seasons in the Na Thap River from June 2005 to December 2007. The minimum densities of the dominant genera were zero, except for <i>Euglena</i> sp., which had 447 cells L21 and 82 cells L21 during rainy seasons in the freshwater zone and in the estuarine zone, respectively. Sample sizes of genera were 13 and 51 for the rainy and the dry seasons, respectively, in the freshwater zone, and 15 and 65, respectively, in the estuarine zone.	39
2 Factor analysis (loadings, 0.2 omitted).	42
3 Coefficients and standard errors (in parenthesis) from fitting the multivariate multiple regression model with all nine predictors.	

Coefficients with p-values ≤ 0.05 are omitted; those adjudged not honestly statistically significant are shown in italics, and those with p-values ≤ 0.01 are shown in bold. The detailed abbreviation of each genus is shown in Table 1; factor 1 refers to salinity; factor 2 refers to heavy metal; factor 3 refers to water clarity; factor 4 refers to quantity of bacteria; factor 5 refers to dissolved oxygen.

Prince of Songkla University
Pattani Campus

List of Figures

Figures		Page
2.1	The Na Thap River and sampling stations (labeled 1-10)	17
2.2	Dataset of the study of Phytoplankton abundance	19
2.3	Steps for development of model to assess phytoplankton density	20
2.4	The Na Thap River and sampling stations (labeled 1-5)	23
2.5	Dataset of the study of Polychaetes abundance	25
2.6	Path diagram showing roles of variables	25
3.1	Water salinity in Na Thap River from June 2005 to May 2010 by station	33
3.2	Water salinity in Na Thap River from June 2005 to May 2010 by month	33
3.3	Natural logarithms of Polychaeta organisms per square meter in Na Thap River from June 2005 to May 2010 by station	34
3.4	Natural logarithms of Polychaeta organisms per square meter in Na Thap River from June 2005 to May 2010 by month	34
The first article		
1	Sampling sites (labeled 1–10) in the Na Thap River with Universal Transverse Mercator (UTM) coordinates (1000 3 1000 m).	37
2	Patterns of physicochemical variables: salinity, total sulfate, turbidity, transparency, water pH, phosphate–phosphorus, ammonia–nitrogen, and nitrate–nitrogen in the Na Thap River from June 2005 to December 2007 by site (left panel) and month (right panel).	40

Figures	Page
3	Patterns of physicochemical variables: dissolved oxygen, total iron, cadmium, copper, biochemical oxygen demand, oil and grease, total coliform bacteria, and fecal coliform bacteria in the Na Thap River from June 2005 to December 2007 by site (left panel) and month (right panel). 41
The second article	
1	Map with sampling stations 49
2	Histograms of Polychaeta organisms per square meter before (left panel) and after transformation using zero-corrected natural logarithms (right panel). 50
3	Polychaeta densities observed by months elapsed over duration of study with 95% confidence interval for gradient. The legend gives the number of overlapping points for observations from the 5 sites. The occasions when no Polychaeta were observed are as follows: at one site after 9, 32, 42 and 54 months, at two sites after 1, 3, 5 and 44 months, never at three sites, at four sites after 7 months, and at all five sites after 28 months. 50
4	Polychaeta densities by month elapsed adjusted for salinity (left panel), and by salinity adjusted for trend (right panel). 51