

## เอกสารอ้างอิง

- กรมวิชาการเกษตร. 2555. ถั่วหรั่ง. (Online). Available from: [www.jobdoa.com](http://www.jobdoa.com). [5 เมษายน 2555]
- จิราพร ศิริรัตน์. 2550. การศึกษาด้วยยีสโปรตีนจากถั่วหรั่งและการประยุกต์ใช้ วิทยานิพนธ์ปริญญาโท สาขาวิชาชีวเคมี มหาวิทยาลัยสงขลานครินทร์ วิทยาเขตปัตตานี
- ชนินันท์ ลิ้มปิษฐ์ชาล. 2551. ผลของปริมาณโปรตีนในกัมสำรองต่อสมบัติอิมัลชัน ปริมาณกรดไขมันอิสระ และความสามารถต้านออกซิเดชัน วิทยานิพนธ์ปริญญาโท สาขาวิชาวิทยาศาสตร์การอาหาร มหาวิทยาลัยเกษตรศาสตร์
- ณัฐวุฒิ ส่งแสง. 2550. การผลิตสารปรุงแต่งกลิ่นรสทะเลจากโปรตีนไฮโดรไลเสตจากถั่วเขียวโดยเอนไซม์โปรตีนเอส วิทยานิพนธ์ปริญญาโท สาขาวิชาเทคโนโลยีชีวเคมี มหาวิทยาลัยเทคโนโลยีพระจอมเกล้าธนบุรี
- ณรงค์ นิยมวิทย์. 2538. องค์ประกอบและการเปลี่ยนแปลงทางเคมีกายภาพของอาหาร พิมพ์ครั้งที่ 1. กรุงเทพฯ: ฟอร์แมทพรีนติ้ง
- นิธิยา รัตนานพนธ์. 2551. เคมีอาหาร พิมพ์ครั้งที่ 3. กรุงเทพฯ: โอเดียนสโตร์
- ปิยรัตน์ ศิริวงศ์ไพศาล. 2548. การปรับปรุงแป้งและสตาร์ชจากถั่วหรั่ง รายงานวิจัย มหาวิทยาลัยสงขลานครินทร์ วิทยาเขตหาดใหญ่
- ปราณี อ่านเปรื่อง. 2547. เอนไซม์ทางอาหาร พิมพ์ครั้งที่ 4. สำนักพิมพ์แห่งจุฬาลงกรณ์มหาวิทยาลัย
- พรชนัน เทียวทั่ว. 2548. การปรับปรุงสมบัติเชิงหน้าที่ของโปรตีนโอคาราโดยกระบวนการทางเคมี วิทยานิพนธ์ปริญญาโท สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหารลาดกระบัง
- ชั้นยพร จันทน์แสนโรจน์. 2550. การผลิตและสมบัติเชิงหน้าที่ของโปรตีนไฮโดรไลเสตจากหอยเป่าชื่อ *Haliotis asinina* วิทยานิพนธ์ปริญญาโท ภาควิชาเทคโนโลยีทางอาหารจุฬาลงกรณ์มหาวิทยาลัย
- ภูมิสันต์ จีวิพันธ์พงษ์ และ ธนาพร วีระประดิษฐ์ศิลป์. 2536. การศึกษาคุณสมบัติของถั่วหรั่งและการนำไปใช้ ภาควิชาเทคโนโลยีทางอาหาร คณะวิทยาศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย
- เยาวภา เขียรศิลากุล. 2549. กิจกรรมการเป็นสารต้านออกซิเดชันและคุณสมบัติเชิงหน้าที่ของโปรตีนไฮโดรไลเสตจากกล้ามเนื้อปลาหูแหก (*Decapterus maruadsi*) วิทยานิพนธ์ปริญญาโท สาขาเทคโนโลยีอาหาร มหาวิทยาลัยสงขลานครินทร์ วิทยาเขตหาดใหญ่
- รัตนา ชุมสาย ณ อยุธยา. 2525. วิเคราะห์องค์ประกอบของถั่วหรั่ง สาขาวิจัยเคมี และผลิตผลของเกษตรเคมี กรมวิชาการเกษตร: กรุงเทพฯ

วินัย ตะห์ลัน. 2542. ผลิตภัณฑ์โปรตีน (ออนไลน์) สืบค้นจาก:

[http://www.elibonline.com/doctor/food\\_protein01.html](http://www.elibonline.com/doctor/food_protein01.html) [17 เมษายน 2554]

สุภาวดี ทรัพย์ศิริไพบูลย์. 2550. ความสามารถในการเป็นสารต้านออกซิเดชันของโปรตีนถั่วเหลืองสกัดที่ผ่านการดัดแปรด้วยวิธีทางเอนไซม์และวิธีทางเคมี วิทยานิพนธ์ปริญญาโท สาขาวิชาวิทยาศาสตร์การอาหาร สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหารลาดกระบัง

A.O.A.C. 2000. Official methods of analysis. 16<sup>th</sup> ed. Washington, DC.: The Association of official analytical chemists.

Adebowale, K.O., Afolabi, T.A. and Lawal, O.S. 2002. Isolation, chemical modification and physicochemical charecterisation of bambara groundnut (*Voandzeia subterranea*) starch. *Nahrung/Food*. 46: 311-316.

Adler, N.J. 1986. Enzymic hydrolysate of food protein. Barking, UK.: Elsevier Applied Science Publisher.

Amarteifio, J.O. and Moholo, D. 1998. The chemical composition of four legumes consumed in Botswana. *Journal of Food Composition and Analysis*. 11: 329-332.

Anonymous. 2000. Novozyme. (Online). Available from: [www.Novozymes.com](http://www.Novozymes.com). [28 December 2010].

Belitz, H.D., Grosch, W. and Schieberle, P. 2009. Food Chemistry. 4<sup>th</sup> ed. Amino acids, Peptides, Proteins. Berlin, Germany: Springer – Verlag Berlin Heidelberg. 8-92.

Benjakul, S. and Morrissey, M.T. 1997. Protein hydrolysates from Pacific whiting solid wastes. *Food Chemistry* 145: 3423–3430.

Binsan, W., Benjakul, S., Visessanguan, W., Roytrakul, S., Tanaka, M. and Kishimura, H. 2008. Antioxidative activity of Mungoong, an extract paste, from the cephalothorax of white shrimp (*Litopenaeus vannamei*). *Food Chemistry*. 106: 185–193.

Borget, M. 1989. Les légumineuses vivrières tropicales. France: Maisson neuve et Larose et ACCT: 1-142.

Chabanon, G., Chevalot, I., Framboisier, X., Chenu, S. and Mare, I. 2007. Hydrolysis of rapeseed protein isolates: kinetics, characterization and functional properties of hydrolysates. *Process Biochemistry*. 42: 1419-1428.

- Chanput, W., Theerakulkait, C. and Nakai, S. 2009. Antioxidative properties of partially purified barley hordein, rice bran protein fractions and their hydrolysates. *Journal of Cereal Science*. 49: 422-428.
- Chen, H.M., Muramatoto, K., Yamauchi, F., Fujimoto, K. and Nokihara, K. 1998. Antioxidative properties of histidine-containing peptides designed from peptide fragments found in the digests of soybean protein. *Journal of Agricultural and Food Chemistry*. 46: 49–53.
- Chew, P., Casey A.J and Johnson S.K. 2003. Protein quality and physico - functionality of Australian sweet lupin (*Lupinus angustifolius cv. Gungurru*) protein concentrates prepared by isoelectric precipitation or ultrafiltration. *Food Chemistry*. 83: 575–583.
- Chuan-He, T., Jing, P., Da-Wen, Z. and Zhong, C. 2009. Physicochemical and antioxidant properties of buckwheat (*Fagopyrum esculentum Moench*) protein hydrolysates. *Food Chemistry*. 115: 672 – 678.
- Churuch, F.C., Porter, D.H., Catignani, G.L. and Swaisgood, H.E. 1985. An ophthaldehyde spectrophotometric assay for proteinases. *Analytical Biochemistry*. 146: 343-348.
- Chobert, J.M., Bertrand-Harb, C. and Nicolus, M.G. 1988. Solubility and emulsifying properties of caseins and whey proteins modified enzymatically by trypsin. *Journal of Agricultural and Food Chemistry*. 36: 883–892.
- Clemente, A., Vioque, J., Sánchez-Vique, R., Pedroche, J., Bautista, J. and Millán, F. 1999. *Food Chemistry*. 67: 269-274.
- Damodaran, S. and Paraf, A. 1997. *Food Protein and Their Application*. New York: Marcel Dekker, Inc.
- Decker, E.A. and Crum, A.D. 1993. Antioxidant activity of canosine in cooked ground pork. *Meat Science*. 58: 337-341.
- Dinis, T.C., Madeira, V.M. and Almeida, L.M. 1994. Action of phenolic derivatives (acetoaminophen, salicylate and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Archives of Biochemistry and Biophysics*. 315: 161-169.
- Duke, J.A., Okigbo, B.N., Reed, C.F. and Weder, J.K.P. 1986. *Voandzeia subterranean* (L). Thouars, *Handbook of Legume of World Economics Importance*. 307-310.

- Francis, G., Makkar, P.S. Harinder. and Becker, K. 2001. Antinutritional factors present in plant derived alternate fish feed ingredients and their effects in fish. *Aquaculture Research*. 199: 197-227.
- Gbogouri, G.A., Linder, M., Fanni, J. and Parmentier, M. 2004. Influence of hydrolysis degree on the functional properties of salmon byproduct hydrolysates. *Journal of Food Science*. 69: 615–622.
- Gordon, M. 2001. Antioxidants and food stability. In: Pokorny, J., Yanishlieva, N. and Gordon, M. (Eds.), *Antioxidant in Food*: 7–21. New York, USA: CRC Press. 158-322.
- Govindaraju, K. and Srinivas, H. 2004. Studies on the effect of enzymatic hydrolysis on functional and physico – chemical properties of arachin. *LWT-Food Science and Technology*. 56: 183-191.
- Gupta, Y.P. 1987. Anti-nutritional and toxic factors in food legumes: a review. *Plant Foods for Human Nutrition*. 37: 201–228.
- Halling, P.J. 1981. Protein-stabilized foams and emulsions. *Food Science and Nutrition*. 13: 155-203.
- Hamerstrand, G.E., Black, L.T. and Glover, J.D. 1981. Trypsin inhibitors in soy products: modification of the standard analytical procedure. *Cereal Chemistry*. 58: 42-45.
- Hill, R.L. 1965. *Advance in Protein Chemistry*. New York: John Wiley and Sons.
- Horiguchi, N., Horiguchi, H. and Suzuki, Y. 2005. Effect of wheat gluten hydrolysate on the immune system in healthy human subjects. *Bioscience, Biotechnology and Biochemistry*. 69: 2445–2449.
- Jamdar, S.N., Rajalakshmi, V., Pednekar, M.D., Juan, F., Yardi, V. and Sharma, A. 2010. Influence of degree of hydrolysis on functional properties, antioxidant activity and ACE inhibitory activity of peanut protein hydrolysate. *Food Chemistry*. 121: 178–184.
- Jookyeong, L. 2011. Soy protein hydrolysate; Solubility, Thermal stability, Bioactivity, and Sensory acceptability in tea bevaerage. A thesis submitted to the faculty of the graduate school. University of Minnesota.
- Kinsella, J.E. 1976. Functional properties of food proteins: a review. *Critical Reviews in Food Science and Nutrition*. 7: 219-280.

- Klompong, V., Benjakul, S., Kantachote, D. and Shaidi, F. 2007. Antioxidative activity and functional properties of protein hydrolysates of yellow stripe trevally (*Dearoides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. *Food Chemistry*. 102: 1317–1327.
- Klompong, V., Benjakul, S., Kantachote, D. and Shaidi, F. 2009. Characteristics and use of Yellow stripe trevally hydrolysate as culture media. *Journal of Food Science*. 74: 219-225.
- Kristinsson, G.H. and Rasco, A.B. 2000. Kinetics of the hydrolysis of Atlantic salmon (*Salmo salar*) muscle proteins by alkaline proteases and a visceral serine protease mixture. *Process Biochemistry*. 36: 131–139.
- La-ongdao, W., Premwadee, S., Kalyanee, J. and Parichat, H. 2011. Potential use of antioxidative mungbean protein hydrolysate as an anticancer Asiatic acid carrier. *Food Research International*. 44: 812–817.
- Lawal, O.S., Adebowale, K.O. and Adeb Owale, Y.A. 2007. Functional properties of native and chemically modified protein concentrates from bambarra groundnut. *Food Research International*. 40: 1003-1011.
- Li, G.H., Le, G.W., Liu, H. and Shi, Y.H. 2005. Mung-bean protein hydrolysates obtained with alcalase exhibit angiotensin I-converting enzyme inhibitory activity. *Food Science and Technology International*. 11: 281–287.
- Li, B., Chen, F., Wang, X., Ji, B. and Wu, Y. 2007. Isolation and identification of antioxidative peptides from porcine collagen hydrolysate by consecutive chromatography and electrospray ionization-mass spectrometry. *Food Chemistry*. 102: 1135–1143.
- Li, Y., Jiang, B., Zhang, T., Mu, W. and Liu, J. 2008. Antioxidant and free radical scavenging activities of chickpea protein hydrolysate (CPH). *Food Chemistry*. 106: 444 – 450.
- Linder, M., Fanni, J. and Parmentier, M. 1996. Functional properties of veal bone hydrolysates. *Journal of Food Science*. 61: 712–716.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with the Folin phenol reagent. *Biological Chemistry*. 193: 265–275.
- Mackie, I.M. 1982. Fish protein hydrolysate. *Process Biochemistry*. 20: 26–30.

- Material safety data sheet. Tannic acid MSDS. 2013. (Online). Available from:  
<http://www.sciencelab.com/msds.php?msdsId=9927618>. [7 February 2013].
- Mccarthy, T.L., Kerry, J.P., Kerry, J.F., Lynch, P.B. and Buckley, D.J. 2001. Assessment of the antioxidant potential of natural food and plant extracts in fresh and previously frozen pork patties. *Meat Science*. 57: 177-184.
- Mondor, M., Aksay, S., Drolet, H., Roufik, S., Farnworth, E. and Boye, I.J. 2009. Influence of processing on composition and antinutritional factors of chickpea protein concentrates produced by isoelectric precipitation and ultrafiltration. *Innovative Food Science and Emerging Technologies*. 10: 342-347.
- Moore, S. and Stein, W.H. 1948. Photometric ninhydrin method for use in chromatography of amino acids. *Biological Chemistry*. 176: 367-388.
- Moure, A. Dominguez, H. and Parajo, J.C. 2006. Antioxidant properties of ultrafiltration recovered soy protein fractions from industrial effluents and their hydrolysates. *Process Biochem*. 41: 447-456.
- Mune, M.A.M., Minka, S.R., Mbome I.L. and Etoa F.X. 2011. Nutritional potential of Bambara bean protein concentrate. *The Journal of Nutrition*. 10: 112-119.
- Mutilangi, W.A.M., Panyam, D. and Kilara, A. 1996. Functional properties of hydrolysates from proteolysis of heat-denatured whey protein isolate. *Journal of Food Science*. 61: 270-274.
- Nicorescu, I., Loisel, C., Vial, C., Riaublanc, A., Djelveh, G., Cuvelier, G. and Legrand, J. 2008. Combined effect of dynamic heat treatment and ionic strength on denaturation and aggregation of whey proteins-part I. *Food Research International*. 41: 707-713.
- Nalinanon, S., Benjakul, S., Kishimura H. and Shahidi F. 2011. Functionalities and antioxidant properties of protein hydrolysates from the muscle of ornate threadfin bream treated with pepsin from skipjack tuna. *Food Chemistry*. 124: 1354-1362.
- National academy of science. 1979. Bambara groundnut, In *Tropical Legumes: resources for the future*. Washington, DC.: National academy of sciences. 47-53.
- Panyam, D. and Kilara, A. 1996. Enhancing the functionality of food proteins by enzymatic modification. *Trends in Food Science and Technology*. 7: 120-125.

- Pearce, K.N. and Kinsella, J.E. 1978. Emulsifying properties of proteins: evaluation of a turbidimetric technique. *Journal of Agricultural and Food Chemistry*. 26: 716–723.
- Pena – Ramos, A.E. and Xiong, L.Y. 2003. Whey and soy protein hydrolysates inhibit lipid oxidation in cooked pork patties. *Meat Science*. 64: 259–263.
- Phanipa, P., Soottawat, B., Wonnop, V. and Sittiruk, R. 2010. Use of pyloric caeca extract from bigeye snapper (*Priacanthus macracanthus*) for the production of gelatin hydrolysate with antioxidative activity. *Journal of Food Science and Technology*. 43: 86–97.
- Phillips, L.G., Whitehead, D.M and Kinsella, J.E. 1994. Protein stabilized foams, Structure function of food Proteins. New York: Academic Press, Inc. 131–152.
- Pomeranz, Y. 1991. Functional properties of food components. 2<sup>nd</sup> Ed. San Diego, CA: Academic Press, Inc. 248–298.
- Prommool, S., Jongjareonrak, A. and Benjaku, S. 2010. Study on antioxidant activities of protein hydrolysate from Skipjack tuna (*Katsuwonus pelamis*) viscery. 35<sup>th</sup> Congress on Science and Technology of Thailand.
- Purseglove, J.W. 1977. Tropical Crop; dicotyledons. London: Longman Group Ltd. 719.
- Radha, C., Kumar Ramesh, P. and Prakash, V. 2007. Preparation and characterization of a protein hydrolysate from an oilseed flour mixture. *Food Chemistry*. 106: 1166-1174.
- Rafat, A., Philip, K. and Muniandy, S. 2010. Antioxidant potential and content of phenolic compounds in ethanolic extracts of selected parts of *Andrographis paniculata*. *Medicinal Journal of Plants Research*. 4: 197-202.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M. and Rice, E.C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorisation assay. *Free Radical Biology and Medicine*. 26: 1231-1237.
- Rebeca, B.D., Pena-vera, M.T. and Diaz-Castaneda, M. 1991. Production of fish protein hydrolysates with bacteria proteases; yield and nutritional value. *Journal of Food Science*. 56: 309–314.
- Reddy, N.R., Pierson, M.D., Sathe, S.K. and Salunkhe, D.K. 1985. Dry bean tannins a review of nutritional implications. *The American Oil Chemists Society*. 62: 541–549.
- Reed, J.D. 1995. Nutritional toxicology of tannin and related polyphenols in forage legumes. *Journal of Animal Science*. 73: 1516-1528.

- Ryan, M., McEvoy, E., Duignan, S., Crowley, C., Fenelon, M., O'Callaghan, D.M. and Fitzgerald, F.J. 2008. Thermal stability of soy protein isolate and hydrolysate ingredients. *Food Chemistry*. 108: 503-510.
- Sara, E., Molina, O. and Jorge, R.W. 2002. Hydrolysates of native and modified soy protein isolates: structural characteristics, solubility and foaming properties. *Food Research International*. 35: 511-518.
- Schofield, D.M. and Mbugua, A.N. 2001. Analysis of condensed tannins: a review. *Animal Feed Science and Technology*. 53: 21-40.
- Seutsuna, K. and Chen, J.R. 2002. Isolation and characterization of peptides with antioxidative activity derived from wheat gluten. *Food Science and Technology Research*. 8: 227-230.
- Shahidi, F., Han, X.Q. and Synowiecki, J. 1995. Production and characteristics of protein hydrolysates from capelin (*Mallotus villosus*). *Food Chemistry*. 53: 285-293.
- Shimada, K., Fujikawa, K., Yahara, K. and Nakamura, T. 1992. Antioxidative properties of xanthan on the antioxidation of soybean oil in cyclodextrin emulsion. *Journal of Agricultural and Food Chemistry*. 40: 945-948.
- Sikorski, Z.E. and Naczk, M. 1981. Modification of technological properties of fish protein concentrate. *CRC Critical Reviews in Food Science and Nutrition*. 14: 201-230.
- Silvestre, M.P.C. 1997. Review of methods for the analysis of protein hydrolysate. *Food Chemistry*. 60: 263-271.
- Séverin, S. and Wen-shui, X. 2006. Nutritional evaluation of caseins and whey proteins and their hydrolysates from Protamex. *Journal of Zhejiang University Science*. 7: 90-98.
- Singh, U. 1988. Antinutritional factors of chick pea and pigeon pea and their removal by processing. *International Journal of Food Science and Nutrition*. 38: 251-261.
- Sinha, R., Radha, C., Prakash, J. and Kaul, P. 2007. Whey protein hydrolysate: Functional properties, nutritional quality and utilization in beverage formulation. *Food Chemistry*. 101: 1484-1491.
- Sittiwat, L. and Narissara, Ki. 2000. Antioxidant activity and fatty acid constituents in durian seeds. *Nutrition Research*. Council Thailand. 33: 153-161.
- Skanderby, M. 1994. Protein hydrolysates: their functionality and applications. *European Food Research and Technology International*. 10: 141-144.



- Slinkard, K. and Singleton, V.L. 1977. Total phenolic analysis: Automation and comparison with manual methods. *American Journal of Enology and Viticulture*. 28: 49-55.
- Smyth, M. and Fitzgerald, R.J. 1998. Relationship between some characteristics of WPC hydrolysates and the enzyme complement in commercially available proteinase preparations. *International Dairy Journal*. 8: 819-827.
- Stohs, S.J. and Bagchi, D. 1995. Oxidative mechanisms in the toxicity of metal ions. *Free Radical Biology and Medicine*. 18: 321–336.
- Taha, F.S. and Ibrahim M.A. 2002. Effect of degree of hydrolysis on the functional properties of some oilseed proteins. *Grasasy Aceites*. 53: 273-281.
- Thammarat, K., Soottawat, B. and Wonnop, V. 2011. Effect of NaCl on thermal aggregation of egg white proteins from duck egg. *Food Chemistry*. 125: 706–712.
- Thiansilakul, Y., Benjakul, S. and Shahidi, F. 2007. Compositions, Functional properties and Antioxidant activity of protein hydrolysates prepared from round scad (*Decapterus maruadsi*). *Food Chemistry*. 103: 1385-1394.
- Tsumura, K., Saito, T., Tsuge, K., Ashida, H., Kugimiya, W. and Inouye, K. 2005. Functional properties of soy protein hydrolysates obtained by selective proteolysis. *Journal of Food Science and Technology*. 38: 255-261.
- Urbano, G., Lopez-Jurado, M., Aranda, P., Vidal-Valverde, C., Tenorio, E. and Porres, J. 2000. The role of phytic acid in legumes: antinutrient or beneficial function. *Journal of Physiological and Biochemistry*. 56: 283–294.
- Vaštag, Z., PopoviĆ, L., PopoviĆ, S., Krimer, V. and PeriĆin, D. 2010. Hydrolysis of pumpkin oil cake protein isolate and free radical scavenging activity of hydrolysates: Influence of temperature, enzyme/substrate ratio and time. *Food and Bioproducts Processing*. 88: 277 – 282.
- Whitaker, J.R. 1994. The proteolytic enzyme. In *Principles of Enzymology for the food Science*. New York: Marcel Dekker Inc. 469 -498.
- Wu, H.C., Chen, H.M. and Shiau, C.Y. 2003. Free amino acid and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Research International*. 36: 949-957.

- Wu, S.U. and Brew, M.S. 1994. Soy protein isolate antioxidant effect on lipid peroxidation of ground beef and microsomal lipids. *Journal of Food Science*. 49: 702-706.
- Yoshie-Stark, Y. and Wasche, A. 2004. In vitro binding of bile acids by lupin protein isolate and their hydrolysates. *Food Chemistry*. 88: 179–184.
- Yoshie-Stark, Y., Wada, Y. and Wasche, A. 2008. Chemical composition, functional properties and bioactivities of rapeseed protein isolates. *Food Chemistry*. 107: 32–39.
- Yusuf, A.A., Ayedun, H. and Sanni, L.O. 2008. Chemical composition and functional properties of raw and roasted Nigerian benniseed (*Sesamum indicum*) and Bambara groundnut (*Vigna subterranean*). *Food Chemistry*. 111: 277–282.
- Zayas, Joseph F., 1997. *Functionality of protein in food*. Berlin, Germany: Springer – Verlag Heidelberg. 261-267.
- Zhu, K., Zhou, H. and Qian, H. 2006. Antioxidant and free radical-scavenging activities of wheat germ protein hydrolysates (WGPH) prepared with alcalase. *Process Biochemistry*. 4: 1296-1302.

Prince of Songkla University  
Pattani Campus