เอกสารอ้างอิง

วัลลภ สันติประชา. 2531. เทคโนโลยีเมล็ดพันธุ์. คณะทรัพยากรธรรมชาติ มหาวิทยาลัยสงขลานครินทร์. หน้า 51-57.

Dekkers, J. A., Rao, N. A. and Goh, J. C. 1991. In vitro storage of multiple shoot cultures of gingers at ambient temperature of 24-29°C. Scientia Horticulturae. 47:157-167.

Fukai, S. 1990. Cryopreservation of chrysanthemun shoot tips. Scientia Horticulturae. 45:167-174.

Harither, G. and Ying, H. Y. 1984. The low fruit set that fallows conventional hand pollination in Hevea brasiliensis:Insufficiency of pollen as a couse. RRIM Journal. 32:20-29.

Heszky, E. L., Jekkel, Z. and Ali, H. A. 1990. Effect of cooling rate, cryoprotectant and holding time at different transfer temperature on the survival of cryoprerserved cell suspension culture (Puccinellia distana L.). Plant cell, Tissue and Organ Culture. 21:217-226.

Ko, H. W., Hwang, C. S. and Ku, M. F. 1991. A new technique for storage of meristem-tip cultures of Cavendish banana. Plant Cell, Tissue and Organ Culture. 25:179-183.

Reed, M. B. 1990. Survival of in vitro grown apical meristem of pyrus following cryopreservation. HortScience. 25(1):111-113.

Richard, W. J., Kumar, P. P. and Thorpe, A. T. 1991. Long-term storage of somatic embryogenic white spruce tissue at ambient temperature. Plant cell, Tissue and Organ Culture. 25:53-60.

Sharma, K. K. 1990. In vitro propagation of Malus alba through nodal segment. Scientia Horticulturae. 42:307-320.

Zamora, B. A., Damasco, P. O. and Landicho, S. 1989. Development of a minimal growth medium for *in vitro* storage of banana and plantain (*Musa* spp.) germplasm. The Philipphine Agriculturist. 72(4):466-472.