



**Induction of *Agrobacterium rhizogenes* Transformed Hairy Root Cultures
from *Mitragyna speciosa* (Roxb.) Korth.**

Siriwan Phongprueksapattana

**A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of
Master of Pharmacy in Pharmaceutical Sciences
Prince of Songkla University**

2007

Copyright of Prince of Songkla University

Thesis Title Induction of *Agrobacterium rhizogenes* Transformed Hairy Root Cultures from *Mitragyna speciosa* (Roxb.) Korth.
Author Miss Siriwan Phongprueksapattana
Major Program Pharmaceutical Sciences

Major Advisor

.....
(Assist. Prof. Dr. Juraithip Wungsintaweekul)

Co-advisor

.....
(Assoc. Prof. Dr. Waraporn Putalun)

Examining Committee:

.....Chairperson
(Assoc. Prof. Dr. Pharkphoom Panichayupakaranant)

.....Committee
(Assoc. Prof. Dr. Sompop Prathanturarug)

.....Committee
(Assist. Prof. Dr. Juraithip Wungsintaweekul)

.....Committee
(Assoc. Prof. Dr. Waraporn Putalun)

The Graduate School, Prince of Songkla University, has approved this thesis as partial fulfillment of the requirements for the Master of Pharmacy Degree in Pharmaceutical Sciences

.....
(Assoc. Prof. Dr. Kerkchai Thongnoo)
Dean of Graduate School

ชื่อวิทยานิพนธ์	การเหนี่ยวนำให้เกิดรากขนอ่อนจากต้นกระท่อมด้วยเชื้ออะโกราแบคทีเรียมไรโซจีนเนส
ผู้เขียน	นางสาวศิริวรรณ พงศ์พุกษาพัฒนา
สาขาวิชา	เภสัชศาสตร์
ปีการศึกษา	2550

บทคัดย่อ

รากขนอ่อนของต้นกระท่อม ได้จากการเหนี่ยวนำด้วยเชื้ออะโกราแบคทีเรียมไรโซจีนเนส ATCC 15834 จากชิ้นส่วนบริเวณเส้นกลางใบ รากขนอ่อนที่ได้ถูกนำมาวิเคราะห์หาชิ้นที่ก่อให้เกิดราก ได้แก่ *rolA* และ *rolB* ด้วยวิธี polymerase chain reaction ผลการทดลองพบว่าประสิทธิภาพของการถ่ายยีนจาก เชื้ออะโกราแบคทีเรียมไรโซจีนเนส มีค่าประมาณ 60% ลักษณะรากขนอ่อนที่ได้มีลักษณะแตกแขนง และมีขนอ่อนจำนวนมาก เมื่อศึกษาผลของอาหารเพาะเลี้ยงต่อการเจริญเติบโตของรากขนอ่อน จากอาหารเพาะเลี้ยงรวม 9 สูตรอาหาร พบว่าสูตรอาหาร WPM ที่เสริมด้วย 0.5 mg/l 1-naphthalene acetic acid เป็นสูตรอาหารที่เหมาะสมที่สุดต่อการเจริญเติบโตของรากขนอ่อน เมื่อศึกษาถึงการสร้างสารทุติยภูมิในรากขนอ่อน โดยการเตรียมเป็นสารสกัดเฮกเซนและสารถูกแยกด้วยเทคนิคคอลัมน์โครมาโตกราฟี ผลการทดลองได้สาร MSF1 และ MSF2 ในปริมาณ 0.016% (w/w) และ 0.027% (w/w) ตามลำดับ ทำการวิเคราะห์โครงสร้างของสารทั้งสองด้วยเทคนิค ^1H - และ ^{13}C -NMR spectroscopy พบว่าสาร MSF1 เป็นสารผสมระหว่าง β -sitosterol และ stigmasterol (อัตราส่วน 1:1) และสาร MSF2 เป็นสารผสมระหว่าง ursolic acid และ oleanolic acid (อัตราส่วน 5:1) จากนั้นได้สร้างกราฟเจริญเติบโตและกราฟการสร้างสาร ursolic acid ในรากขนอ่อน กราฟการเจริญเติบโตแสดงให้เห็นว่ารากขนอ่อนมีวงจรการเจริญเติบโตประมาณ 40 วัน และจากกราฟการสร้างสาร ursolic acid แสดงให้เห็นว่ารากขนอ่อนนั้นสามารถสร้างสาร ursolic acid และมีปริมาณ ursolic acid สะสมสูงสุดคือ 3.47 ± 0.03 มก.ต่อน้ำหนักแห้ง ในช่วงปลายของระยะ linear phase ในการศึกษาครั้งนี้ยังได้ศึกษาการขยายพันธุ์ต้นกระท่อมได้ตลอดทดลอง โดยศึกษาถึงการเพิ่มจำนวนยอดจากตาข้างของต้นกระท่อม ในสูตรอาหาร WPM ที่เสริมด้วย 0.1 mg/l thidiazuron โดยต้นที่เจริญขึ้นใหม่ที่มีลักษณะต้นที่สมบูรณ์แล้ว ได้ถูกนำมาปรับสภาพเพื่อนำมาเลี้ยงในสภาวะธรรมชาติ พบว่าในกระบวนการนี้ต้นกระท่อมมีอัตราการรอดประมาณ 60% เมื่อวิเคราะห์หาปริมาณ mitragynine ของต้นกระท่อมที่ได้จากการทดลองและต้นที่เพาะจากเมล็ด (ที่เลี้ยงในสภาวะทดลอง) อายุ 5 เดือน ด้วยวิธี TLC-densitometric method ผลการทดลองพบว่าต้นที่เจริญขึ้นใหม่มีการสะสมสาร mitragynine ในปริมาณ 14.25 ± 0.25 มก.ต่อน้ำหนักแห้ง ส่วนต้นที่เพาะจากเมล็ดมีการสะสมสาร mitragynine ในปริมาณ 4.45 ± 0.09 มก.ต่อน้ำหนักแห้ง

Thesis Title Induction of *Agrobacterium rhizogenes* Transformed Hairy Root Cultures from *Mitragyna speciosa* (Roxb.) Korth.

Author Miss Siriwan Phongprueksapattana

Major Program Pharmaceutical Sciences

Academic Year 2007

ABSTRACT

The hairy root culture of *Mitragyna speciosa* was induced with the wild type *Agrobacterium rhizogenes* ATCC 15834 from leaf vein explants. The hairy roots were identified for the rooting genes loci of *rolA* and *rolB* by the polymerase chain reaction. The results showed that the efficiency of *A. rhizogenes* genes transformation was about 60%. The transformed hairy roots appeared branching with pubescent hairs. Study on the appropriate medium for growth of the hairy roots suggested that among 9 different kinds of media, WPM supplemented with 0.5 mg/l 1-naphthalene acetic acid was suitable medium for promoting the growth. Study on secondary metabolite production, the *n*-hexane extract of the hairy roots was prepared and the compounds were isolated by silica gel column chromatography, affording the MSF1 and MSF2 with the yield of 0.016% (w/w) and 0.027% (w/w), respectively. Elucidation of their structures using ¹H- and ¹³C-NMR spectroscopic methods, indicated that MSF1 was a mixture of β -sitosterol and stigmasterol (ratio 1:1) and MSF2 was a mixture of ursolic acid and oleanolic acid (ratio 5:1). The growth curve and the ursolic acid production curve of the hairy root culture were plotted. The growth cycle of the hairy root culture was realized within 40 days. The production curve of the hairy roots indicated that ursolic acid was contained in the *M. speciosa* hairy roots and reached the maximum yield of 3.47 ± 0.03 mg/g dry wt at a late linear phase. In this study, the micropropagation of *M. speciosa* was also performed. Shoot multiplication was successfully induced from axillary buds in WPM supplemented with 0.1 mg/l thidiazuron. The plantlet regeneration was successfully acclimatized with 60% survival rate. The mitragynine content of the regenerated plant and the *in vitro* plantlets at 5-months old were determined using the TLC-densitometric method. The results showed that the regenerated plant contained mitragynine of 14.25 ± 0.25 mg/g dry wt, whereas the *in vitro* plantlet contained 4.45 ± 0.09 mg/g dry wt.

ACKNOWLEDGEMENTS

I wish to express my deepest appreciation and grateful thanks to my advisor, Assistant Professor Dr. Juraithip Wungsintaweekul for her helpful guidance, suggestion and encouragement throughout the course of this work.

My sincere thanks are expressed to my thesis co-advisor Associate Professor Dr. Waraporn Putalun, for her kindness and helpful suggestions.

I would like to extend my sincere thanks to Assistant Professor Dr. Niwat Keawpradub for his kindly provided the authentic compounds. I would like to thank Mr. Rajagopal Ragavan for english language consultant.

I would also thank the Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Prince of Songkla University for their support in scientific equipment. I would like to thank all staffs of Faculty of Pharmaceutical Sciences, Prince of Songkla University for their kindness and help.

Finally, I would to thank my friends for their love and encouragement.

Siriwan Phongprueksapattana

CONTENTS

	Page
Contents	vi
List of tables	ix
List of figures	x
List of abbreviations and symbols	xii
Chapter	
1. Introduction	1
1.1 Background and rationale	1
1.2 Review of literature	2
1.2.1 Botanical aspects of <i>Mitragyna speciosa</i> (Roxb.) Korth.	2
1.2.2 Chemical constituents of <i>M. speciosa</i>	3
1.2.3 Biological activities	11
1.2.3.1 Indole alkaloids	11
1.2.3.2 Triterpenoids : ursolic acid	11
1.2.4 Genetic transformation and plant tissue cultures of Rubiaceous plant	12
1.2.5 Biosynthesis in <i>M. speciosa</i>	16
1.2.5.1 Mitragynine biosynthesis	16
1.2.5.2 Ursolic acid biosynthesis	17
2. Materials and methods	20
2.1 Materials and Equipments	20
2.1.1 Plant material	20
2.1.2 Chemicals	20
2.1.3 Equipments	22
2.1.4 Media preparations and solutions	23
2.1.5 Plasmid isolation kits and enzymes kits	24

CONTENTS (Continued)

	Page
2.2 Methods	24
2.2.1 Preparation of <i>M. speciosa</i> plantlets	24
2.2.2 Bacterial strain and preparation	24
2.2.3 Plant tissue culture and media preparation	25
2.2.4 Induction of hairy roots	26
2.2.5 Identification of transformed hairy roots	26
2.2.6 Agarose gel electrophoresis	28
2.2.7 Medium manipulation	29
2.2.8 Isolation of secondary metabolites from transformed hairy roots	30
2.2.9 NMR spectroscopy	30
2.2.10 Establishment of the growth and production curves of <i>M. speciosa</i> hairy roots	32
2.2.11 Plant regeneration of <i>M. speciosa</i>	33
2.2.11.1 Induction of shoots and roots	33
2.2.11.2 Acclimatization	34
2.2.12 Establishment of thin layer chromatography-densitometric method	34
3. Results	37
3.1 Establishment of transformed hairy root culture of <i>M. speciosa</i>	37
3.1.1 Induction of hairy roots	37
3.1.2 Identification of <i>rolA</i> and <i>rolB</i> genes	41
3.2 Medium manipulation	43
3.3 Structure elucidation of isolated compounds from <i>M. speciosa</i> hairy root culture	48
3.3.1 MSF1	48
3.3.2 MSF2	50
3.4 Ursolic acid production in <i>M. speciosa</i> hairy roots	52
3.5 Micropropagation of <i>M. speciosa</i>	56

CONTENTS (Continued)

	Page
3.6 Mitragynine production of the regenerated <i>M. speciosa</i> plantlet	61
4. Discussion	64
5. Conclusions	67
References	69
Appendix	74
Vitae	99

LIST OF TABLES

Table	Page
1.1 Chemical constituents of <i>M. speciosa</i> (Roxb.) Korth.	4
1.2 Plant tissue cultures of the Rubiaceae plants	15
2.1 Primers used in this study	27
2.2 Standard procedure for PCR	27
3.1 Induction of hairy roots from different explants	38
3.2 Concentrations and purities of total DNA extracted from hairy root cultures	42
3.3 The appearances of the hairy roots cultures in different media	44
3.4 The dry weight of the hairy roots in different media	46
3.5 HMBC: ¹ H and ¹³ C NR spectroscopic data for ursolic acid	51
3.6 Biomass and ursolic acid contents in untransformed roots and hairy roots	54
3.7 Shoot number of plantlet regeneration in WPM medium	58
3.8 Mitragynine content	63
A-1 Inorganic salt and vitamin composition of MURASHIGE&SKOOG MEDIUM (MS)	74
A-2 Inorganic salt and vitamin composition of McCOWN WOODY PLANT	75
A-3 Inorganic salt and vitamin composition of Gamborg's B5 medium	76

LIST OF FIGURES

Figure	Page
1.1 <i>Mitragyna speciosa</i> (Roxb.) Korth. (Rubiaceae)	3
1.2 Chemical structures of indole alkaloids found in <i>M. speciosa</i>	6
1.3 Chemical structures of flavonoids, phenylpropanoids, lignan, and triterpenoid found in <i>M. speciosa</i>	9
1.4 Interaction of <i>Agrobacterium rhizogenes</i> to plant cell	13
1.5 Biosynthesis of indole alkaloids	16
1.6 Formation of squalene	18
1.7 Biosynthesis of ursolic acid	19
2.1 <i>M. speciosa</i> (Roxb.) Korth.: the globular fruit	20
2.2 Scheme of isolation of compounds	31
2.3 Scheme of preparation of the crude alkaloid extract	36
3.1 <i>M. speciosa</i> grown in WPM solid medium supplemented with BA 1.0 mg/l	37
3.2 Steps of the hairy roots induction from <i>M. speciosa</i> plantlets	38
3.3 Characteristics of hairy roots, induced from leaf veins and stems	39
3.4 Characteristic of hairy roots culture in WPM solid medium	40
3.5 Characteristics of transformed hairy roots and untransformed roots	40
3.6 1.2% Agarose gel electrophoresis	42
3.7 Growth curves of hairy roots in selected media	47
3.8 Relative growth of hairy roots in selected media	47
3.9 Structures of β -sitosterol and stigmasterol	49
3.10 Structures of ursolic acid and oleanolic acid	51
3.11 Calibration curves of ursolic acid	52
3.12 HPLC-chromatograms of ursolic acid	53
3.13 Growth curve and production curve of hairy root culture	55
3.14 Growth curve and production curve of untransformed root culture	55
3.15 Shoot regeneration from <i>M. speciosa</i> hairy root culture	56

LIST OF FIGURES (Continued)

Figure	Page
3.16 Plant regeneration from the <i>M. speciosa</i> hairy root culture	57
3.17 Shoot multiplication of the axillary buds	58
3.18 Root initiation of the regenerated plantlets of <i>M. speciosa</i> in WPM medium	59
3.19 Acclimatization protocol for <i>M. speciosa</i> regenerated plant	60
3.20 Calibration curve of authentic mitragynine	61
3.21 TLC densito-chromatograms and UV absorption	62
4.1 Proposed biosynthetic pathway of mitragynine	65
B-1 ¹ H NMR spectrum of mixture of MSF1 in CDCl ₃	79
B-2 ¹³ C NMR spectrum of mixture of MSF1 in CDCl ₃	80
B-3 ¹ H NMR spectrum of MSF2 in CDCl ₃ :CD ₃ OD	88
B-4 ¹³ C NMR spectrum of MSF2 in CDCl ₃ :CD ₃ OD	89
B-5 DEPT 90 of MSF2 in CDCl ₃ :CD ₃ OD	94
B-6 DEPT 135 of MSF2 in CDCl ₃ :CD ₃ OD	95
B-7 COSY of MSF2 in CDCl ₃ :CD ₃ OD	96
B-8 HMQC spectrum of MSF2 in CDCl ₃ :CD ₃ OD	97
B-9 HMBC spectrum of MSF2 in CDCl ₃ :CD ₃ OD	98

LIST OF ABBREVIATIONS AND SYMBOLS

bp	=	base pair
B5	=	Gamborg's B5 medium
BA	=	benzyladenine
DNA	=	deoxyribonucleic acid
dNTP	=	deoxynucleoside triphosphate
g	=	gram
h	=	hour
HPLC	=	high performance liquid chromatography
NAA	=	1-naphthalene acetic acid
mg	=	milligram
min	=	minute
ml	=	mililiter
MS	=	Murashige-Skoog (MS) medium
WPM	=	McCown Woody plant medium
µg	=	microgram
µl	=	microliter
OD	=	optical density
pH	=	-log hydrogen ion concentration
rpm	=	round per minute
PCR	=	polymerase chain reaction
TLC	=	thin layer chromatography
TDZ	=	thidiazuron
T _m	=	melting temperature
fig.	=	figure
UV	=	ultraviolet
v/v	=	volume by volume
w/v	=	weight by volume

LIST OF ABBREVIATIONS AND SYMBOLS (Continued)

w/w	=	weight by weight
YEB	=	yeast extract broth
°C	=	degree celsius

CHAPTER 1

INTRODUCTION

1.1 Background and rationale

Mitragyna speciosa (Roxb.) Korth. (Rubiaceae) is an endemic plant found in tropical Southeast Asia. It is of particular medicinal importance known as “Kratom” in Thailand, “Biak-Biak” in Malaysia (เต็ม สมิตินันท์, 2544). For folklore medicine, Kratom has been used as an opium-substitute for pain relief and treatment of diarrhea. Many studies reported the wide variety of indole alkaloid from Kratom’s leaves including mitragynine, paynantheine, rhynchophylline, etc. (see Table 1.1). Moreover, it also contains flavone, flavonol, flavonoid, lignan, phenylpropanoid and triterpenoid (Information from NAPRALERT database). The pharmacological activities of mitragynine and its derivatives have been reported from studies in tested animals such as analgesic activity, antinociceptive activity (Watanabe *et al.*, 1997), anti-stress activity, muscle relaxant activity (Aji *et al.*, 2001) and inhibition of gastric acid secretion (Tsuchiya *et al.*, 2002). The mechanism of action of mitragynine for analgesic activity is binding to the opioid receptors, similar to morphine. Interestingly, mitragynine has a characteristic of less addiction than morphine. Therefore, it highlights the relevance to be an alternative to opioid analgesic drug (Thongpraditchote *et al.* 1998).

Contrary to the potential of *M. speciosa*, it is recognized as an illegal plant and growing of this plant is prohibited in Thailand. As such the basic knowledge of mitragynine biosynthesis is very little. From the biosynthesis point of view, the steps of mitragynine formation are still unknown. As part of our interest on plant biosynthesis, the plant cell cultures, such as callus and suspension cultures are established in order to use as a model instead of the intact plant. For those reasons the model plant is controlled under specific conditions. Therefore, the chemicals produced by the plant cell culture do not change by seasons. However, the disorganized cells, sometimes, do not produce the secondary metabolites as found in the intact plant. Techniques of medium manipulation and elicitation then are used to stimulate the disorganized

cells growing, which are able to produce the secondary metabolites. On the other hand, establishing the organ culture is probably making cells complex enough for production of secondary metabolites. One of the techniques widely used for inducing the organ culture is the genetic transformation of *Agrobacterium* to plant cells. For instance, *Agrobacterium rhizogenes*, the causative agent of hairy root disease in plants, was transformed into plant cells and later forming the hairy root culture. This technique has emerged as an important method for secondary metabolic production. An advantage of using transformed hairy root culture is practically scaling up in the bioreactor for commercial purposes. For the sources of indole alkaloid production, hairy roots of *Catharanthus roseus*, *Rauwolfia serpentina* and *R. micrantha* have been reported to accumulate medicinal compounds such as ajmaline, ajmalicine and serpentine (Sim *et al.*, 1994, Benjamin *et al.*, 1994, Sudha *et al.*, 2003). These cultures have been currently used in the biosynthetic studies of indole alkaloid.

In this study, we aim to induce the hairy root culture of *M. speciosa* with a wild type *Agrobacterium rhizogenes* ATCC 15834. The potential of secondary metabolites production was evaluated. Then, secondary metabolites accumulated in the *M. speciosa* hairy roots are isolated and elucidated for their structures by NMR spectroscopy. In addition, we also report the protocol of micropropagation of *M. speciosa*. The regenerated plantlets are evaluated for mitragynine accumulation using TLC-densitometric method. The hairy root culture and the regenerated plantlets of *M. speciosa* may be good models for the study of mitragynine biosynthesis.

1.2 Review of literature

1.2.1 Botanical aspects of *Mitragyna speciosa* (Roxb.) Korth.

Mitragyna speciosa (Roxb.) Korth. (Fig. 1.1) belongs to the Rubiaceae family, found in the regions of Africa and Asia. The genus, named *Mitragyna* was given by Korthals due to the shape of the stigmas in the species. However the nomenclature has frequently been confused; the genres have been variously named and are consistently recognized as *Naucleaeae*, *Sarcocephalus*, *Stephegyne* and *Uncaria*. In Thailand, there are four species of *Mitragyna*, which

are *M. speciosa*, *M. hirtusa*, *M. diversifolia* and *M. rotundifolia* (เต็ม สมิตินันท์, 2544). Taxonomically, plant species can be categorized by types of indole alkaloids (Keawpradub, 1990). Recently, these species were authenticated by molecular approach, based on rDNA ITS sequence (Sukrong *et al.*, 2007).

For botanical aspects, *M. speciosa* is characterized by the globular flowering head each containing up to 120 florets (Fig. 1.1 B). Each floret is surrounded by many overlapping bracteoles which completely cover the developing florets during the flower-bud stage. The inflorescence is a dichasial cyme. The fruit is a capsule containing numerous small flat seeds. The young woody shoots bear 10-12 leaves arranged in opposite and decussate each pair of leaves being accompanied by two interpetiolar stipules (Fig. 1.1 A) which initially are closely appressed and protect the apical bud.




Figure 1.1 *Mitragyna speciosa* (Roxb.) Korth. (Rubiaceae) (A); Globular flowering head (B).

1.2.2 Chemical constituents of *M. speciosa*

Leaves of *M. speciosa*, found in Thailand and Malaysia, contain common indole alkaloids. Takayama (2004) reviewed the abundance of mitragynine in crude alkaloidal extract of Thai and Malay strains. It has been reported that the *M. speciosa* of Thai contained mitragynine up to 66% of the total alkaloid content, whereas only 12% was found in the *M. speciosa* of Malay sample. Reviews of the distribution of indole alkaloids in leaves, twigs, stem barks and root barks

of *M. speciosa* are summarized in Table 1.1. Chemical structures of selected indole alkaloids were shown in Fig 1.2 and of other components were shown in Fig 1.3.

Table 1.1 Chemical constituents of *M. speciosa* (Roxb.) Korth.

<i>Plant part</i>	<i>Category</i>	<i>Chemical substance</i>	<i>Reference</i>
leaves	alkaloid	ajmalicine, akuammigine, angustine, corynantheidine, corynantheidaline, corynantheidalinic acid, corynoxine, corynoxine B, hirsutine, hirsuteine, isocorynoxine, isomitraphylline, isorhynchophylline, isocorynantheidine, javaphylline, mitraciliatine, mitragynine oxindole B, mitrajavine, mitraphylline, mitrasulgynine, mitragynaline, mitragynalinic acid, mitralactonal, paynantheine, mitragynine, pinoresinol, speciociliatine, speciogynine, 3-isoajmalicine, 3,4,5,6-tetrahydromitragynine, 7 α -hydroxy-7H-mitragynine	Phillipson <i>et al.</i> , 1973; Phillipson <i>et al.</i> , 1973; Shellard <i>et al.</i> , 1966; Shellard <i>et al.</i> , 1978a; Shellard <i>et al.</i> , 1978b Takayama, 2004
	flavones	apigenin, apigenin-7-O-rhamnoglucoside, cosmosiin	Hinou and Harvala, 1988
	flavonol	astragalinalin, hyperoside, kaempferol, quercetin, quercitrin, quercetin-3-galactoside-7-rhamnoside, quercitrin, rutin	Harvala and Hinou, 1988

Table 1.1 (continued)

<i>Plant part</i>	<i>Category</i>	<i>Chemical substance</i>	<i>Reference</i>
leaves	phenylpropanoid	caffeic acid, chlorogenic acid	Hinou and Harvala, 1988
	flavonoid	(-)-epicatechin	Houghton and Said, 1986
	lignin	(+)-pinoresinol	Takayama <i>et al.</i> , 1998
	triterpene	ursolic acid	Said <i>et al.</i> , 1991
young twigs, stem bark	alkaloid	ciliaphylline, rhynchociline, ciliaphylline, isomitraphylline, isorhynchophylline, isospecionoxeine, javaphylline, mitraciliatine, mitragynine oxindole A, mitragynine oxindole B, mitraphylline, rhynchociline, rhynchophylline, speciogynine, speciociliatine, specionoxeine	Shellard <i>et al.</i> , 1978a; Shellard <i>et al.</i> , 1978b
root bark	alkaloid	ciliaphylline, corynoxine, isocorynoxine, isomitraphylline, isorhynchophylline, isospecionoxeine, mitraciliatine, mitraphylline, rhynchociline, rhynchophylline, speciociliatine, speciogynine, specionoxeine	Shellard <i>et al.</i> , 1978b; Houghton and Shellard, 1974







Figure 1.2 Chemical structures of indole alkaloids found in *M. speciosa*




corynantheidalinic acid




mitralactonal




mitraciliatine




ajmalicine




mitraphylline



isomitraphylline




isorhynchophylline




hirsutine


Figure 1.2 (Continued)




hirsuteine




corynoxine B



corynoxine



corynoxine



isocorynoxine

Figure 1.2 (Continued)




Figure 1.3 Chemical structures of flavonoids, phenylpropanoids, lignan, and triterpenoid found in *M. speciosa*




Figure 1.3 (Continued)

1.2.3 Biological activities

1.2.3.1 Indole alkaloids

As mentioned earlier, *M. speciosa* is a major source of indole alkaloids. Mitragynine is the most abundant component that is present in the leaves of *M. speciosa*. For biological activities of alkaloids obtained from *M. speciosa*, the crude extract and pure alkaloid-mitragynine were used as materials *in vitro* and *in vivo* experiments. In 1996, a Japanese group investigated the antinociception of mitragynine in mice and the results showed that mitragynine acted as opioid receptors agonist in the brain of mice (Matsumoto *et al.*, 1996a). Later, they found that the mechanisms for antinociceptive effects differed from those of morphine in mice (Matsumoto *et al.*, 1996b). Mitragynine preferred to bind μ and δ opioid receptors, unlike morphine that binds only to μ opioid receptor specifically. Therefore, mitragynine exhibits 10 fold less potent than morphine (Thongpraditchote *et al.*, 1998). Caused by the affinity to opioid receptor, mitragynine also acts as morphine-like action on inhibition of gastric secretion (Tsuchiya *et al.*, 2002). Studies on the synthesis of mitragynine-related indole alkaloids discovered mitragynine pseudoindoxyl and 7-hydroxymitragynine, which exhibited opioid agonistic activity with higher potency than morphine (Takayama *et al.*, 2002). With regards to the psychological effects, *M. speciosa* extract was known to have a stimulatory effect on the dorsal raphe nucleus and an antidepressant-like activity. Stimulation of this brain area has been known to cause antinociception (Kumarnsit *et al.*, 2007).

1.2.3.2 Triterpenoids : ursolic acid

Ursolic acid was found in *M. speciosa* leaves (Said *et al.*, 1991). To date, there is no report about the biological activity of ursolic acid from *M. speciosa*. However, ursolic acid normally exists in plants, especially the Lamiaceae family. Contemporary scientific researches revealed and confirmed that ursolic acid exhibited several pharmacological effects, such as antitumour, hepatoprotective, anti-inflammatory, anti-ulcer, antimicrobial, anti-hyperlipidemic and antiviral activity (Liu, 1995). Ursolic acid was also identified as one of the active principles in

Rosmarinus officinalis (Lamiaceae) to inhibit the growth of some food borne pathogens (Kamatou *et al.*, 2007). Somova *et al.*, 2003 found that African wild olives contain oleuafrecein (mixture of oleanolic and ursolic acid) exhibiting the antihypertensive, diuretic/natriuretic, antihyperlipidemic, hypoglycemic and antioxidant activities. Ursolic acid from *Bouchea fluminensis* has anti-inflammatory activity inhibiting significantly the edema induced by either carrageenan or serotonin (Costa *et al.*, 2003). The anti-inflammatory effects of ursolic acid have been attributed to various mechanisms including inhibition of lipoxygenase and cyclooxygenase activities, inhibition of elastase and inhibition of C3-convertase (Costa *et al.*, 2003).

1.2.4 Genetic transformation and plant tissue culture of Rubiaceae plant

The most popular bacteria, used for genetic transformations, are *Agrobacterium tumefaciens* and *A. rhizogenes*. *A. tumefaciens* contains tumor-inducing plasmid (Ti-plasmid), therefore, causing the crown-gall tumor to plant cells, whereas *A. rhizogenes* contains root-inducing (Ri-plasmid), causing the hairy roots (Oksman-Caldentey and Hiltunen, 1996). In this study, we used *A. rhizogenes* for genetic transformation to *M. speciosa* cells in order to induce the hairy root cultures.

A. rhizogenes is a gram negative soil bacterium and is supplied in many varieties such as wild-type strain-ATCC 15834, A4 (Sevón and Oksman-Caldentey, 2002). *A. rhizogenes* carries the Ri plasmid, its size about 200 to greater than 800 kb. It contains T-DNA and a virulence (*vir*) region, responsible for induction of hairy root formation (Oksman-Caldentey and Hiltunen, 1996) The Ri-plasmid carries the borders of T_L and T_R (or LB and RB), which are expressed after stimulation of virulence (*vir*) region. The *vir* region is about 35 kb and encodes six transcriptionals: *vir* A, B, C, D, E and G, which have important functions in gene transfer.

Fig. 1.4 illustrates the pathogenic responses of the interaction between *A. rhizogenes* and plant cells. After plant cells are wounded, they then release the phenolic compounds e.g. acetosyringone, which activates the genes of *vir*-region of the bacterial plasmid. Finally, the T-DNA is cleaved (at the region of LB and RB), transferred to plant cells and later incorporated into the nuclear DNA of host plant (Sevón and Oksman-Caldentey, 2002). The

resulting genetic-transformed plant cells synthesize novel metabolites e.g. opine, which are essential nutrients for auxin production (Oksman-Caldentey and Hiltunen, 1996).

Mechanism of *Agrobacterium*-plant cell interaction




Figure 1.4 Interaction of *Agrobacterium rhizogenes* to plant cell.

In conclusion, the plasmid itself has three regions, those are important for the gene transfer, which are 1) the T-DNA, which is integrated into the genome of the plant cells; 2) the border sequences of the T-DNA and 3) the virulence area (*vir*), which does not enter the plant cell but causes the transfer of the T-DNA to the plant cells (Oksman-Caldentey and Hiltunen, 1996).

A. rhizogenes Ri-plasmid transformed to plant cells causes the hairy roots and the transformed cells are more sensitive to extracellular supplied auxins than the normal roots (Sevón *et al.*, 1998). Susceptibility of plant species to *Agrobacterium* strains varies greatly. It is dependent upon factors, which are 1) transformation ability of different strains of *Agrobacterium*; 2) age and differentiation status of plant tissues; 3) presence of phenolic compounds e.g. addition of acetosyringone. For identification of gene-loci of Ri-plasmid, the genetic transformation can be confirmed by assays: 1) detection of opine production by paper electrophoresis; 2) detection of T-DNA by Southern blot hybridization and 3) detection of genes in *A. rhizogenes* by polymerase

chain reaction (PCR) (Oksman-Caldentey and Hiltunen, 1996). The hairy roots are morphologically characterized by laterally highly branched roots, fast growing, and are able to grow in hormone-free medium. They are genetically stable and produce high contents of secondary metabolites. The ability of secondary metabolite production of hairy roots is highly linked to cell differentiation.

For the studies on plant tissue culture of the Rubiaceae plants, they reported the establishment of callus, cell suspension and organ cultures under specific conditions. Table 1.2 summarized the plant tissue cultures of the Rubiaceae plants. It can be noted that they were able to produce the alkaloids and triterpenoids. An example of the hairy roots of *Ophiorrhiza pumila* convinced the utility of hairy root culture for camptothecin production in the industrial scale (Saito *et al.*, 2007). Suspension culture of *Uncaria tomentosa* could produce ursolic acid and oleanolic acid but no alkaloid was present in the culture (Feria Romero *et al.* 2005).

Table 1.2 Plant tissue cultures of the Rubiaceous plants

<i>Plant source</i>	<i>Chemicals</i>	<i>Culture conditions</i>	<i>Reference</i>
<i>Cinchona ledgerina</i>	cinchonine, quinidine, quinine	suspension culture : Gamborg B5 basal medium containing 2,4-dichlorophenoxyacetic acid 1 mg/l and saccharose 20 mg/l	Hamill <i>et al.</i> , 1989
<i>Ophiorrhiza pumila</i>	Camptothecin	hairy root culture : Gamborg B5 medium containing sucrose 2%(w/v)	Saito <i>et al.</i> , 2007
<i>Psychotria umbellata</i>	umbellatine	embryogenic regeneration : Murashige and Skoog medium containing 1-naphthaleneacetic acid 1 mg/l and kinetin 1 mg/l	Paranhos <i>et al.</i> , 2005
<i>Rudgea jasminoides</i>	-	callus cell culture : Murashige and Skoog medium containing kinetin, 2.22 μ M and picloram 2.07 μ M	Stella <i>et al.</i> , 2002
<i>Uncaria rhynchopylla</i>	hirsuteine, hirsutine, 3 α -dihydrocadambine, ursolic acid	callus culture : Gamborg B5 containing indole acetic acid 10^{-4} M and benzyladenine 3×10^{-5} M and sucrose 2% (w/v)	Kohda <i>et al.</i> , 1996
<i>U. tomentosa</i>	ursolic acid, oleanolic acid	suspension culture : Nitsch & Nitsch medium containing 2,4-dichlorophenoxyacetic acid 2 mg/l, kinetin 2 mg/l and sucrose 20 g/l	Feria Romero <i>et al.</i> , 2005

1.2.5 Biosynthesis in *M. speciosa*

1.2.5.1 Mitragynine biosynthesis

Mitragynine is corynanthe-type indole alkaloid. Generally, the indole nucleus is obtained from the condensation of one molecule of tryptamine, decarboxylation of tryptophan and one molecule of secologanin, iridoid moiety. Tryptamine is supplied from the shikimate pathway, whereas secologanin originated from terpenoids pathway. Therefore, the alkaloid group is named as “monoterpenoid-indole alkaloids” (TIAs). Condensation of tryptamine and secologanin is catalyzed by strictosidine synthase, forming the first intermediate strictosidine. (Mizukami *et al.*, 1979). Strictosidine serves as common precursor for all indole alkaloids. Distance between strictosidine and mitragynine is still unknown. Fig. 1.5 shows the condensation reaction forming the general precursor of indole alkaloids.




Figure 1.5 Biosynthesis of indole alkaloids. Dashed arrows indicate multi- step reactions.

1.2.5.2 Ursolic acid biosynthesis

Ursolic acid is the triterpene acid that is biosynthesized from six molecules of isoprene units. The two molecules of farnesyl diphosphate are joined tail to tail yielding the hydrocarbon squalene (Fig. 1.6). Then squalene is cyclized to polycyclic triterpene via the intermediate squalene-2,3-oxide (oxidosqualene) by a flavoprotein requiring O_2 and NADPH as cofactors. The methyls and hydrides groups are migrated by the sequences of Wagner-Meerwein rearrangements (Dewick, 2002). After a pentacyclic ring system is formed, giving a new five-membered ring and a lupenyl cation, the 1,2-alkyl shift is then occurred to form the olenyl cation, which is a precursor of oleanolic acid. The lupenyl cation is a branch point to taraxasteryl cation, a precursor of ursolic acid (Dewick, 2002). A five-membered ring is not highly strained as evidenced by all the examples encountered. The cyclizations and Wagner-Meerwein rearrangements appear to be catalyzed by a single enzyme, which converts squalene oxide into the final product, e.g. ursolic acid or oleanolic acid (Fig. 1.7).




Figure 1.6 Formation of squalene




Figure 1.7 Biosynthesis of ursolic acid

CHAPTER 2

MATERIALS AND METHODS

2.1 Materials and equipments

2.1.1 Plant material

Mature plants of *M. speciosa* (6 years-old), grown in the open field in Hat Yai district, Songkhla, Thailand, were used in this study. A voucher specimen of this plant was deposited in the Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat Yai campus, Songkhla, Thailand. Seeds were collected from the globular fruit (Fig. 2.1A) and flat seeds (Fig. 2.1B) were separated. These are used for plantlet preparation and further for genetic transformation with *A. rhizogenes* ATCC 15834.




Figure 2.1 *M. speciosa* (Roxb.) Korth.: the globular fruits (A); flat seeds (B)

2.1.2 Chemicals

The authentic mitragynine was kindly provided by Assist. Prof. Dr. Niwat Keawpradub, Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Prince of Songkla University. Chemicals used in this study are listed below.

<i>Chemicals</i>	<i>Company, Country</i>
acetic acid, glacial	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
agarose gel	Research Organics, USA
agar (Bacto)	Himedia laboratories, Ltd., India
amberlite XAD-4	Sigma, USA
cefotaxime disodium	M&H Manufacturing, Thailand
chloroform	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
clorox [®]	Clorox, USA
DNA markers	Sib-enzyme, Russia 2-Log DNA Ladder (0.1-10.0 kb) NEB (New England Biolabs), UK
ethanol (95%v/v)	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
ethidium bromide	Bio Basic INC, Canada
ethyl acetate	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
Gamborge's B5 medium	Duchefa Biochemie , The Netherlands
<i>n</i> -hexane	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
McCown Woody plant (WPM) medium	Duchefa Biochemie, The Netherlands
Murashige-Skoog (MS) medium	Duchefa Biochemie, The Netherlands
methanol	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
NH ₃ 25%(v/v)	BDH AnalaR [®] , England
petroleum ether	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
plant agar	Duchefa Biochemie, The Netherlands
sucrose	MITR PHOL, Thailand
thidiazuron	Sigma, USA

Chemicals (continued)

<i>Chemicals</i>	<i>Company, Country</i>
TLC-plate silica gel GF ₂₅₄	Merck, Germany
ursolic acid	Sigma, USA
yeast extract	Lab-scan Asia Co., Ltd., Bangkok, Thailand.
6-benzyladenine (BA)	Fluka, Switzerland
1-naphthaleneacetic acid (NAA)	Fluka, Switzerland
1-propanol	Lab-scan Asia Co., Ltd., Bangkok, Thailand.

2.1.3 Equipments

<i>Equipments</i>	<i>Company</i>
autoclave	Model HA-3D, Hirayama, Japan
balances	Explorer, Ohaus, USA; Avery Berkel, USA
centrifuge	Hermle Z 323 K, Germany
electrophoresis chamber	Mupid α Mini electrophoresis system, Japan
gel documentation	Gel Doc model 1000, BIO-RAD, USA Molecular Analyst [®] Software, Windows Software for BioRad's Image Analysis Systems Version 1.4
gene amplifier	GeneAmp, PCR system 9600, Perkin Elmer, USA
hot air oven	Memmert, Germany
hot plate and stirrer	Model PC-101, CORNING; Fisher Scientific, USA
incubating block	Thermomixer comfort, Eppendorf, Germany
laminar air flow cabinet	Holten, Denmark
micropipette	Socorex, Switzerland
microwave oven	LG, Thailand
pH meter	pH meter Model 710A, ORION Research, USA
platform shaker	Innova 2300, USA

<i>Equipments</i>	<i>Company</i>
refrigerator	4°C: Sanden Intercool, Thailand; -20°C: Whirlpool, Thailand; -80°C, Forma Scientific, USA
rotary evaporator	aspirator A-3S, EYELA, Japan
rotary shaking	orbital Shaker OS-20
UV-VIS spectrophotometer	Labomed, USA
TLC-scanner	TLC SCANNER 3 and software cats version 4.01, CAMAG, Switzerland
UV-CABINET II for 254, 366 nm	CAMAG, USA
vortex	Vortex-Genie 2 TM , USA
water bath	Digital water bath SB-1000, EYELA, Japan

2.1.4 Media preparations and solutions

Anisaldehyde spray reagent	Anisaldehyde (5 ml) was mixed with 10 ml of acetic acid and 5ml of HCl (37%), and then adjusted the volume to 100 ml with distilled water.
Cefotaxime stock solution (1g/5 ml)	Cefotaxime Na (1 g) was dissolved in 5 ml of sterile distilled water. The solution was sterilized using 0.22 µm membrane.
Dragendorff's spray reagent	<u>Solution A</u> : bismuth subnitrate (850 mg) was dissolved in 40 ml of distilled water and 10 ml of glacial acetic acid. <u>Solution B</u> : potassium iodide (8 g) was dissolved in 20 ml of distilled water. The mixture was prepared mixing by 5 ml of solution A and 5 ml of solution B. Then 20 ml of glacial acetic acid was added and the volume was adjusted to 100 ml with distilled water.
Ethidium bromide solution	Ethidium bromide (10 µl) was mixed in 100 ml of distilled water.

Loading buffer	Loading buffer contains glycerol (30% v/v) and bromophenol blue (0.025% w/v).
TAE buffer, 50x	The concentrated TAE (Tris-acetic acid-EDTA) buffer composed of Tris base (121 g), EDTA.3Na (19.7 g), glacial acetic acid (35ml), which were dissolved in distilled water, pH was adjusted with HCl (37%) to 8.0 and the volume was adjusted to 500 ml with distilled water.
YEB medium	The mixture contained beef extract (5 g), yeast extract (1 g), peptone (5 g), sucrose (5 g) and 10% w/v MgSO ₄ (5 ml) and distilled water. The volume was adjusted to 1 L with distilled water. For YEB solid medium, 1.5 g Bacto-agar was added to 100 ml of mixture. The culture medium was sterilized by autoclaving at 121°C, 15 lb/in ² , for 20 minutes.

2.1.5 Plasmid isolation kits and enzyme kits

GFX Micro Plasmid Prep Kit (GE Healthcare, USA)

DNeasy Plant Mini Kit (Qiagen, Germany)

ThermoPol *taq* polymerase (NEB (New England Biolabs), USA)

2.2 Methods

2.2.1 Preparation of *M. speciosa* plantlets

Flat seeds (2.1.1) of *M. speciosa* were surface sterilized by rinsing with 70% (v/v) ethanol for 5 min, then sterilizing with 20% (v/v) Clorox[®] for 5 min and finally rinsing thoroughly with sterile distilled water (x3). Sterilized seeds were placed on WPM solid medium supplemented with BA (1.0 mg/l) for seed germination. After incubating at 25°C and under light

for 16 h/day for 1 week, the seedlings germinated. The 2-month old plantlets were the explants for bacterial infection.

2.2.2 Bacterial strain and preparation

Agrobacterium rhizogenes wild-type strain ATCC 15834 (TISTR 1450) was obtained from the Microbiological Resources Centre, Pathumthani, Thailand. *A. rhizogenes* ATCC 15834 was grown and maintained in YEB medium. Before bacterial infection, *A. rhizogenes* was activated by culturing on YEB solid medium and incubated at 28°C for overnight (16 h). A single bacterial colony was obtained and used for inoculation into 5 ml YEB liquid medium. The culture was then placed on rotary shaker (218 rpm) and incubated at 28°C overnight. The bacterial suspension was harvested by centrifugation at 3,500 rpm for 10 min. The pellet was then re-suspended. The bacterial concentration was adjusted with sterile WPM liquid medium to obtain an OD₆₀₀ about 0.5-0.6. Finally, the bacterial suspension was further used to infect the explants using needle injection method (Dhakulkar *et al.*, 2005).

2.2.3 Plant tissue culture media preparations

The composition of the MS, WPM and B5 media (Duchefa Biochemie) are shown in table A-1, table A-2 and table A-3, respectively (see Appendix). For MS medium preparation, the MS powder (4.4 g) and sucrose (30 g) were dissolved in distilled water. The pH of the solution was adjusted with 1 N NaOH or 1 N HCl to 5.8. For WPM medium preparation, the McCown WPM powder (2.5 g) and sucrose (20 g) were dissolved in distilled water in the final volume of 1 L. The pH of WPM medium was adjusted to 5.7. For B5 medium preparation, the B5 powder (3.2 g) and sucrose (20 g) were dissolved in distilled water; volume was adjusted to 1 L. The pH of B5 medium was adjusted to 5.5. Plant growth regulator was added as needed. For solid medium preparation, plant agar was added to the final concentration of 0.8% (w/v), and the mixture was heated gently with continuous stirring until a clear solution was obtained. All kinds of culture media were sterilized by autoclaving at 121°C, 15 lb/in² ch, for 15 minutes.

2.2.4 Induction of hairy roots

For hairy roots induction, explants (young stems and leaves) were infected with *A. rhizogenes* (2.2.2). Firstly, the explants were wounded with needle and then the wounded explants were submerged in bacterial suspension for further 30 min. The excess bacterial suspension was washed thoroughly with sterile distilled water for 3 times. After removing the water with sterile filtered paper, the infected explants were transferred to hormone-free WPM solid medium and incubated the culture at 25 °C under darkness. To kill the residual *A. rhizogenes*, those explants were transferred to WPM solid containing 500 mg/l of cefotaxime after infection for 3 days. The cultures were further incubated for 1 week. Then, the cefotaxime concentration was reduced each week from 500, 250, 100 mg/l, respectively. Finally the cultures, free from *A. rhizogenes*, were transferred to WPM solid medium without plant growth regulators. For the controlled root culture, the untransformed root culture was induced from the adventitious roots of plantlets. Adventitious roots were excised from 2-month old plantlets and cultured in WPM liquid medium. All cultures were incubated at 25°C, on rotary shaker (80 rpm) and under dark condition. The hairy root culture and untransformed root culture were sub-cultured into fresh medium every 2 months. For large scale production, the hairy roots and untransformed roots were sub-cultured to 200 ml of WPM medium in 1 L Erlenmeyer flask.

2.2.5 Identification of transformed hairy roots

The hairy root culture was identified for the genetic transformation from *A. rhizogenes*. The presence of the virulence genes of *rolA* and *rolB* were detected using the specific primers (Table 2.1), which were designed from the nucleotide sequence of T-DNA region of *A. rhizogenes* (GenBank) (Oksman-Caldentey and Hiltunen, 1996). The DNA fragment was amplified using polymerase chain reaction (PCR). The genomic DNA of the hairy roots was isolated using the DNeasy Plant Mini Kit (Qiagen), which was used as the DNA template in the PCR reaction. For negative control reaction, the DNA template was the genomic DNA of untransformed roots, which were isolated using the same procedure. For positive controlled reaction, the DNA template was the *A. rhizogenes* plasmid, which was isolated from the *A.*

rhizogenes cells using the GFX Micro Plasmid Prep Kit (GE Bioscience). The protocols of plant genomic DNA and bacterial plasmid isolation were summarized in the Appendix. PCR reactions were performed for the detection of a 300 base pairs fragment of the *rolA* gene and a 780 base pairs fragment of the *rolB* gene amplifications. The PCR reaction was performed in 50 μ l of mixture (Table 2.2) and the PCR conditions were summarized as shown below.

Table 2.1 Primers used in this study

<i>Name of sequence</i>	<i>Nucleotide sequence (5' to 3')</i>	<i>T_m (°C)</i>
<i>RolA-1</i>	CAGAATGGAATTAGCCGGACTA	60.8
<i>RolA-2</i>	CGTATTAATCCCGTAGGTTTGTTT	59.4
<i>RolB-1</i>	ATGGATCCCAAATTGCTATTCCTTCCACGA	66.0
<i>RolB-2</i>	TTAGGCTTCTTTCTTCAGGTTTACTGCAGC	66.0

Table 2.2 Standard procedure for PCR

<i>Reagents</i>	<i>Volume/reaction (μl)</i>	<i>Final concentration</i>
DNA Template	5	ca. 6 pg
dNTP mix, 10 mM each	1	200 μ M of each dNTP
10 μ M primer 1	2.5	0.5 μ M
10 μ M primer 2	2.5	0.5 μ M
10X ThermoPol Buffer	5	1x
<i>Taq</i> DNA polymerase (NEB)	0.5	2.5 units/reaction
Sterilize distilled water	33.5	
Total volume	50	

PCR conditions for *rolA* amplification

<i>Segment</i>	<i>Step</i>	<i>Temperature</i> (°C)	<i>Time</i> (min)	Number of cycles
1	Denaturing	94	2	1
2	Denaturing	94	2	1
	Annealing	58	2	
3	Extension	72	2	24
	Denaturing	94	1	
	Annealing	58	1	
4	Extension	72	1	1
	Extension	72	10	
	Holding	4	∞	

PCR conditions for *rolB* amplification

<i>Segment</i>	<i>Step</i>	<i>Temperature</i> (°C)	<i>Time</i> (min)	Number of cycles
1	Denaturing	94	3	1
2	Denaturing	94	1	30
	Annealing	55	1	
3	Extension	72	3	1
	Extension	72	10	
	Holding	4	∞	

2.2.6 Agarose gel electrophoresis

To determine the size of the DNA fragments obtained from the PCR reactions, agarose gel (1.2% (w/v)), was prepared. Agarose (0.48 g) and TAE buffer (50x) were mixed with 40 ml of distilled water. The mixture was heated using the microwave oven until a clear solution was obtained. Then, the mixture was poured on a plastic tray, and a comb was placed to mold the wells for sample applications. After the agarose gel was completely set (30-45 minutes at room

temperature), the comb was carefully removed and the gel was installed on the platform in the electrophoresis tank containing 1x TAE buffer (dilution from TAE buffer (50x) with distilled water). The PCR products (10 μ l) including loading buffer (1 μ l) were loaded into the wells of 1.2% w/v agarose gel using micropipette. The gel was run at 100 volts. The electrophoresis process was stopped when the loading dye had migrated to two third of the agarose gel (about 30 min). After that, the agarose gel was stained with ethidium bromide solution for 10 minutes and the DNA patterns were observed under UV 312 nm (a UV light box equipped with Gel Doc model 1000, BIO-RAD, USA) and a photograph was taken.

2.2.7 Medium manipulation

After obtaining the positive clones, the hairy root cultures (clone no.4) were grown in different liquid media in order to optimize the growth media for the hairy root cultures. In order to increase the growth of the hairy root cultures, the culture media were manipulated as well as the plant growth regulator-NAA was added in the final concentration of 0.5 mg/l. The manipulated culture media were WPM, MS, $\frac{1}{2}$ WPM, $\frac{1}{2}$ MS, $\frac{1}{2}$ B5, WPM plus NAA, MS plus NAA, B5 plus NAA, and $\frac{1}{2}$ B5 plus NAA. The control cultures were the untransformed roots in WPM and WPM plus NAA. The culture conditions were at 25°C on rotary shaker (80 rpm) and under dark condition. The morphology of cultures was observed after sub-culture for 1 month. The growth of cultures was evaluated by weighing the dry weight. The dry weight was obtained from weight of the hairy roots, which were dried at 50°C overnight in hot air oven. The time course of growth was also established for the period of 40 days of culture. The samples (n=3) were consecutively taken every five days of culture. Only the hairy roots cultures grown in WPM, WPM plus NAA, MS plus NAA, B5 plus NAA and $\frac{1}{2}$ B5 plus NAA were used to construct the growth curves. To decrease the error of the inoculation size, the growth curve of the hairy roots cultures was constructed by plotting the relative growth (in comparison to day 5th of culture) against the age of culture.

2.2.8 Isolation of secondary metabolites from transformed hairy root culture

The potential of secondary metabolites production by the hairy roots culture was evaluated. Mass preparation of hairy roots was prepared (2.2.4). The hairy roots were harvested when the age of culture reached 1 month. The hairy roots were washed thoroughly with distilled water to get rid of excess culture medium. Then, the hairy roots were dried at 50°C overnight in the hot air oven and ground in mortar. The dried powder (37.4 g) was further macerated with methanol (100 ml, 3 times), filtered and evaporated. Crude methanol extract (8.5 g) was obtained and then re-dissolved in methanol (100 ml). The methanol fraction was partitioned with *n*-hexane (100 ml, 3 times) in the 1 L separatory funnel. The *n*-hexane extract was pooled and evaporated. The crude *n*-hexane (403 mg) was obtained and further purified by silica gel column chromatography. Silica gel column (size 3x18 cm) was prepared. Then, the crude *n*-hexane extract was loaded on the top column and eluted with sequential solvent of *n*-hexane: ethyl acetate; 9:1, 8:2, 7:3 and 5:5. The column was then eluted with polar solvent of ethyl acetate, ethyl acetate: methanol; 8:2 and 5:5. Finally, the column was washed with methanol. After following the elution fraction by TLC analysis, the 11 fractions were obtained. The scheme of isolation of secondary metabolites from the *M. speciosa* hairy roots was shown in Fig. 2.2.

For further purification, fractions of F6 (10 mg) and F10 (17.05 mg) were purified for re-crystallization. The fraction F6 was re-crystallized with the ratio of solvent of CHCl₃ and MeOH; 7:3. The white needle crystal was obtained, resulting in MSF1 (6.1 mg). The fraction F10 was washed with *n*-hexane. The white amorphous solid was obtained, resulting in MSF2 (10.11 mg). The resulting MSF1 and MSF2 were elucidated for chemical structures by ¹H- and ¹³C-NMR spectroscopy.

2.2.9 NMR spectroscopy

¹H and ¹³C-Nuclear magnetic resonance (¹H-125 MHz and ¹³C NMR-500 MHz), DEPT, COSY, HMQC, HMBC spectra were recorded on Fourier transformed NMR (FTNMR), Varian Unity Innova (USA), measured at the Scientific Equipment center, Prince of Songkla

university. CDCl_3 and CD_3OH were used as solvents and tetramethylsilane (TMS) was used as the reference.




Figure 2.2 Scheme of isolation of compounds from the *n*-hexane extract of *M. speciosa* hairy roots

2.2.10 Establishment of the growth and production curve of *M. speciosa* hairy roots

The growth curve and production curve of the hairy roots, cultured in WPM medium supplemented with 0.5 mg/l of NAA, were constructed. Time course study was performed. The samples (n=3) were taken every 5 days of culture, for a period of 40 days. The dry weight of each culture was determined and the data were plotted against the age of culture. For constructing the production curve, the ursolic acid content was determined from each sample (total of 24 samples) using the HPLC method. The production curve showed the relationship between the ursolic acid content and the age of culture. The ability of *M. speciosa* hairy roots for ursolic acid production was compared to the untransformed roots. In this study, the growth curve and production curve of the untransformed roots culture were also determined in the same procedure.

Determination of dry weight

The hairy roots cultures and the untransformed roots cultures were harvested and washed with distilled water. The excess water was removed by vacuum. Then, the cultures were dried at 50 °C in a hot air oven overnight and ground into powder. The cultures were weighed for the dry weight (g).

Determination of the ursolic acid content

The ursolic acid content was determined using HPLC method as described in Chen *et al.* (2003) with slight modification. The samples were prepared for *n*-hexane extract. After the samples were weighed for dried weight, the dried powder (100 mg) was extracted with *n*-hexane (50 ml, 3 x) under reflux for an hour and filtered. After that the filtrates were evaporated to dryness and were dissolved in acetonitrile. The volume was adjusted in a 5 ml-volumetric flask. The solution was further filtered through a 0.45 µm membrane filter prior to subjection to the HPLC system. Identification of the peak was performed from the peak retention time and UV absorption spectra of the corresponding peak, in comparison with the authentic ursolic acid (Sigma). The calibration curve of ursolic acid was constructed from the authentic compound. Standard ursolic acid 4.68 mg was dissolved in of acetonitrile and adjusted the volume to 10 ml. It was diluted with acetonitrile to concentration of 117, 58.5, 29.25, 14.625, 7.3125 µg/ml. The standard solutions were injected (10 µl) and run for calibration curves.

To calculate the ursolic acid content, the areas under the peaks were converted to concentrations by using the calibration curve. The HPLC analysis of each sample was performed in triplicate. The HPLC condition for separation of the ursolic acid was described below. The ursolic acid content was reported as mg per g dry weight.

HPLC conditions:

HPLC Column	: WATERS [®] USA, μ -BONDAPAK C18, 5 μ m particle size column size 3.9 x300 mm
Mobile phase	: acetonitrile : 0.1% H ₃ PO ₄ in H ₂ O (70:30), isocratic elution
Flow rate	: 1.0 ml/min
Detector	: UV 206 nm (photodiode array)
Injection volume	: 10 μ l

2.2.11 Plant regeneration of *M. speciosa*

The plant regenerations were observed in some clones of the hairy root culture, using the stems as explants. After excising the plantlets from the hairy root culture, they were placed on WPM solid medium, incubated at 25 °C under daylight for 16 h/day. Root initiation was observed after culture for 1 month. Therefore, in this study, we decided to study the micropropagation of *M. speciosa*, the regenerated plant that obtained from the hairy root culture. The regenerated plants were evaluated for the existing genes rooting loci for *rolA* and *rolB* of *A. rhizogenes* (2.2.5). After that, the mitragynine content of the regenerated plant was determined using the TLC-densitometric method (2.2.12), in comparison with the plantlet grown *in vitro* culture.

2.2.11.1 Induction of shoots and roots

The regenerated plantlets, initiated from the hairy root culture, were maintained in hormone-free WPM solid medium for 5 months. Then, the stems containing axillary buds were cut and placed on the WPM solid medium. For the experiment of multiple shoots induction, the

cytokinins including BA (1 mg/ml), thidiazuron (TDZ, 0.1 mg/l, 0.5 mg/l) were investigated by adding into the WPM solid medium. The cultures were kept at 25 °C under daylight for 16 h/day. About 5-7 samples of each medium were manipulated. The periods of shoot initiation and the shoot formations were recorded and calculated as the shoot number per explant. For root induction, shoots (containing 2-pairs of leaves) were excised and directly placed on the WPM solid medium without plant growth regulator. The roots were initiated after one week of culture.

2.2.11.2 Acclimatization

In order to transfer an *in vitro* plant to greenhouse, the plantlets (2.2.11.1), the regenerated plants were kept in the culture condition for 2 months. The plantlets were removed from the medium and the roots were gently washed in tap water. Then the plantlets were transferred to a bottle containing a mixture of sterilized soil and vermiculite (ratio 1:1 (w/w)), and further kept under *in vitro* condition for 1 month. To acclimatize the plantlets, they were transferred to a pot containing soil and placed under greenhouse conditions. Plant regenerations were irrigated every day. The survival rate of regenerated plants was calculated after 2 months of culture.

2.2.12 Establishment of thin layer chromatography-densitometric method

In this study, we tried to establish the rapid and simple method of thin-layer chromatography (TLC) densitometric method in order to simplify the method of mitragynine content determination. To prepare the crude alkaloid extract, the samples including hairy roots, untransformed root, and regenerated plants were harvested. They were dried at 50°C overnight in a hot air oven and ground. The dried powder (200 mg) was refluxed with methanol (50 ml, 3x) at 60 °C for 1 h and filtered. The methanolic filtrates were pooled and evaporated to dryness. The dried residue re-dissolved in the mixture of glacial acetic acid (2 ml) and distilled water (30 ml) and filtered. Acidic filtrate was washed with petroleum ether and basified with 25% (v/v) NH₃ to pH 9. Then, the filtrate was partitioned with chloroform (x ml, 3x), pooled and evaporated to dryness, obtaining the crude alkaloid extract. Finally, the crude alkaloid was re-dissolved in

chloroform (5 ml) and used for TLC-densitometric analysis. The scheme of the crude alkaloid extract preparation was shown in Fig. 2.3 (Keawpradub, 1990)

For calibration curve of mitragynine, the authentic mitragynine was used to prepare the concentration of 0.3-5.0 mg/ml in chloroform. Data points were obtained from independent 3-TLC plates and samples were performed in four-replicates at each concentration. The standard compound (5 μ l) was subjected to TLC plate, dried and run in the appropriate condition (see below). After removing from the tank, the TLC plate was dried and placed in the CAMAG TLC scanner. The R_f -value of mitragynine was 0.7. The peak area was integrated from that peak. The calibration curve was constructed by plotting the peak areas against the tested concentrations. The linear regression constant was determined.

The crude alkaloid extracts of samples including hairy roots (30 days old), untransformed root (30 days old), regenerated plants (5 months old) and plantlets (5 months old) were subjected to the TLC plate in the same procedure of preparation of the calibration curve. Integration of peak at R_f -value of 0.7 obtained the concentration of the sample by the conversion of peak area to concentration of calibration curve. The TLC densitometric analysis of each sample was performed in triplicate. The mitragynine content was reported as mg per g dry weight.

TLC densitometry condition:

TLC plate	: Aluminium sheet of silica gel 60 F-254
Mobile phase	: chloroform: methanol (9:1)
Plate size (width x height)	: 20.0 x 10.0 cm
Layer thickness	: 250 μ m
Application position Y	: 10.0 mm
Position of solvent	: 97.0 mm
Scan start position Y	: 3.0 mm
Scan end position X	: 95.0 mm
Scan start position X	: 15.0 mm
Distance between tracks X	: 20.0 mm
Lamp	: Deuterium
Wavelength	: ultraviolet light at 254 nm
Slit dimension	: 10 x 0.2 mm




Figure 2.3 Scheme of preparation of the crude alkaloid extract from *M. speciosa* hairy roots

CHAPTER 3

RESULTS

3.1 Establishment of transformed hairy root cultures of *M. speciosa*

3.1.1 Induction of hairy roots

Sterile seedlings were prepared from flat seeds of *M. speciosa*, *in vitro* culture. They were germinated within 1 week, and then maintained under culture condition for 2 months. Fig. 3.1 shows the sterile plantlets and parts of the plant including leaves, stems and adventitious roots, all used in this study.




Figure 3.1 *M. speciosa* grown in WPM solid medium supplemented with BA 1.0 mg/l A: 10 days old, B: plantlets after growing for 2 months.

According to Dhakulkar *et al.* (2005), the transformed hairy root culture of *M. speciosa* was induced with *Agrobacterium rhizogenes* ATCC 15834 with slight modification. Firstly, the hairy roots were induced by wounding at sites of the whole plantlet, including leaf veins and stems. Steps of hairy roots induction are summarized as shown in Fig. 3.2. From the results, the rootlets were initiated from injured sites and observed within 10 days. It can be noted that the frequency of hairy root induction was different in between leaf veins and stems.




Figure 3.2 Steps of the hairy roots induction from *M. speciosa* plantlet

- A. 2 months plantlets
- B. infection with *A. rhizogenes* ATCC 15834
- C. incubated the infected plantlet for 3 days
- D. culture in WPM plus cefotaxime 500 mg/l for 7 days
- E. culture in WPM plus cefotaxime 250 mg/l for 7 days
- F. culture in WPM plus cefotaxime 100 mg/l for 7 days

Therefore, a further experiment was done to determine an appropriate explant for the hairy roots induction. The leaves and stems of the isolated explants were then excised and infected with *A. rhizogenes* ATCC 15834. The results of hair roots induction were recorded on the 28th day after infection, as shown in Fig. 3.3.

Table 3.1 Induction of hairy roots from different explants

<i>Explants</i>	<i>Number of cultures used for genetic transformation</i>	<i>Number of cultures showing hairy root induction</i>	<i>% Hairy root induction</i>
Stem	189	128	67.72
Leaf vein	339	289	85.25




Figure 3.3 Characteristics of hairy roots, induced from leaf veins and stems

Fig. 3.3 illustrates the hairy roots characteristics, which were induced from the leaf vein and the stem. From Fig. 3.3, infection sites of the leaf vein gave a number of hairy roots ranging from 2-25 hairy roots per leaf vein, whereas the stem could only produce 3-8 hairy roots. Notably, the hairy roots obtained from the leaf vein appeared as thin and long and contained small pubescence. In contrast, the hairy roots obtained from the stem contained nodules (undeveloped hairy roots) and thick roots. To determine the potential of hairy roots formation in these explants, the percentages of hairy roots induction were calculated from the number of cultures used for *A. rhizogenes* infection and the number of cultures showing the hairy roots. As shown in Table 3.1, explants of leaf veins gave better number of hairy roots formation of 85%. Therefore, the explant of leaf vein was used for further genetic transformation with *A. rhizogenes* ATCC 15834 in order to get more clones for genes identification.

After the hairy roots grew in cefotaxime-free WPM medium, they were excised from the explant and placed on the surface of WPM solid medium when its length was about 3-4 cm. The hairy root culture was maintained under culture condition for 1 month (Fig. 3.4). After that, the hairy roots were transferred to WPM liquid medium and incubated at 25 °C, on a rotary shaker 80 rpm under dark condition.



M. speciosa hairy root culture of WPM solid medium

Figure 3.4 The characteristics of hairy root culture in WPM solid medium

For the untransformed root culture, the adventitious roots (Fig. 3.1) were excised and cultured in WPM liquid medium. Fig. 3.5 shows the characteristics of the transformed and untransformed roots after transferring to WPM liquid medium. Morphologically, transformed hairy roots induced by *A. rhizogenes* ATCC 15834 were thin and branching (Fig. 3.5A), whereas untransformed roots were thick, branching and shorter (Fig. 3.5B). From Fig. 3.5, young roots appeared white and gradually turned to slightly yellow-brown, when the cultures were aging. Under eye observation, both cultures were relatively grown under similar conditions.




Figure 3.5 Characteristics of transformed hairy roots (A) and untransformed root (B) cultures in WPM liquid medium at 22 days of culture.

3.1.2 Identification of *rolA* and *rolB* genes

The hairy roots (10 clones) were obtained from *A. rhizogenes* infections and used as materials for *rolA* and *rolB* identifications. The genomic DNAs from 10 clones of hairy roots (1 month old) were separately isolated and used as template in the PCR reactions. The genomic DNA of untransformed roots and plasmid DNA of *A. rhizogenes* were used as templates in the PCR reactions for the negative control and positive control, respectively. After performing the genomic DNA isolations, the concentrations of the DNA were determined by measuring the absorbance at 260 nm (A_{260}) in a spectrophotometer. The total DNA content was calculated according to the equation (see Appendix). To check the quality of isolated DNA, the absorbance at 280 nm (A_{280}) was also measured. The A_{280} indicated the amount of proteins, contaminated in the DNA fraction. Therefore, the ratio of A_{260}/A_{280} exhibited the purity of the DNA. Theoretically, the purity of total DNA should have an A_{260}/A_{280} ratio of 1.7-1.9.

Table 3.2 shows the concentrations of isolated DNAs and their purities. Data indicated that the total DNA concentrations were about 0.05-0.09 $\mu\text{g}/\mu\text{l}$. Calculation of A_{260}/A_{280} ratio of DNAs revealed that the isolated DNA contaminated with protein, meaning less quality of DNA was obtained (A_{260}/A_{280} about 0.2-0.6). Nevertheless, the total DNAs were further used as templates in the PCR reactions. After that the PCR reactions were performed, aliquots of reaction (5 μl) were analyzed for the DNA fragment in 1.2% agarose gel electrophoresis as shown in Fig. 3.6. For positive control (lane 2), DNA fragments, corresponding to *rolA* and *rolB*, were observed at the sizes of *ca.* 300 base pair and 780 base pair, respectively. None of DNA fragment appeared in the negative control (lane 3). Clones no. 4, 5, 6, 8, 9 and 10 (lanes 7, 8, 9, 11, 12, 13) were shown to carry the *A. rhizogenes* genes since the DNA fragments of 300 base pair and 780 base pair have been observed (Fig. 3.6). From this results, the genetic transformation rate of *A. rhizogenes* to *M. speciosa* leaf vein explant was accounted for 6/10 clones or 60%.

Table 3.2 Concentrations and purities of total DNA, extracted from the hairy root cultures

<i>Clone no.</i>	A_{260}	A_{280}	<i>Concentration of total DNA</i> <i>($\mu\text{g}/\mu\text{l}$)</i>	<i>Purity</i> <i>(A_{260}/A_{280})</i>
1	0.028	0.071	0.056	0.394
2	0.023	0.103	0.046	0.223
3	0.032	0.040	0.064	0.800
4	0.043	0.089	0.086	0.483
5	0.027	0.076	0.054	0.355
6	0.027	0.045	0.054	0.600
7	0.026	0.102	0.052	0.254
8	0.040	0.058	0.080	0.689
9	0.044	0.081	0.088	0.543
10	0.014	0.023	0.028	0.608




Figure 3.6 1.2% Agarose gel electrophoresis. PCR was performed with primers for the rooting locus genes *roIA* (A, 300 bp) and *roIB* (B, 780 bp). Lane 1 marker (10 kb DNA ladder), lane 2 *A. rhizogenes* ATCC 15834, lane 3 untransformed roots, lane 4-13 transformed hairy root (lines 1-10)

3.2 Medium manipulation

The hairy roots, in the WPM medium, grew slowly. The doubling time was about 2 months. In order to reduce the doubling time of the hairy root culture, the type of medium and the addition of plant growth regulator were manipulated. As described in 2.2.7 (Materials and methods), 9 types of media were investigated in the hairy roots (clone no.4). The appearances of the culture were observed after 1 month (Table 3.3). The untransformed roots in WPM medium and WPM plus NAA (0.5 mg/l) were used as control cultures. For time course determination, dry weights of hairy roots were determined from the hairy roots, which grew in WPM, WPM plus NAA, MS plus NAA and B5 plus NAA. The results were plotted and constructed for the growth curve (Fig. 3.7) and the relative growth plot (Fig. 3.8).

Table 3.3 shows the appearances of the hairy roots cultures of *M. speciosa* in different media. In the case of the untransformed roots culture in WPM plus NAA (0.5 mg/l), the roots were transformed to callus-like tissue. This culture was unlike the hairy roots, which cultured in media: WPM, MS or B5, supplemented with NAA. The hairy roots in $\frac{1}{2}$ WPM, MS and $\frac{1}{2}$ MS media grew badly. In contrast, the hairy roots could grow faster in the media, with the addition of NAA (0.5 mg/l). Interestingly, the appearances of the hairy roots in MS plus NAA and B5 plus NAA were slightly white-green. The time course of growths of the hairy roots were established in order to determine an appropriate medium for the hairy root culture.

Table 3.4 summarizes the dry weights of the hairy roots, which grew in WPM, WPM plus NAA, MS plus NAA and B5 plus NAA media, in comparison with the untransformed roots in WPM medium. The hairy roots were sampled every 5 days for over 40 days. The results were used to construct the growth curves (Fig. 3.7). The results revealed that the manipulated media could promote growth of the hairy roots as reduction of doubling time from 2 months to within 1 month. Growth curves were re-plotted in terms of the relative growth in order to eliminate the error of inoculation size. The relative growth plots indicated that additional 0.5 mg/l NAA in WPM, MS and B5 gave similar results. With the omission of NAA, hairy roots significantly grew slowly (Fig. 3.8). Therefore, the WPM plus 0.5 mg/l NAA was selected as an appropriate medium for biomass production, and the hairy roots were further evaluated for secondary metabolites production.

Table 3.3 The appearances of the hairy roots cultures in different media







<i>Type</i>	<i>Medium</i>	<i>Characteristics</i>	<i>After culture for 30 days</i>
Untransformed root	WPM	thick and short	
Untransformed root	WPM medium supplemented with plus 0.5 mg/l NAA	callus, brown color	
Hairy root	WPM	brown color and white color	
	MS	brown color	
	1/2 B5	dark brown and not grown	
	1/2 MS	brown color	

Table 3.3 (continued)





<i>Type</i>	<i>Medium</i>	<i>Characteristics</i>	<i>After culture 30 days</i>
	1/2WPM	dark brown and not grown	
	WPM medium plus 0.5 mg/l NAA	dark brown and white color	
	B5 plus 0.5 mg/l NAA	green color	
	MS plus 0.5 mg/l NAA	green color	

Table 3.4 The dry weights of the hairy roots in different media (clone no.4). (n=4)

<i>Day of culture</i>	Dry weight (mg \pm S.D.)				
	Untransformed roots biomass	Hairy roots biomass			
		WPM	WPM	WPM plus NAA*	B5 plus NAA*
5	2.2 \pm 0.2	26.6 \pm 2.2	27.9 \pm 4.6	8.8 \pm 0.3	7.1 \pm 0.4
10	2.2 \pm 0.4	36.0 \pm 0.2	25.3 \pm 1.6	11.4 \pm 0.6	9.1 \pm 0.6
15	8.9 \pm 2.5	40.8 \pm 0.9	38.0 \pm 6.2	15.8 \pm 0.1	12.4 \pm 0.4
20	13.5 \pm 0.8	43.1 \pm 0.6	59.4 \pm 1.6	18.3 \pm 0.1	14.0 \pm 0.8
25	14.8 \pm 1.9	46.4 \pm 2.3	66.0 \pm 11.5	20.3 \pm 0.5	17.3 \pm 0.3
30	25.0 \pm 4.0	53.7 \pm 2.3	72.6 \pm 9.1	24.9 \pm 0.3	20.7 \pm 0.5
35	33.6 \pm 2.0	60.7 \pm 3.3	72.1 \pm 10.6	26.5 \pm 0.8	22.6 \pm 0.7
40	31.9 \pm 2.6	54.3 \pm 0.3	72.4 \pm 4.1	27.8 \pm 1.5	26.6 \pm 1.2

*The culture medium was supplemented with NAA at the concentration of 0.5 mg/l.




Figure 3.7 Growth curves of hairy roots in selected media. Error bars represent standard deviations. (n=4)




Figure 3.8 Relative growth of hairy roots in selected media. Error bars represent standard deviations. (n=4)

3.3 Structure elucidation of MSF1 & MSF2 from *M. speciosa* hairy roots culture

For preliminary screening, the hairy roots culture of *M. speciosa*, grown in WPM medium supplemented with NAA (0.5 mg/l) was extracted and subjected to TLC plate. After developing, the TLC plate was then sprayed with Dragendroff's reagent to evaluate the potential of alkaloid formation. In the beginning, 2 spots of alkaloids were detected and they were positive to the Dragendroff's reagent. However, those spots were absent when the hairy roots were sub-cultured for 2-3 passages. This could probably have been due to the unstable compounds or because the alkaloids were from parts of the explant. Nevertheless, our attempts were to try to isolate the compound in order to evaluate the secondary metabolites. Therefore, *M. speciosa* hairy root culture was scaled up as described earlier (2.2.4). The compounds were isolated (2.2.8) and the structures of MSF1 and MSF2 were elucidated (2.2.9). MSF1 and MSF2 were obtained in percent yields of 0.016% and 0.027%, respectively.

3.3.1 MSF1

MSF1 was isolated from the *n*-hexane extract of transformed hairy root of *M. speciosa*, obtaining white needle crystal solids. In comparison with authentic β -sitosterol, the TLC analysis suggested that MSF1 was sterols. Analyses of ^1H - , ^{13}C -NMR spectra revealed that MSF1 was a mixture of β -sitosterol ($\text{C}_{29}\text{H}_{50}\text{O}$) and stigmasterol ($\text{C}_{29}\text{H}_{48}\text{O}$) (6.1 mg, 0.016 %w/w).

β -sitosterol, ^1H NMR (CDCl_3 , 500 MHz): δ H3.52 (1H, *m*, H-3), 5.35 (1H, *dd*, H-6), 0.68 (3H, *s*, Me-18), 0.98 (3H, *s*, Me-19), 0.91(3H, *d*, $J = 6.4$ Hz, Me-21), 0.83 (3H, *d*, $J = 6.8$ Hz, Me-26), 0.81 (3H, *d*, $J = 6.9$ Hz, Me-27), 0.85 (3H, *t*, $J = 7.8$ Hz, Me-29) (Fig. B-1).

Stigmasterol, ^1H NMR (CDCl_3 , 500 MHz): δ H3.53, (1H, *d*, H-3), 5.35 (1H, *d*, H-6), 0.66 (3H, *s*, Me-18), 0.98 (3H, *s*, Me-19), 0.90 (3H, *d*, $J = 6.4$ Hz, Me-21), 0.81 (3H, *d*, $J = 6.8$ Hz, Me-26), 0.79 (3H, *d*, $J = 6.9$ Hz, Me-27), 0.81 (3H, *t*, $J = 7.8$ Hz, Me-29) (Fig. B-1).

β -sitosterol, ^{13}C NMR (CDCl_3 , 125 MHz): δ C37.21 (C-1), 31.61 (C-2), 71.80 (C-3), 42.17 (C-4), 140.71 (C-5), 121.72 (C-6), 31.86 (C-7), 31.86 (C-8), 50.3 (C-9), 36.47 (C-10), 21.04 (C-11), 39.72 (C-12), 42.25 (C-13), 56.72 (C-14), 24.28 (C-15), 28.23 (C-16), 55.98 (C-17), 11.83 (C-18), 19.38 (C-19), 36.12 (C-20), 18.75 (C-21), 33.89 (C-22), 26.3 (C-23), 45.77 (C-24), 29.07 (C-25), 19.81 (C-26), 18.89 (C-27), 23.01 (C-28), 11.96 (C-29) (Fig. B-2).

Stigmasterol, ^{13}C NMR (CDCl_3 , 125 MHz): δ C37.21 (C-1), 31.61 (C-2), 71.80 (C-3), 42.25 (C-4), 140.71 (C-5), 121.72 (C-6), 31.86 (C-7), 31.86 (C-8), 50.07 (C-9), 36.47 (C-10), 21.04 (C-11), 39.63 (C-12), 42.25 (C-13), 56.82 (C-14), 24.34 (C-15), 28.92 (C-16), 55.92 (C-17), 11.96 (C-18), 19.38 (C-19), 40.51 (C-20), 21.04 (C-21), 138.32 (C-22), 129.21 (C-23), 51.21 (C-24), 31.86 (C-25), 21.09 (C-26), 18.95 (C-27), 25.4 (C-28), 12.02 (C-29) (Fig. B-2).

Analysis of chemical shifts, integration and spin coupling patterns from ^1H NMR data were in agreement with NMR data of the published report (Subhadhirasakul and Pechongs, 2005). From ^{13}C -spectrum, δ at 138.32 (C-22) and at 129.21 (C-23) were observed. Both signals corresponded to the double bond in the structure of stigmasterol. Therefore, signals of H-22 and H-23 in ^1H -NMR were integrated and calculated for the molarities. The results suggested that MSF1 was a mixture of β -sitosterol and stigmasterol, present in the ratio of 1:1. The structures of both compounds are shown in Fig. 3.9. All spectra are shown in the Appendix.




Figure 3.9 Structures of β -sitosterol and stigmasterol

3.3.2 MSF2

MSF2 was isolated from the *n*-hexane extract of the transformed hairy root of *M. speciosa*, obtaining an amorphous solid. The ^1H spectrum of MSF2 was suggested that MSF2 exhibited the structure as typical as the structure of triterpene compound. Later, MSF2 was sent to perform for the ^{13}C -NMR, DEPT, 2D-NMR spectra. Analyses of all spectra indicated that MSF2 was ursolic acid ($\text{C}_{30}\text{H}_{48}\text{O}_3$), co-existing with oleanolic acid ($\text{C}_{30}\text{H}_{48}\text{O}_3$) (10.11 mg, 0.027 %w/w).

Ursolic acid, ^1H NMR (500 MHz, CDCl_3 : CD_3OD): 3.20 (t, 1H, $J = 7.5$ Hz, H-3), 5.24 (m, $J = 3.5$ Hz 1H, H-12), 2.20 (d, $J = 11.0$ Hz, 1H, H-18), 2.02-1.15 (m, H-22), 0.98 (s, 3H, Me-23), 0.78 (s, 3H, Me-24), 0.93 (s, 3H, Me-25), 0.82 (s, 3H, Me-26), 1.10 (s, 3H, Me-27), 0.86 (d, $J = 6.5$ Hz, 3H, Me-29), 0.95 (d, $J = 6.5$ Hz, 3H, Me-30) (Fig. B-3)

Ursolic acid, ^{13}C -NMR (125 MHz, CDCl_3 : CD_3OD): 38.4 (C-1), 26.5 (C-2), 78.5 (C-3), 39.2 (C-4), 55.0 (C-5), 18.0 (C-6), 32.8 (C-7), 39.2 (C-8), 47.3 (C-9), 36.7 (C-10), 23.9 (C-11), 125.2 (C-12), 137.9 (C-13), 41.8 (C-14), 23.9 (C-15), 23.0 (C-16), 47.6 (C-17), 52.6 (C-18), 38.8 (C-19), 38.7 (C-20), 30.4 (C-21), 36.6 (C-22), 27.7 (C-23), 15.3 (C-24), 15.1 (C-25), 16.6 (C-26), 23.2 (C-27), 180.5 (C-28), 16.7 (C-29), 20.9 (C-30) (Fig. B-4)

DEPT 90 experiment CH (CDCl_3 : CD_3OD): 78.5 (C-3), 55.0 (C-5), 47.3 (C-9), 125.2 (C-12), 52.6 (C-18), 38.8 (C-19), 38.7 (C-20) (Fig. B-5)

DEPT 135 experiment CH_2 (CDCl_3 : CD_3OD): 38.4 (C-1), 26.5 (C-2), 18.0 (C-6), 32.8 (C-7), 23.9 (C-11), 23.9 (C-15), 23.0 (C-16), 30.4 (C-21), 36.6 (C-22); CH_3 : 27.7 (C-23), 15.3 (C-24), 15.1 (C-25), 16.6 (C-26), 23.2 (C-27), 16.7 (C-29), 20.9 (C-30) (Fig. B-6)

From ^{13}C -NMR spectrum, MSF2 composed of 30 carbons in its structure. DEPT90 and DEPT135 spectra suggested the types of carbons. The presence of δ at 78.5 (C 3; COH), 125.2 (C 12), 137.9 (C 13), 180.5 (C 28; C=C) suggested that MSF2 was ursolic acid. 2D-NMR: COSY (Fig. B-7), HMQC (Fig. B-8), HMBC (Fig. 3.12 and Fig B-9) indicated the

correlation of ^{13}C - ^{13}C , ^1H - ^{13}C and long-length coupling ^1H - ^{13}C , respectively. Analysis of NMR spectra of MSF2 was in agreement with the previous report (Güvenalp *et al.*, 2006). HMBC spectrum suggested the correlation of carbon and hydrogen atoms as shown in Table 3.5. The presence of signal at δ 5.28 MHz in ^1H -NMR spectra indicated that MSF2 contained not only ursolic acid but also oleanolic acid. This evidence was in line with the presence of signals at δ 122 MHz and 144 MHz in ^{13}C -NMR spectrum (Fig. B-4). Integration of H-12 signal in ^1H -NMR indicated that MSF2 was a mixture of ursolic acid and oleanolic acid in the ratio of 5:1. All spectra are shown in Appendix.

Table 3.5 HMBC: ^1H and ^{13}C NMR spectroscopic data for ursolic acid

<i>H/C</i>	δ ^1H (J, Hz)	δ ^{13}C (ppm)	HMBC correlations
3	3.2 (<i>t</i> , 7.5 Hz)	78.5	C-1, C-2, C-23, C-24
12	5.24(<i>m</i> , 3.5 Hz)	125.2	C-9, C-14, C-18
18	2.2 (<i>d</i> , 11 Hz)	52.6	C-13, C-16, C-28
25	0.93 (<i>s</i>)	15.1	C-1, C-5, C-9
26	0.82 (<i>s</i>)	16.6	C-7, C-9, C-14
27	1.10 (<i>s</i>)	23.2	C-8, C-13, C-14, C-15
29	0.86 (<i>d</i> , 6.5 Hz)	16.7	C-18, C-20
30	0.95 (<i>d</i> , 6.5 Hz)	20.9	C-19, C-21




Figure 3.10 Structures of ursolic acid and oleanolic acid

3.4 Ursolic acid production in *M. speciosa* hairy roots

Since the hairy roots of *M. speciosa* were maintained in WPM medium and supplemented to NAA (0.5 mg/l), they could produce ursolic acid. Previously it had been reported that ursolic acid exhibits varieties of pharmacological activities (Liu, 1995). In this study, ursolic acid production was, therefore, determined. The time course of ursolic acid production was investigated. Calibration curve of an authentic ursolic acid was constructed. The elution time of ursolic acid was 15.5 min under HPLC condition (2.2.10) (Fig. 3.12). The linearity was ranged from 7.31-117 $\mu\text{g/ml}$ with the linear regression (R^2) of 0.9999 (Fig. 3.11).




Figure 3.11 Calibration curves of authentic acid ursolic acid.

To construct the growth curve and production curve, the samples ($n=3$) were taken every 5 days for over 40 days of culture. The ursolic acid content of sample was determined after weighing the dry weight. The untransformed root culture was used as a control culture. The hairy roots and the untransformed hairy roots were extracted and prepared for acetonitrile fractions (2.2.10). The sample (10 μl) was subjected to the HPLC system. The chromatograms of authentic ursolic acid, the extracts from the untransformed roots and the hairy roots are shown in

Fig. 3.12. The identities of ursolic acid were confirmed by UV absorption spectra (200-300 nm) using the diode array detector.





Figure 3.12 HPLC-chromatograms of (A) authentic ursolic acid (RT=15.5 min); (B) the untransformed roots extract; (C) the hairy roots extract and (D) UV absorption spectra of peak at 15.5 min using the HPLC-UV diode array detector.

Table 3.6 Biomass and ursolic acid contents in untransformed roots and hairy roots (n=4)

Day	Untransformed roots		Transformed hairy roots	
	biomass (g dry wt./flask)	ursolic acid content (mg/g dry wt. \pm S.D.)	biomass (g dry wt./flask)	ursolic acid content (mg/g dry wt. \pm S.D.)
5	0.003 \pm 0.001	1.578 \pm 0.012	0.025 \pm 0.006	1.520 \pm 0.004
10	0.004 \pm 0.004	1.742 \pm 0.008	0.028 \pm 0.006	1.902 \pm 0.023
15	0.007 \pm 0.004	1.891 \pm 0.007	0.042 \pm 0.009	2.490 \pm 0.023
20	0.013 \pm 0.001	1.691 \pm 0.007	0.058 \pm 0.018	1.649 \pm 0.012
25	0.016 \pm 0.003	2.090 \pm 0.009	0.062 \pm 0.012	2.004 \pm 0.040
30	0.027 \pm 0.005	2.410 \pm 0.008	0.071 \pm 0.010	3.468 \pm 0.034
35	0.035 \pm 0.004	1.740 \pm 0.023	0.085 \pm 0.027	1.981 \pm 0.027
40	0.035 \pm 0.007	2.010 \pm 0.004	0.086 \pm 0.028	2.398 \pm 0.010

The area under peak at 15.5 min was integrated and converted to ursolic acid concentration using the calibration curve. The results are summarized in Table 3.6. Growth curves and production curves of the hairy root culture and the untransformed root culture are shown in Fig. 3.13 and Fig. 3.14, respectively. The results indicated that both types of culture were able to produce ursolic acid. However, at 30 days of culture, the hairy roots accumulated the ursolic acid with the yield of 3.468 ± 0.034 mg/g dry wt, whereas the untransformed roots could produce 2.410 ± 0.008 mg/g dry wt. The results show that the ursolic acid was produced and accumulated at the late linear phase (Fig. 3.13) for both types of culture.

Fig. 3.12 indicates that the cell cycle of the *M. speciosa* hairy roots was within 30 days of culture. It can be seen that the cycle composed on 3 phases: days 1-10 for lag phase, days 10-20 for exponential phase, days 20-30 for linear phase and days 30-40 for stationary phase. The production curve of the hairy roots show that ursolic acid was usually present in the cells and maintained at a particular level of *ca.* 2 mg/g dry wt. However, the hairy roots were able to accumulate maximum yield at the late linear phase. Thus, the ursolic acid was utilized by the cells and acted as a secondary metabolite in this *in vitro* culture.




Figure 3.13 Growth curve and production curve of *M. speciosa* hairy root culture. Error bars represent standard deviations.




Figure 3.14 Growth curve and production curve of the *M. speciosa* untransformed root culture. Error bars represent standard deviations.

3.5 Micropropagation of *M. speciosa*

During the establishment of *M. speciosa* hairy root culture, shoot regeneration was observed from cultures, which were maintained in WPM liquid medium. This evidence was observed when the hairy roots contained part of the stem (cut from the original explant). Nevertheless, the shoot regeneration disappeared when the culture was further sub-cultured. Shoot development from the hairy roots is shown in Fig. 3.15. After 2 months, the plantlets were removed and placed in WPM solid medium. The roots were then initiated after culture for 10 days in the same medium. The *in vitro* plantlets were used as materials for micropropagation.




Figure 3.15 Shoot regeneration from *M. speciosa* hairy root culture, maintained in WPM liquid medium

Micropropagation began with the *M. speciosa* plantlet, which was grown in WPM solid medium. The cultures were maintained at 25 °C under daylight for 16 h/day (Fig. 3.16). Axillary buds were excised from the stems and used in the experiments for shoot multiplication.




Figure 3.16 Plant regeneration from the *M. speciosa* hairy root culture (4 months old)

To increase the number of shoots in the *M. speciosa* plantlet, the axillary buds were excised and manipulated in WPM solid medium supplemented with two kinds of cytokinin. In this study, cytokinins, BA, TDZ at 0.1 mg/l and TDZ at 0.5 mg/l, were used for the shoot multiplication. The results were observed during 1 month of culture. Fig. 3.7 illustrates the formation of shoots in manipulated media. After 2 weeks, the shoots were initiated and appeared in red. This characteristic was the same as the young shoots of *M. speciosa* in nature. The shoot numbers were counted and calculated as the shoot number per explant. Table 3.7 shows the shoot number per explant of cultures, grown in different cytokinins. The result indicated that TDZ was shown to be an appropriate cytokinin for shoot multiplication with about 6 shoots per explant. Concentration of 0.1 mg/l and 0.5 mg/l of TDZ effected insignificantly the shoot multiplication. Therefore, *M. speciosa* shoots multiplied from the axillary buds using the WPM medium supplemented with TDZ at the concentration of 0.1 mg/l. The cultures were placed at 25°C under daylight for 16 h/day for 2 months before root initiation.




Figure 3.17 Shoot multiplication of the axillary buds, cultured in different cytokinins

A: in WPM supplemented with 1 mg/l BA; A-1, 1st day of culture; A-2, 12 days old, A-3: 26 days old

B: in WPM supplemented with 0.1 mg/l TDZ; B-1, 1st day of culture; B-2, 12 days old, B-3: 26 days old

C: in WPM supplemented with 0.5 mg/l TDZ; C-1, 1st day of culture; C-2, 12 days old, C-3: 26 days old

Table 3.7 Shoot number of plantlet regeneration in WPM solid medium

Hormone	Number of axillary buds	Shoot number/explant (Mean \pm S.D.)
without hormone	5	not determined
1 mg/l BA	6	2.83 \pm 1.47
0.1 mg/l TDZ	6	6.33 \pm 1.63
0.5 mg/l TDZ	7	6.29 \pm 1.25



Figure 3.18 Root initiations of the regenerated plantlet of *M. speciosa* in WPM solid medium (4 months old)

Root initiation of the regenerated plantlet of *M. speciosa* was simply performed. The shoots (containing 2-4 leaves) were cut from explants and placed on WPM solid medium without any plant growth regulator. Roots were initiated after 1 week of culture. Then, the complete plantlets were grown for 2 months, before transferring to greenhouse condition.

M. speciosa regenerated plantlets were successfully acclimatized by sequential changing of the supporting materials (Fig. 3.19). The acclimatization protocol composed of two steps of culture. Firstly, the plantlets were transferred to culture in sterilized mixture of vermiculite and soil (1:1) and grown under controlled conditions for 30 days (Fig 3.19B). Finally, they were further grown in the pot containing soil for 1 month (Fig 3.19C). The plantlets were watered every day. After 2 months, the percentage of survival was evaluated at 60%. The characteristics of regenerated plants appeared the same as the original plants (Fig 3.19D).



Figure 3.19 Acclimatization protocol for *M. speciosa* regenerated plant

A: culture in WPM solid medium for 2 months

B: culture in vermiculite: soil (1:1) for 30 days

C: culture in soil for 30 days

D: culture in soil after 6 months

3.6 Mitragynine production of the regenerated *M. speciosa* plantlets

Mitragynine contents of *in vitro* plantlets and regenerated plantlets were determined. The *in vitro* plantlets were obtained from seed germination and grew under controlled conditions. The simple and rapid method of mitragynine quantification was established using the TLC-densitometric method. The TLC plate was developed and mitragynine was migrated at R_f value of 0.7 (Fig. 3.21). The detector was set at 254 nm at the maximum absorbance of mitragynine. First of all, the calibration curve of the authentic mitragynine was established. The linearity of the calibration curves was in the range of 0.3-5 mg/ml with R^2 of 0.9984 (Fig. 3.20). The peak identities were performed by scanning the UV absorbance (200-600 nm) as shown in Fig. 3.20.




Figure 3.20 Calibration curve of authentic mitragynine

The extracts from the whole plant of *in vitro* and regenerated plantlets (5 months olds) were prepared for the alkaloid extracts (2.2.12). The samples (5 μ l) were subjected to the TLC plate and developed. The TLC was further developed for the chromatograms at the wavelength of 254 nm. The TLC-densitochromatograms are shown in Fig. 3.21. The peaks at R_f value of 0.7 were integrated. The area under peaks was converted to concentrations using a calibration curve. Table 3.8 summarizes the mitragynine content from 5-month old plantlets, grown under culture conditions.




Figure 3.21 TLC densito-chromatograms (at $\lambda = 254$ nm) and UV absorption (200-600 nm) of A: authentic mitragynine, B: *in vitro* plantlet, C: regenerated plant, D: leaves of regenerated plant.

The 5 months of *in vitro* and regenerated plantlets of *M. speciosa* were shown to produce mitragynine in different contents. As shown in Table 3.8, mitragynine content in the

regenerated plantlet was 14.25 ± 0.25 mg/g dry wt., about 3.2 times higher than that *in vitro* plantlet, which was 4.45 ± 0.09 mg/g dry wt. Leaves of the regenerated plantlet were isolated and the mitragynine content was determined. The results show that mitragynine was produced and accumulated in leaves.

Table 3.8 Mitragynine content determination from 5 months old plants

<i>Sample</i>	<i>mitragynine content^d</i> <i>(mg/g dry wt. \pm S.D.)</i>
<i>in vitro</i> plantlet ^a	4.45 \pm 0.09
regenerated plant ^b	14.25 \pm 0.25
leaves from regenerated plant ^c	12.5 \pm 0.25

^afrom plantlet that germinated from seeds and grown in WPM medium supplemented with BA (1 mg/l)

^bfrom regenerated plantlets (3.6)

^ccut leaves from regenerated plantlets (3.6)

^dSamples were determined in triplicate.

CHAPTER 4

DISCUSSION

Mitragyna speciosa (Roxb.) Korth. (Rubiaceae) is a source of mitragynine, a monoterpenoid indole alkaloid (MIA), which acts as opium-like effect. The main objective of this investigation was to establish hairy roots culture of *M. speciosa* by infection with *Agrobacterium rhizogenes* ATCC 15834. Since the susceptibility of plant cells to *A. rhizogenes* was dependent upon age and differentiation status of plant tissue (Sevón and Oksman-Caldentey, 2002). Therefore, the hairy root culture, in this study, was induced from small plantlets, which were germinated under controlled conditions. The meristematic tissues of stems and leaves were used as explants due to ease for differentiation. The transformed hairy roots were identified for the *rolA* and *rolB* as genetic markers. Six of ten clones were positive to have genes from *A. rhizogenes* ATCC 15834 (transformation rate of 60%), which means low frequency of transformation. Unlike the transformation of *Gmelina arborea*, the herbaceous plants, transformation rate was about 100% (Dhakulkar *et al.*, 2005). Type of woody plant of *M. speciosa* affected the transformation rate which may be discussed. However, the chance of successful transformation may be obtained from infection with different strains of *A. rhizogenes* (Sevón and Oksman-Caldentey, 2002).

Theoretically, the hairy root culture should grow relatively fast and can grow in plant growth regulator free medium (Guillon *et al.*, 2006). But the *M. speciosa* hairy roots obtained from this study could grow slowly. For this study, a low concentration of NAA (0.5 mg/l) is required for promoting their growth. This evidence is unusual for normal hairy root culture (Sevón and Oksman-Caldentey, 2002). In the case of other Rubiaceae plants, *Ophiorrhiza pumila* hairy root culture in B5 medium (containing 2% (w/v) sucrose) could grow fast and produce the camptothecin, a MIA-anticancer agent (Saito *et al.*, 2007). It can be suggested that the hormonal balance in *M. speciosa* hairy roots was not appropriate for root proliferation (Guillon *et al.*, 2006).




Figure 4.1 Proposed biosynthetic pathway of mitragynine

The hairy roots of *M. speciosa* could accumulate compounds, which are triterpenoids; ursolic acid and oleanolic acid (with the ratio of 5:1) and phytosterols, β -sitosterol and stigmasterol (with the ratio of 1:1). Nevertheless, many studies reported that the alkaloids such as mitraphylline, rhynchophylline have been found in the root bark of *M. speciosa* (Houghton and Shellard, 1974; Shellard *et al.* 1978b). But none of the alkaloids were found in this culture. Intermediates of alkaloids were probably unstable and degraded during culture (Fig 4.1). However, the hairy root culture accumulated the ursolic acid, which is pharmacologically important for antibacterial, anti-inflammatory, antiviral activities, etc. (Liu, 1995). It produced in the late linear phase a yield of 3.468 ± 0.034 mg/g dry wt. It can be noted that *M. speciosa* hairy roots could accumulate high yield of ursolic acid when compared to the *Uncaria tomentosa* cell suspension culture (1.680 ± 0.039 mg/g dry wt.) (Feria-Romero *et al.*, 2005). This evidence concludes that the differentiated cells such as hairy roots preferred to produce higher amount of secondary metabolites.

For the biosynthetic point of view (Fig. 4.1), *M. speciosa* hairy roots could not produce any alkaloid, especially mitragynine. It may be caused by a lack of precursors and enzymes, which is involved in the terpenoid indole alkaloid, since primary metabolites need special cell compartments for storage and degradation (Luckner, 1984). The presence of ursolic acid, coexisting with oleanolic acid, indicated that at least isoprene units have been produced in this culture as shown in Fig. 4.1. These isoprene units are precursors for both triterpenoids and phytosterols (Luckner, 1984), which means their biosynthesis was active in this culture. Therefore, the *M. speciosa* hairy root culture is not only useful for ursolic acid production but also have a potential to be a model plant culture for triterpenoid biosynthesis.

Unexpectedly, plantlet regeneration of *M. speciosa* was obtained. Micropropagation of this plant was performed successfully transferring the plants to nature. From this study, TDZ was shown to be an efficient cytokinin for shoot multiplication. TDZ acts as a substitute for both auxin and cytokinin requirements to induce organogenesis and somatic embryogenesis in many species (He, 2007). From this study, the regenerated plantlets produced and accumulated the considerable amount of mitragynine, when compared to the *in vitro* plantlets (from seed germination). It was postulated that this may be caused by the effect of TDZ. Nevertheless, it is still unknown.

CHAPTER 5

CONCLUSION

- 5.1 *Mitragyna speciosa* hairy root culture was successfully induced from leaf veins of explants using *Agrobacterium rhizogenes* ATCC 15834, at a transformation rate of 85.25%. Hairy roots were morphologically characterized by phenotype and genotype. Hairy roots were fast growing, thin, and branching with pubescence hairs, distinguished from untransformed roots. Experiments of PCR identified the presence of the *rolA* and *rolB* genes with an incorporation rate of 60%.
- 5.2 Optimal medium that promoted the hairy roots growth was investigated in the presence of low concentration of NAA. The results concluded that WPM, B5 and MS media containing 0.5 mg/l NAA promoted the hairy roots growth. The optimal conditions for the hairy root culture were in WPM medium (containing 2% (w/v) sucrose), supplemented with NAA (0.5 mg/l) and the cultures were incubated at 25°C under darkness.
- 5.3 Secondary metabolites accumulated in the hairy roots were evaluated. No alkaloid was found. Triterpenoids; ursolic acid and oleanolic acid, and phytosterols; β -sitosterol and stigmasterol, were isolated from the *n*-hexane fraction of the hairy roots with yields of 0.027% (w/w) and 0.016% (w/w), respectively
- 5.4 The ursolic acid content was determined using HPLC analysis. The growth curve and production curve of *M. speciosa* hairy roots were established. The production curve suggested that ursolic acid was produced and accumulated at the late linear phase with a yield of 3.468 ± 0.034 mg/g dry wt. about 1.4 times higher than that from untransformed roots.
- 5.5 For micropropagation, TDZ at 0.1 mg/l and 0.5 mg/l in WPM solid medium could induce the shoot multiplication for 6.33 ± 1.63 and 6.29 ± 1.25 shoots per explant, respectively. Micropropagation of *M. speciosa* regenerated plants successfully with a 60% survival rate.

5.6 A simple and rapid technique of TLC-densitometry was established for mitragynine content quantification. Authentic mitragynine was used to construct the calibration curve. Samples of regenerated plantlets, leaves of regenerated plantlets, *in vitro* plantlets were determined for mitragynine production. The results showed that regenerated plantlets accumulated the mitragynine yielding an amount of 14.25 ± 0.25 (mg/g \pm S.D.), that was 3 times higher than that of the *in vitro* plantlets.

REFERENCES

- เต็ม สมิตินันท์. 2544. ชื่อพรรณไม้แห่งประเทศไทย. กรุงเทพฯ: สวนพฤกษศาสตร์ป่าไม้ สำนักวิชาการป่าไม้ กรมป่าไม้.
- Aji, A.M., Effraim, K.D. and Onyeyili, A.P. 2001. Antistress activity of *Mitragyna africanus* (Wild) stembark extract. *Sciences* 1, 105-107.
- Benjamin, B.D., Roja, G. and Heble, M.R. 1994. Alkaloid synthesis by root cultures of *Rauwolfia serpentine* transformed by *Agrobacterium rhizogenes*. *Phytochemistry* 35, 381-383.
- Chen, J.H., Xia, Z.H. and Tan, R.X. 2003. High-performance liquid chromatographic analysis of bioactive triterpenes in *Perilla frutescens*. *J. Pharm. Biomed. Anal.* 32: 1175-1179.
- Costa, V.B., Coube, C.S., Marinho, B.G., Matheus, M.E., Leitao, S.G. and Fernandes P.D. 2003. Anti-inflammatory and analgesic activity of *Bouchea fluminensis*. *Fitoterapia* 74, 364–371.
- Dewick, P.M. 2002. Medicinal natural product: a biosynthetic approach. 2nd, *John Wiley and sons*, UK, 167-225.
- Dhakulkar, S., Ganapathi, T.R., Bhargava, S. and Bapat, V.A. 2005. Induction of hairy roots in *Gmelina arborea* Roxb. and production of verbascoside in hairy roots. *Plant Sci.* 169, 812-818.
- Feria-Romero, I., Lazo, E., Ponce Noyola, T., Cerda Garcia Rojas, C.M. and Ramos Valdivia, A.C. 2005. Induced accumulation of oleanolic acid and ursolic acid in cell suspension cultures of *Uncaria tomentosa*. *Biotechnol. Lett.* 27, 839-843.
- Flores-Sanchez, I.J., Ortega-Lopez, J., Montes-Horcasitas, M.C. and Ramos-Validivia, A.C. 2002. Biosynthesis of sterols and triterpenes in cell suspension cultures of *Uncaria tomentosa*. *Plant Cell Physiol.* 43: 1502-1509.
- Gamborg, O.L., Miller, R.A., and Ojima, K. 1968. Nutrient requirement of suspensions cultures of soybean root cells. *Exp. Cell. Res.* 50, 151-158.
- Guillon, S., Tremouillaux Guiller, J., Pati, P.K., Rideau, M. and Gantet, P. 2006. Hairy root research: recent scenario and exciting prospects. *Plant Biol.* 9, 341-346.
- Guvenalp, Z., Kilic, N., Kazaz, C., Kaya, Y. and Demirezer, O. 2006. Chemical constituents of *Galium tortumense*. *Turk. J. Chem.* 30, 515-523.

- Hamill, J.D., Robins, R.J. and Rhodes, M.J.C. 1989. Alkaloid production by transformed root cultures of *Cinchona ledgeriana*. *Planta Med.* 55, 354-357.
- He, S.S., Liu, C.Z. and Saxena, P.K. 2007. Plant regeneration of an endangered medicinal plant *Hydrastis canadensis* L. *Scientia Horticulturae* 113, 82-86.
- Hinou, J. and Harvala, C. 1988. Polyphenolic compounds from the leaves of *Mitragyna speciosa*. *Fitoterapia* 59, 156.
- Houghton, P.J. and Said, I.M. 1986. 3-dehydromitragynine: An alkaloid from *Mitragyna speciosa*. *Phytochemistry* 25(12), 2910-2912.
- Houghton, P.J. and Shellard, E.J. 1974. The alkaloidal pattern in *Mitragyna rotundifolia* from Burma. *Planta med.* 26, 104-112.
- Kamatou, G.P.P., Van Vuuren, S.F., Van Heerden, F.R., Seaman, T. and Viljoen, A.M. 2007. Antibacterial and antimycobacterial activities of South African *Salvia* species and isolated compounds from *S. chamelaeagnea*. *S. Afr. J. Bot.* 73, 552-557.
- Keawpradub, N. 1990. Alkaloids from the fresh leaves of *Mitragyna speciosa*. Master's Thesis, Graduate School, Chulalongkorn University, Bangkok.
- Kohda, H., Namera, A., Koyama, A., Yamasaki, K. and Tani, T. 1996. Indole alkaloid production in callus cultures of *Uncaria rhynchophylla*. *Chem. Pharm. Bull.* 44, 352-357.
- Kumarnsit, E., Vongvatcharanon, U., Keawpradub, N. and Intasaro, P. 2007. Fos-like immunoreactivity in rat dorsal raphe nuclei induced by alkaloid extract of *Mitragyna speciosa*. *Neurosci. Lett.* 416, 128-132.
- Liu, J. 1995. Pharmacology of oleanolic acid and ursolic acid. *J. Ethnopharmacol.* 49, 57-68.
- Lloyd, G. and McCown, B. 1980. Commercially-feasible micropropagation of mountain laurel, *Kalmia latifolia*, by use of shoot-tip culture. *Int. Plant Prop. Soc. Proc.* 30, 421-427.
- Luckner, M. 1984. Secondary metabolism in microorganisms, plants and animals. *Springer Verlag Berlin Heidelberg*, New york, 1-563.
- Matsumoto, K., Mizowaki, M., Thongpraditchote, S., Murakami, Y., Takayama, H., Sakai, S-L., Aimi, N. and Watanabe, H. 1996a. Central antinociceptive effects of mitragynine in mice: contribution of descending noradrenergic and serotonergic pathways. *Eur. J. pharmacol.* 371, 75-81.

- Matsumoto, K., Mizowaki, M., Thongpraditchote, S., Takayama, H., Sakai, S-L., Aimi, N. and Watanabe, H. 1996b. Antinociceptive action of mitragynine in mice : evidence for the involvement of supraspinal opioid receptors. *Life Sci.* 59, 1149-1155.
- Mizukami, H., Nordlov, H., Lee, S.L. and Scott, A.I. 1979. Purification and properties of strictosidine synthetase (an enzyme condensing tryptamine and secologanin) from *Catharanthus roseus* cultured cells. *Biochemistry* 28, 3760.
- Murashige, T., and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15, 473-497.
- Oksman-Caldentey, K.M., and Hiltunen, R. 1996. Transgenic crops for improved pharmaceutical products. *Field Crops Res.* 45, 57-69.
- Paranhos, J.T., Fragoso, V., Henriques, A.T., Ferreira, A.T. and Fett-Neto, A.T. 2005. Regeneration of *Psychotria umbellata* and production of the analgesic indole alkaloid umbellatine. *Tree Physiol.* 25, 251–255.
- Phillipson, J.D., Tantivatana, P., Tarpo, E. and Shellard, E.J. 1973. Hirsuteine and Mitrajavine from *Mitragyna hirsute*. *Phytochemistry* 12, 1507-2048.
- Said, I.M., Chun, N.C. and Houghton, P.J. 1991. Ursolic acid from *Mitragyna speciosa*. *Planta Med.* 57, 398.
- Saito, K., Sudo, H., Yamazaki, M., Koseki-Nakamura, M., Kitajima, M., Takayama, H. and Aimi, N. 2007. Feasible production of camptothecin by hairy root culture of *Ophiorrhiza pumila*. *Plant Cell Rep.* 20, 267-271.
- Sevón, N., Hiltunen, R. and Oksman-Caldentey, K.M. 1998. Somaclonal variation in *Agrobacterium* transformed roots and protoplast-derived hairy root clones of *Hyoscyamus muticus*. *Planta Med.* 64, 37-41.
- Sevón, N. and Oksman-Caldentey, K.M. 2002. *Agrobacterium rhizogenes*-Mediated transformation: Root cultures as a source of alkaloids. *Planta Med.* 68, 859-868.
- Shellard, E.J., Beckett, A.H., Tantivatana, P., Phillipson, J.D. and Lee, C.M. 1966. Alkaloids from *Mitragyna javanica* Hoord and Valetton and *Mitragyna hirsuta* Havil. *J. Pharm. Pharmac.* 18, 553-555.
- Shellard, E.J., Houghton, P.J. and Resha, M. 1978a. The *Mitragyna* species of Asia Part XXXI. The alkaloids of *Mitragyna speciosa* Korth. from Thailand. *Planta Med.* 34, 26-36.

- Shellard, E.J., Houghton, P.J. and Resha, M. 1978b. The *Mitragyna* species of Asia Part XXXII. The distribution of alkaloids in young plants of *Mitragyna speciosa* Korth. Grown from seed obtained from Thailand. *Planta Med.* 34, 253-263.
- Sim, S.J., Chang, H.N., Liu, J.R. and Jung, K.H. 1994. Production and secretion of indole alkaloid in hairy root cultures of *Catharanthus roseus*: effects of *in situ* adsorption, fungal elicitation and permeabilization. *J. Ferment. Bioeng.* 78, 229-234.
- Somova, L.I., Shode, F.O., Ramnanan, P. and Nadar, A. 2003. Antihypertensive, antiatherosclerotic and antioxidant activity of triterpenoids isolated from *Olea europaea*, subspecies *africana* leaves. *J. Ethnopharmacol.* 84, 299-305.
- Stella, A. and Braga, M.R. 2002. Callus and cell suspension cultures of *Rudgea jasminoides*, a tropical woody Rubiaceae. *Plant cell, Tiss. Org. Cult.* 68, 271-276.
- Sudha, C.G., Obul Reddy, B., Ravishankar, G.A. and Seeni, S. 2003. Production of ajmalicine and ajmaline in hairy root cultures of *Rauvolfia micrantha* Hook f., a rare and endemic medicinal plant. *Biotechnol. Lett.* 25, 631-636.
- Subhadhirasakul, S. and Pechpongs, P. 2005. Terpenoid and two steroids from *Mammea siamensis*. *Songklanakarin J. Sci. Technol.* 27, 555-561.
- Sukrong, S., Zhu, S., Ruangrunsi, N., Phadungcharoen, T., Palanuvej, C. and Komatsu, K. 2007. Molecular analysis of the genus *Mitragyna* existing in Thailand based on rDNA ITS sequences and its application to identify a narcotic species: *Mitragyna speciosa*. *Biol. Pharm. Bull.* 30, 1284.
- Takayama, H., Ishikawa, H., Kurihara, M., Kitajima, M., Aimi, N., Ponglux, D., Koyama, F., Matsumoto, K., Moriyama, T., Yamamoto, L.T., Watanabe, K., Murayama, T. and Horie, S. 2002. Studies on the synthesis and opioid agonistic activities of *Mitragynine*-related indole alkaloids: discovery of opioid agonists structurally different from other opioid ligands. *J. Med. Chem.* 45, 1949-1956.
- Takayama, H., Kurihara, M., Kitajima, M., Said, I.M., and Aimi, N. 1998. Alkaloids from the leaves of Malaysian *Mitragyna speciosa*. *Tetrahedron* 54(29), 8433-8440.
- Takayama, H. 2004. Chemistry and Pharmacology of analgesic indole alkaloids from the Rubiaceae plant, *Mitragyna speciosa*. *Chem. Pharm. Bull.* 52, 916-928.

- Thongpradichote, S., Matsumoto, K., Tohda, M., Takayama, H., Aimi, N., Sakai, S. and Watanabe, H. 1998. Identification of opioid receptor subtypes in antinociceptive actions of supraspinally-administered mitragynine in mice. *Life Sci.* 62, 1371-1378.
- Tsuchiya, S., Miyashita, S., Yamamoto, M., Horie, S., Sakai, S.I., Aimi, N., Takayama, H. and Watanabe, K. 2002. Effect of mitragynine, derived from Thai folk medicine, on gastric acid secretion through opioid receptor in anesthetized rats. *Eur. J. Pharmacol.* 443, 185–188.
- Watanabe, K., Yano, S., Horie, S. and Yamamoto, L.T. 1997. Inhibitory effect of mitragynine, an alkaloid with analgesic effect from Thai medicinal plant *Mitragyna speciosa*, on electrically stimulated contraction of isolated guinea-pig ileum through the opioid receptor. *Life Sci.* 60, 933-942.

APPENDIX I

Table A-1 Inorganic salt and vitamin composition of Murashige & skoog medium (MS)
(DUCHEFA BIOCHEMIE B.V.) (Murashige and Skoog, 1962)

<i>Constituent</i>	<i>Concentration (mg/liter)</i>
Macronutrients:	
CaCl ₂	332.02
KH ₂ PO ₄	170.00
KNO ₃	1,900.00
MgSO ₄	180.54
NH ₄ NO ₃	1650.00
Micronutrients:	
CoCl ₂ .6H ₂ O	0.025
CuSO ₄ .5H ₂ O	0.025
FeNaEDTA	36.70
H ₂ BO ₃	6.20
KI	0.83
MnSO ₄ .H ₂ O	16.90
NaMoO ₄ .2H ₂ O	0.25
ZnSO ₄ .7H ₂ O	8.60
Sucrose (g)	30.00
Vitamins:	
Glycine	2.00
myo-Inositol	100.00
Nicotinic acid	0.50
Pyridoxine hydrochloride	0.50
Thiamine hydrochloride	0.10
pH	5.8

To prepare the MS medium using the mixed powder 4.4 g/l

Table A-2 Inorganic salt and vitamin composition of McCown Woody Plant (DUCHEFA BIOCHEMIE B.V.) (Lloyd and McCown, 1980)

<i>Constituent</i>	<i>Concentration (mg/liter)</i>
Macronutrients:	
CaCl ₂	72.50
Ca(NO ₃) ₂ ·4H ₂ O	471.26
KH ₂ PO ₄	170.00
K ₂ SO ₄	990.00
MgSO ₄	180.54
NH ₄ NO ₃	400.00
Micronutrients:	
CuSO ₄ ·5H ₂ O	0.25
FeNaEDTA	36.70
H ₃ BO ₃	6.20
MnSO ₄ ·H ₂ O	22.30
NaMoO ₄ ·2H ₂ O	0.25
ZnSO ₄ ·7H ₂ O	8.60
Sucrose (g)	20.00
Vitamins:	
Glycine	2.00
myo-Inositol	100.00
Nicotinic acid	0.50
Pyridoxine hydrochloride	0.50
Thiamine hydrochloride	1.00
pH	5.7

To prepare the WPM medium using the mixed powder 2.5 g/l

Table A-3 Inorganic salt and vitamin composition of Gamborg's B5 medium (DUCHEFA BIOCHEMIE B.V) (Gamborg *et al.*, 1968)

<i>Constituent</i>	<i>Concentration (mg/liter)</i>
Macronutrients:	
CaCl ₂	113.23
KNO ₃	2500.00
MgSO ₄	121.56
NaH ₄ PO ₄	130.44
(NH ₄) ₂ SO ₄	134.00
Micronutrients:	
CoCl ₂ .6H ₂ O	0.025
CuSO ₄ .5H ₂ O	0.025
Disodium EDTA	36.70
H ₃ BO ₃	3.00
KI	0.75
MnSO ₄ .H ₂ O	10.00
NaMoO ₄ .2H ₂ O	0.25
ZnSO ₄ .7H ₂ O	2.00
Sucrose (g)	20.00
Vitamins:	
myo-Inositol	100.00
Nicotinic acid	1.00
Pyridoxine hydrochloride	1.00
Thiamine hydrochloride	10.00
pH	5.5

To prepare the B5 medium using the mixed powder 3.2 g/l

APPENDIX II

1. Plasmid isolation from *Agrobacterium rhizogenes* ATCC 15834

Plasmid of *A. rhizogenes* (positive control) was isolated from the overnight culture using GFX Micro Plasmid Prep Kit (Qiagen). Overnight culture (1.5 ml) was transferred to a microcentrifuge tube and centrifuged at 13,000 rpm for 30 s to pellet the cells. The supernatant matters were removed. The pellet was resuspended in solution I (150 μ l) in a vigorous vortex. Solution II (150 μ l) was added to the pellet and mixed for 10-15 times. Then solution III (300 μ l) was added and mixed by inverting the tube until a flocculent precipitate appeared. After that the mixture was centrifuged at 13,000 rpm for 5 min to pellet cell debris. The supernatant matters were transferred to a GFX column and incubated for 1 min at room temperature. The column was centrifuged at 13,000 rpm for 30 s. The flow-through was discarded. The column was added with solution III (300 μ l) and centrifuged at 13,000 rpm for 30 s. Washing buffer (400 μ l) was added and then centrifuged at 13,000 rpm for 60 s. Then the column was centrifuged at 13,000 rpm for additional 1 min. Finally the GFX column was transferred to a new tube. Tris-EDTA buffer (100 μ l) was added and incubated at room temperature for 1 min, centrifuged at 13,000 rpm for 1 min, and the plasmid DNA was obtained. The purified DNA was kept at -20°C until it was required for use.

2. Genomic DNA isolation from transformed hairy root cultures

Total DNA was isolated from hairy roots as well as from untransformed roots using DNeasy Plant Mini Kit (Qiagen). The plant tissues were ground into fine powder in the presence of liquid N₂. The powder was re-suspended in extraction buffer AP1 (400 μ l) plus 4 μ l of RNase A stock solution (100 mg/ml). The mixture was set to a vigorous vortex. The suspension was incubated for 10 min at 65°C, mixed 2-3 times during incubation by inverting tube. Buffer AP2 (130 μ l) was added to the lysate and incubated on ice for 5 min. The lysate was then loaded into the QIAshredder mini spin column in a 2 ml collection tube and centrifuged at 14,000 rpm for 2 minutes. The flow-through was transferred to a new tube. Buffer AP3/E (1.5

volumes) was added into the cleared lysate and mixed by pipette. The mixture (650 μ l) was moved to a DNeasy mini spin column sitting in a 2 ml collection tube and centrifuged at 8,000 rpm for 1 min. The flow-through was discarded. Buffer AW (500 μ l) was added on DNeasy mini spin column and centrifuged for at 8,000 rpm 1 min. Finally the DNeasy mini spin column was placed in a 1.5 ml microcentrifuge tube and adding buffer AE (100 μ l), incubated for 5 min at room temperature. Finally, the plant DNA was eluted from the column after it was centrifuged for 1 min at 8,000 rpm. The plant DNA was stored at -20°C until it was required for use.

The concentration and purity of DNA were determined by measuring the absorbance at 260 nm (A_{260}) and 280 nm (A_{280}) in a spectrophotometer. An absorbance of 1.0 at 260 nm corresponds to 50 μ g of DNA per ml ($A_{260}=1=50 \mu\text{g/ml}$). The purity was determined by calculating the ratio of absorbance at 260 nm to the absorbance at 280 nm. Pure DNA has an A_{260}/A_{280} ratio of 1.7-1.9. Concentration of total DNA was calculated from the equation below.

$$\text{Concentration of total DNA } (\mu\text{g}/\mu\text{l}) = (A_{260}) (\text{dilution factor}) (50)$$




Figure B-1 ^1H NMR spectrum of mixture of MSF1 in CDCl_3




Figure B-2 ^{13}C NMR spectrum of mixture of MSF1 in CDCl_3





Figure B-2 (Continued)




Figure B-2 (Continued)





Figure B-2 (Continued)




Figure B-2 (Continued)




Figure B-2 (Continued)




Figure B-2 (Continued)





Figure B-2 (Continued)




Figure B-3 ^1H NMR spectrum of MSF2 in $\text{CDCl}_3:\text{CD}_3\text{OD}$




Figure B-4 ^{13}C NMR spectrum of MSF2 in $\text{CDCl}_3:\text{CD}_3\text{OD}$





Figure B-4 (Continued)





Figure B-4 (Continued)




Figure B-4 (Continued)





Figure B-4 (Continued)





Figure B-5 DEPT 90 of MSF2 in $\text{CDCl}_3:\text{CD}_3\text{OD}$




Figure B-6 DEPT 135 of MSF2 in $\text{CDCl}_3:\text{CD}_3\text{OD}$




Figure B-7 COSY of MSF2 in CDCl₃:CD₃OD




Figure B-8 HMQC spectrum of MSF2 in CDCl₃:CD₃OD





Figure B-9 HMBC spectrum of MSF2 in CDCl₃:CD₃OD

VITAE

Name Miss Siriwan Phongprueksapattana

Student ID 4752012

Education Attainment

Degree	Name of Institution	Year of Graduation
Bachelor of Science (Chemistry)	Prince of Songkla University	2003

List of Publication and Proceeding

Siriwan Phongprueksapattana, Waraporn Putalun, Wanchai De-Eknamkul, Jurathip Wungsintaweekul. 2006. *Agrobacterium rhizogenes* transformed hairy root cultures from *Mitragyna speciosa* Korth. Proceeding of 3rd Life Sciences Postgraduate Conference 2006. Universiti Sain Malaysia, Penang, Malaysia, May 24-27, 2006.