CONTENTS

· ·	Page
บทกัดย่อ	(3)
Abstract	
Acknowledgment	(4)
Contents	(6)
List of abbreviations	(7)
List of tables	(8)
List of figures	(9)
Chapter	(10)
1. Introduction	1
2. Literature reviews	3
3. Materials and mehods	61
4. Results	
5. Discussion	69
Bibliography	90
Appendix	100
Vitae	123
-	124

LIST OF ABBREVIATIONS

°C = degree Celsius

IC₅₀ = inhibitory concentration at 50% of maximum response

G = gram

kg = kilogram

M = molar

mg = miligram

ml = milliliter

mM = millimolar

 $\mu M = micromolar$

μg = microgram

ng = nanogram

% = percent

S.E. = standard error

h = hour

ACh = acetylcholine

5-HT = serotonin

KCl = potassium chloride

CaCl₂ = calcium chloride

NaCl = sodium chloride

NaHCO₃ = sodium bicarbonate

MgSO₄ = magnesium sulfate

KH₂PO₄ = potassium dihydrogen phosphate

X = mean

LIST OF TABLES

Γable	Page
1	Substances that may be neurotransmitters or
	neuromodulators in the enteric nervous system
2	Yield and texture of methanolic extract from each plant
3	Effect of methanolic extracts of P. longum, P. sarmentosum
	and Q. infectoria on caecal amoebiasis of mice
4	IC ₅₀ values (concentration producing 50% of maximum
	inhibition of agonist-induced contractions of rat or
	guinea-pig isolated ileums, $n = 5$) of the three plant
	methanolic extracts, loperamide, verapamil, the
	muscarinic receptor antagonist, atropine; the
	serotoninergig receptor antagonist, cyproheptadine and
	the histaminergic receptor antagonist, chlorpheniramine80
5	Comparison of the potencies (on µg/ml concentration
	basis) of the methanolic extracts of P. sarmentosum, P.
	longum and Q. infectoria on the inhibition of the agonist-
	induced contractions of rat or guinea-pig ileums with
	those of: (a) loperamide (Lop) and (b) verapamil (Ver)
	and (c) atropine (Atrop), cyproheptadine (Cypro) and
	chlorpheniramine (CPM)
6	The inhibitory effects of P. sarmentosum, P. longum and Q.
	infectoria methanolic extracts and loperamide on the phasic
	and tonic contraction-induced by KCl of rat ileums
7	The IC ₅₀ values of the three plant extracts and loperamide
	on the inhibition of phasic and tonic contractions of KCl-
	induced contraction of rat ileums and their potency ratios

LIST OF FIGURES

Figure		Page
1	Entamoeba histolytica	4
2	Structural formula of metronidazole	
3	Photograph of medicinal plants	
4	The general organization of the layers of the	
	gastrointestinal tract	24
5	Neural control of the gut wall	
6	Diagram of the wall of the small intestine	
7	Sources of intracellular Ca ²⁺	
8	Positive and negative-feedback loops regulating Ca2+	
	entry	39
9	Putative mechanisms for activation of Ca2+ by	
	agonists	40
10	Calcium-mobilizing intracellular messengers	
11	Mechanisms of smooth muscle contraction and signal	
	transduction mechanisms	45
12	Mechanisms of agonists-induced contraction and	
	relaxation	46
13	The balance between the relaxant or contractile state	
	of smooth muscle	52
14	Signaling cascade initiated by contractile agonist in	
	intestinal longitudinal muscle cells	59
15	Dual signaling cascade initiated by interaction of VIP	
	with two receptor subtypes	60
16	Set up of isolated ileum preparation in organ bath for	
	recording of its contraction	66

LIST OF FIGURES (Continued)

igure	Page
17	Representative recordings of the acetylcholine-
	induced contractions of isolated rat ileums and the
	inhibitions by various concentrations of the
	methanolic extracts of P. sarmentosum, P. longum and
	Q. infectoria73
18	Representative recordings of the serotonin-induced
	contractions of isolated rat ileums and the inhibitons
	by various concentrations of the methanolic extracts
	of P. sarmentosum, P. longum and Q. infectoria74
19	Representative recordings of the histamine-induced
	contractions of isolated guinea-pig ileums and the
	inhibitions by various concentrations of the
	methanolic extracts of P. sarmentosum, P. longum and
	Q. infectoria75
20	Inhibition of acetylcholine-induced contractions of
	isolated rat ileums by atropine, loperamide, verapamil
	and the methanolic extracts of P. sarmentosum,
	P. longum and Q. infectoria
21	Inhibition of serotonin-induced contractions of
	isolated rat ileums by cyproheptadine, loperamide,
	verapamil and the methanolic extracts of
	P. sarmentosum, P. longum and Q. infectoria77
22	Inhibition of histamine-induced contractions of
	isolated guinea-pig ileums by cyproheptadine,
	loperamide, verapamil and the methanolic extracts of
	P. sarmentosum, P. longum and Q. infectoria78

LIST OF FIGURES (Continued)

Figure	•	Page
23	Representative recordings of potassium chloride-	
	induced contractions (each consists of phasic, initial	
	rapid contraction, followed by a sustained tonic	
	contraction) of rat ileums and the inhibition by	
	various concentrations of the methanolic extracts of	
	P. sarmentosum, P. longum and Q. infectoria	82
24	Inhibition of potassium chloride-induced phasic	
	contraction (A) and tonic contraction (B) of rat	
	ileums by loperamide (0.3-10 µg/ml), P. sarmentosum	
	(0.01-1 mg/ml), P. longum (0.01-1 mg/ml) and	
	Q. infectoria (0.1-10 mg/ml)	84
25	Representative recordings of the effects of the methanolic	
	extract of (b) P. longum (0.1-1 mg/ml), (c) P. sarmentosum	
	(0.1-1 mg/ml), (d) Q. infectoria (1-10 mg/ml), (e)	
	loperamide (0.1-1 μ g/ml) and (f) verapamil (0.01-0.1 μ M)	
	on the contractions of guinea pig ileum-induced by	
	calcium chloride (added cumulatively), in comparison with	
	those of time control (a)	86
26	Cumulative concentration-effects of calcium chloride	
	induced-contractions of isolated guinea-pig ileums	
	and the antagonisms by P. sarmentosum methanolic	
	extract and P. longum methanolic extract	87
27	Cumulative concentration-effects of calcium chloride	
	induced-contractions of isolated guinea-pig ileum and	
	the antagonisms by Q. infectoria methanolic extract and	
	loperamide	88

LIST OF FIGURES (Continued)

Figure		Page
28	Cumulative concentration-effects of calcium chloride-	
	induced contraction of isolated guinea-pig ileums and	
	the antagonisms by verapamil	80