

Contents

	Page
บทคัดย่อ	(3)
Abstract	(5)
Acknowledgement	(7)
Contents	(9)
List of Tables	(15)
List of Figures	(17)
Abbreviations and Symbols	(20)
Chapter	
1. Introduction	1
1.1 Introduction	1
1.2 Objectives	6
2. Literature review	7
2.1 Original and geographic distribution	7
2.2 Description of <i>Elaeis guineensis</i> Jacq.	8
2.3 Botanical information	10
2.4 Genetic resources	12
2.5 Palm oil	14
2.6 Economic and production data	15
2.7 Imbibition	21

Contents (Continued)

	Page
2.8 Flow cytometry	22
2.8.1 Principles of flow cytometry	23
2.8.2 The use of reference plant	24
2.8.3 Cell cycle analysis	24
2.8.4 Genomic DNA alterations	25
2.9 Fundamental of plant cell culture	26
2.9.1 Plasticity and totipotency	26
2.9.2 The culture environment	27
2.9.3 Plant cell culture media	28
2.9.4 Carbon sources	28
2.9.5 Gelling agents	28
2.9.6 Plant growth regulators	29
2.10 Review of oil palm <i>in vitro</i> culture works	31
2.10.1 Protoplast isolation and culture	32
2.10.2. Early experiments	33
2.10.3 Importance and use of oil palm protoplast	33
2.10.4 Background of genetic transformation	34
2.10.5 Protoplast culture	35

Contents (Continued)

	Page
3. Materials and methods	38
3.1 Seedling and callus induction	38
3.1.1 Plant materials	38
3.1.2 Seedling and callus initiation	38
3.1.3 Culture media and conditions	39
3.2 Flow cytometric analysis	39
3.2.1 Plant materials	39
3.2.2 Reference plants testing	40
3.2.3 Genomic DNA alteration	40
3.2.4 Cell cycle activity analysis	40
3.2.5 Data analysis	41
3.3 Protoplast isolation and culture	41
3.3.1 Plant materials	41
3.3.2 Protoplast isolation	42
3.3.3 Protoplast purification	43
3.3.4 Yield and viability measurements	43
3.3.5 Protoplast culture methods and protoplast sources investigation	44
3.3.6 Gelling agents and carbon sources investigation	44
3.3.7 Cell division induction	44

Contents (Continued)

	Page
3.3.8 Cell colony formation	45
3.3.9 Data analysis	45
4. Results	46
4.1 Seedling and callus induction	46
4.1.1 Seedling induction	46
4.1.2 Callus formation	49
4.2 Flow cytometric analysis	51
4.2.1 Reference plants testing	51
4.2.2 Genomic DNA alteration	54
4.2.3 Cell cycle activity analysis	57
4.3 Protoplast isolation and culture	60
4.3.1 Protoplast isolation	60
4.3.2 Effect of protoplast culture methods and protoplast sources on cell division	65
4.3.3 Effect of gelling agents and carbon sources on cell division of protoplasts	66
4.3.4 Cell division induction	67
4.3.5 Cell colony formation	69
5. Discussion	72

Contents (Continued)

	Page
5.1 Effect of physical conditions and growth regulators on embryo culture and cell cycle activity	72
5.1.1 Effect of imbibition on seedling and callus induction	72
5.1.2 Effect of imbibition on cell cycle activity	74
5.1.3 Effect of growth regulators on callus induction	76
5.2 Flow cytometry in oil palm	78
5.3 Protoplast isolation and culture	80
5.3.1 Protoplast isolation	81
5.3.2 Protoplast culture	83
5.3.3 Protoplast division and cell colony formation	85
6. Conclusion	87
References	90
Appendix	103
1 Composition of Murashige and Skoog (MS) medium (1962).	104
2 Composition of Eeuwens (Y ₃) medium (1976).	106
3 Composition of CPW salts solution (Power <i>et al.</i> , 1984) in 1000 ml.	108
4 Protocol for determination of protoplast viability (FDA stain) (Widholm, 1972).	109
5 Protocol for preparation of oil palm embryo protoplasts.	110

Contents (Continued)

	Page
6 Protocol for flow cytometric analysis (Pfosser <i>et al.</i> , 1998).	111
7 Lipid grains and storage proteins in oil palm embryos after imbibed for 24-72 h.	113
8 Embryogenic calli of oil palm cultured on MS medium supplemented with 4.5 μ M 2,4-D and 0.1% activated charcoal.	114
9 The results of flow cytometric analysis of oil palm using reference plant as an external standard.	115
Publications	119
Vitae	120

List of Tables

Table	Page
1 Output values of various oilseed crops.	16
2 World production of palm oil by major countries (1000t).	17
3 The 20 highest producing countries of palm oil for the year 2003 (by FAO).	18
4 World mature areas of oil palm (1000 ha).	19
5 The 20 highest exporter countries of palm kernel oil for the year 2002 (by FAO).	20
6 Genome size (2C nuclear DNA content in pg) of <i>E. guineensis</i> Jacq. estimated by comparing with various reference plants. The means are based on 10 replicates per experiment.	52
7 Genome size (2C nuclear DNA content in pg) of oil palm embryos, one-year-old shoot meristem of seedlings and of calli at the 3 rd <i>in vitro</i> subculture. The means are obtained from experiments with 5 replicates.	55
8 Yield (10^5 protoplasts gram ⁻¹ FW) of oil palm protoplasts isolated from seedlings and calli with different enzyme mixture after 3h of incubation.	60
9 Yield of oil palm protoplasts isolated from friable callus cultured in liquid media with enzyme mixture of 1% Cellulase+0.5% Driselase+1% Macerozyme.	61
10 Yield (10^6 protoplasts gram ⁻¹ FW) and viability (%) of oil palm protoplasts isolated from 48h-imbibed embryos with different enzyme mixtures.	63

List of Tables (Continued)

Table	Page
11 Percentage of cell division of embryo-derived oil palm protoplasts cultured on MS medium supplemented with 13.6 μM 2,4-D, 4.4 μM BA and 9.3 μM kinetin using bead technique containing different gelling agents and carbon sources.	67
12 Percentage of cell division of embryo-derived oil palm protoplasts cultured in agarose bead medium containing various types and concentrations of growth regulators (μM).	69
13 Plating efficiency of embryo-derived oil palm protoplasts cultured in different mixtures of growth regulators (μM), after 3 weeks of culture.	71

List of Figures

Figure	Page
1 Oil palm plantation at Horticultural Research Center, Suratthani Province.	9
2 Mature oil palm fruit bunch, <i>Tenera</i> cultivar.	10
3 Young oil palm seed cultivars <i>Pisifera</i> 116 (A); <i>Dura</i> 64 (B) and <i>Tenera</i> 23 (C), 3 months after anthesis.	11
4 Mature oil palm seed cultivar <i>Tenera</i> , 6 months after anthesis.	12
5 Pattern of cell cycle activity from flow cytometric histogram.	25
6 Germination oil palm seedlings from imbibed (a and c) and non-imbibed (b and d) embryos cultured in liquid (a and b) or on solid (c and d) Y3 media without plant growth regulators after culture for 28 days.	47
7 Germination oil palm seedlings from embryos imbibed for 24 48 and 72 h and non-imbibed (control) prior to cultured in liquid Y3 media without plant growth regulators after culture for 56 days.	48
8 Effect of imbibition times on fresh weight, shoot and root length of oil palm seedlings derived from imbibed and non-imbibed embryos cultured in Y3 liquid medium after culture for 56 days. Error bars represent \pm SE.	49

List of Figures (Continued)

Figure	Page
9 Callus formation from embryos imbibed for 0, 24, 48 and 72 hours prior to culture in liquid Y3 media supplemented with the following growth regulators, 4.1 μM picloram and 9.0 μM 2,4-D (media 1); 8.3 μM picloram and 4.5 μM 2,4-D (media 2) and 13.6 μM 2,4-D (media 3) after culture for 56 days.	50
10 Fresh weight of calli derived from embryos imbibed for 0, 24, 48 or 72 hours after culture in Y3 liquid media supplemented with 4.1 μM picloram and 9.0 μM 2, 4-D (media 1); 8.3 μM picloram and 4.5 μM 2, 4-D (media 2) and 13.6 μM 2, 4-D (media 3) after culture for 56 days. Error bars represent $\pm\text{SE}$.	51
11 Histogram of fluorescence intensity of nuclei isolated from oil palm using <i>Glycine max</i> cv. Polanka (A), <i>Lycopersicon esculentum</i> cv. Stupicke (B) and <i>Zea mays</i> line CE-777 (C) as internal reference plants.	54
12 Comparison of fluorescence intensity in nuclei of oil palm among embryos (A), long-term <i>in vitro</i> cultured tissues of shoot meristem (B) and calli (C) using <i>Glycine max</i> cv. Polanka as an internal reference.	57
13 Histogram of propidium iodide fluorescence intensity of oil palm nuclei isolated from embryos imbibed for 24, 48 and 72 hour and non-imbibed embryos. The fluorescence histograms were resolved into 2C (G0/G1) and 4C (G2/M) cell cycle compartments with a peak-reflect algorithm using WinMDI version 2.8. X-axis represents DNA content; Y-axis represents event of cell counts.	58

List of Figures (Continued)

Figure	Page
14 Histogram of propidium iodide fluorescence intensity of oil palm nuclei isolated from embryos imbibed for 48 hour (a), calli (b), shoots (c) and roots (d). The fluorescence histograms were resolved into 2C (G0/G1) and 4C (G2/M) cell cycle compartments with a peak-reflect algorithm using WinMDI version 2.8. X-axis represents DNA content; Y-axis represents event of cell counts.	59
15 Freshly isolated protoplast from basal leaf (A) and callus (B).	61
16 (A) Freshly isolated protoplasts from embryos of oil palm and (B) the same protoplasts when stained with FDA.	64
17 Percentage of cell division of oil palm protoplasts from different sources of explants cultured on MS medium using agarose as gelling agent supplemented with 8.3 μM picloram, 4.5 μM 2,4-D, 8.8 μM BA and 4.6 μM kinetin after culture for 7 days. Error bars represent $\pm\text{SE}$.	65
18 Dividing oil palm protoplasts cultured on MS medium containing 8.3 μM picloram, 4.5 μM 2,4-D, 9.3 μM kinetin and 4.4 μM BA, after 2 days of culture. Bar=10 μm .	68
19 Cell colony formation of oil palm protoplasts cultured in Agarose bead MS medium containing 4.5 μM 2,4-D, 4.5 μM dicamba, 9.3 μM kinetin, 4.4 μM BA and 3.8 μM ABA, after 3 weeks of culture. Bar=20 μm .	70

Abbreviations and Symbols

2,4-D	=	2,4-dichlorophenoxyacetic acid
ABA	=	abscisic acid
AC	=	Activated charcoal
ANOVA	=	Analysis of variance
BA	=	N ⁶ - benzyladenine
bp	=	base pair (s)
Cel	=	Cellulase Onozuka R-10
CFW	=	Calcofluor white
cm	=	centimetre
CPW	=	Cell protoplast wash
cv.	=	cultivar
Dicamba	=	3,6-dichloro-o-anisic acid
DNA	=	Deoxyribonucleic acid
Dri	=	Driselase
FAO	=	Food and Agricultural Organization
FDA	=	Fluorescein diacetate
g (unit of acceleration)	=	gravity
g (unit of weight)	=	gram
gram ⁻¹ FW	=	per gram fresh weight
h	=	hour

Abbreviations and Symbols (Continued)

Kinetin	=	6-furfurylaminopurine
M	=	molar
Mac	=	Macerozyme R-10
MES	=	2-N-morpholinoethanesulphonic acid
mg	=	milligram
mg/l	=	milligram per litre
min	=	minute
ml	=	millilitre
mm	=	millimetre
mM	=	millimolar
MS	=	Murashige and Skoog (MS), 1962
NAA	=	1-naphthaleneacetic acid
nm	=	nanometre
<i>p</i>	=	probability level
PE	=	Plating efficiency
PEG	=	Polyethylene glycol
PI	=	propidium iodide
Picloram	=	4-amino-3,5,6-trichloropicolinic acid
pg	=	picogram
pp	=	protoplast

Abbreviations and Symbols (Continued)

PVP	=	polyvinylpyrrolidone
rpm	=	revolution per minute
SD	=	standard deviation
SE	=	standard error
sh	=	shell (endocarp)
v/v	=	volume by volume
w/v	=	weight by volume
Y ₃	=	Eeuwens (Y ₃), 1976
°C	=	degree of Celsius
μm	=	micrometre
μl	=	microlitre
μmol m ⁻² s ⁻¹	=	micromole per metre square per second
%	=	percentage