

## CONTENTS

	<b>Page</b>
บทคัดย่อ	iii
ABSTRACT	v
ACKNOWLEDGEMENTS	vii
CONTENTS	ix
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xviii
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Literature review	2
1.2.1 Zinnia	2
1.2.1.1 Scientific classification	2
1.2.1.2 Origin/History of zinnia	3
1.2.1.3 The advantages of <i>Zinnia</i>	4
1.2.1.4 The cultivars of <i>Zinnia</i>	7
1.2.2 Propagation of zinnia	8
1.2.2.1 Traditional propagation	8
1.2.2.2 <i>In vitro</i> culture	11
1.2.2.3 Genetic variation	16
1.2.3 Cytogenetic and molecular biological scientific methods	17
1.2.3.1. Flow cytometry	17

## **CONTENTS** (continued)

	<b>Page</b>
1.2.3.2. RAPD-PCR	25
1.3 Objectives	29
<b>CHAPTER 2 MATERIALS AND METHODS</b>	<b>30</b>
Plant materials	30
Methods	35
2.1 Cultivar classification by flow cytometry	35
2.2 Cultivar classification by RAPD	40
2.3 Cultivar classification by morphology of guard cell and plant height	42
2.4 Chromosome numbers	43
2.5 <i>In vitro</i> culture	44
<b>CHAPTER 3 RESULTS</b>	<b>48</b>
3.1 Cultivar classification by flow cytometry	48
3.2 Cultivar classification by RAPD	58
3.3 Cultivar classification by morphology of guard cell and plant height	62
3.4 Chromosome numbers	65
3.5 <i>In vitro</i> culture	67
<b>CHAPTER 4 DISCUSSION</b>	<b>90</b>
4.1 Cultivar classification by flow cytometry	90
4.2 Cultivar classification by RAPD	99

## **CONTENTS** (continued)

	<b>Page</b>
4.3 Cultivar classification by morphology of guard cell and plant height	100
4.4 Chromosome numbers	101
4.5 <i>In vitro</i> culture	102
CHAPTER 5 CONCLUSION	106
REFERENCES	109
APENDICES	124
Technique Specifications of flow cytometry	125
VITAE	129

## LIST OF TABLES

	<b>Page</b>
Table 1.1 Use of <i>Zinnia</i> spp.	5
Table 1.2 List of fluorochrome used for flow cytometry	20
Table 2.1 Comparison of morphology of cultivars	31
Table 2.2 Different media composition for <i>in vitro</i> culture	45
Table 2.3 The concentration combination of two growth regulators	46
Table 3.1 DNA contents and intraspecific genome size variation in <i>Zinnia</i> by PI staining and investigated by FACSCalibur	49
Table 3.2 DNA contents of <i>Zinnia</i> by PI staining and investigated by Partec PAS	52
Table 3.3 DNA contents of <i>Zinnia</i> by DAPI staining with and without 1% PVP and using Partec PAS.	53
Table 3.4 Relative AT-Specific Fluorescence (DF)	57
Table 3.5 The list of used primers	59
Table 3.6 The characteristics of guard cell, plant height and flower sizes of zinnia cultivars	63
Table 3.7 Chromosome numbers of zinnia species and cultivars	66
Table 3.8 Effect of media types on shoot and root formation in zinnia cultivars after 4 weeks in culture	68
Table 3.9 Effect of media types on callus induction in zinnia cultivars after 4 weeks in culture	69
Table 3.10 Effect of type of growth regulators on shoot and root formation in zinnia cultivars after 4 weeks in culture	73

## **LIST OF TABLES** (continued)

	<b>Page</b>
Table 3.11 Effect of type of growth regulators on callus induction in zinnia cultivars after 4 weeks in culture	74
Table 3.12 Effect of kinetin and IBA concentration on shoot and root formation in zinnia cultivars after 4 weeks in culture	78
Table 3.13 Effect of kinetin and IBA concentration on callus induction in zinnia cultivars after 4 weeks in culture	79
Table 3.14 Effect of AgNO <sub>3</sub> on shoot number, shoot height and root formation in three zinnia cultivars	84
Table 3.15 Effect of AgNO <sub>3</sub> on callus induction in zinnia cultivars after 4 weeks in culture	85
Table 4.1 Binding and spectral Properties of Selected DNA Fluorochromes	94

## LIST OF FIGURES

	<b>Page</b>
Figure 1.1 The formula structure of Myene and Germacrene D	6
Figure 1.2 Tracheary element (TE) formation of <i>in vitro</i> isolated <i>Zinnia</i> mesophyll cells	15
Figure 1.3 Example of ploidy analysis of banana ( <i>Musa acuminata</i> )	21
Figure 1.4 Example of genome size estimation of banana ( <i>Musa acuminata</i> )	23
Figure 1.5 Example of Chromosome analyze by flow cytometry	24
Figure 1.6 The process of random amplified polymorphic DNA (RAPD)	25
Figure 2.1 The seed size of <i>Z. angustifolia</i> cv. 'Starbright' (A), <i>Z. haageana</i> cv. 'Persian carpet' (B), <i>Z. elegans</i> cv. 'Dreamland' (C) and <i>Z. elegans</i> cv. 'Profusion' (D)	30
Figure 2.2 <i>Z. angustifolia</i> cv. 'Starbright'	32
Figure 2.3 <i>Z. haageana</i> cv. 'Persian carpet' and 'Chippendale daisy'	32
Figure 2.4 <i>Z. elegans</i> cv. 'Profusion'	32
Figure 2.5 <i>Z. elegans</i> cv. 'Sinnita', cv.'Short stuff', cv.'Dreamland', cv. 'Jupiter', cv. 'Piccolo' and cv. 'Peter pan'	33
Figure 2.6 <i>Z. elegans</i> cv. 'Border beauty', cv. 'Candy cane', cv. 'Jungle', cv. 'Dahlia',cv. 'Giant' and cv. 'Gold medal'	32
Figure 2.7 FACSCalibur in Prince of Songkla university	36
Figure 2.8 Partec PAS in The State Research Center Geisenheim, Germany	36
Figure 3.1 Histogram show <i>Raphanus sativus</i> (M1) compared with <i>Z. angustifolia</i>	50
Figure 3.2 FCM histogram showing combination peaks.	55

## LIST OF FIGURES (continued)

	Page
Figure 3.3 FCM histogram showing peaks of <i>Z. angustifolia</i> (M1) with <i>Z. elegans</i> cv. 'Dreamland' plus A) cv. 'Border beauty' B) cv. 'Giant' C) cv. 'Sinnita' D) cv. 'Jupiter' E) cv. 'Short stuff'	56
Figure 3.4 Primer screening in <i>Z. elegans</i> cv. 'Dreamland'	59
Figure 3.5 Electrophoretic analysis of amplification products obtained with the primer UBC 89	60
Figure 3.6 Dendrogram of <i>Z. elegans</i> cv. 'Dreamland' (Dr) and <i>Z. haageana</i> cv. 'Persian carpet' (Per) accessions (Bionumberic ver 3.0) based on cluster analysis of RAPDs with 5 primers.	61
Figure 3.7 Micrographs of guard cells in <i>Zinia</i>	64
Figure 3.8 Correlations between DNA content and some characteristics of zinnia	65
Figure 3.9 Chromosomes of <i>Z. angustifolia</i> cv. 'Starbright' (A), <i>Z. haageana</i> cv. 'Persian carpet' (B), <i>Z. elegans</i> cv. 'Peter pan' (C) and cv. 'Dreamland'(D)	66
Figure 3.10 <i>Z. angustifolia</i> cultured on MS, ½ MS, FK and KS media after 4 weeks of culture	70
Figure 3.11 <i>Z. haageana</i> cultured on MS, ½ MS, FK and KS media after 4 weeks of culture	70
Figure 3.12 <i>Z. elegans</i> cultured on MS, ½ MS, FK and KS media after 4 weeks of culture	70
Figure 3.13 <i>Z. angustifolia</i> in KS plus 1 µM BA (A), 1 µM BA + 0.1µM 2,4-D (B), 1 µM BA + 0.1µM IBA (C) and 1µM BA + 0.1 µM NAA (D)	75

## LIST OF FIGURES (continued)

	<b>Page</b>
Figure 3.14 <i>Z. angustifolia</i> in KS plus 1 $\mu$ M Kinetin(A), 1 $\mu$ M Kinetin + 0.1 $\mu$ M 2,4-D (B), 1 $\mu$ M Kinetin+0.1 $\mu$ M IBA(C) and 1 $\mu$ M Kinetin+0.1 $\mu$ M NAA(D)	74
Figure 3.15 <i>Z. haageana</i> in KS plus 1 $\mu$ M BA (A), 1 $\mu$ M BA + 0.1 $\mu$ M 2,4-D (B), 1 $\mu$ M BA + 0.1 $\mu$ M IBA (C) and 1 $\mu$ M BA + 0.1 $\mu$ M NAA (D)	76
Figure 3.16 <i>Z. haageana</i> in KS plus 1 $\mu$ M Kinetin(A), 1 $\mu$ M Kinetin + 0.1 $\mu$ M 2,4-D (B), 1 $\mu$ M Kinetin + 0.1 $\mu$ M IBA(C) and 1 $\mu$ M Kinetin+0.1 $\mu$ M NAA(D)	76
Figure 3.17 <i>Z. elegans</i> in KS plus 1 $\mu$ M BA (A), 1 $\mu$ M BA + 0.1 $\mu$ M 2,4-D(B), 1 $\mu$ M BA + 0.1 $\mu$ M IBA (C) and 1 $\mu$ M BA + 0.1 $\mu$ M NAA (D)	77
Figure 3.18 <i>Z. elegans</i> in KS plus 1 $\mu$ M Kinetin(A), 1 $\mu$ M Kinetin + 0.1 $\mu$ M 2,4-D (B), 1 $\mu$ M Kinetin + 0.1 $\mu$ M IBA(C) and 1 $\mu$ M Kinetin+0.1 $\mu$ M NAA(D)	77
Figure 3.19 <i>Z. angustifolia</i> in KS with various concentrations of kinetin and IBA	80
Figure 3.20 <i>Z. haageana</i> in KS with various concentrations of kinetin and IBA	81
Figure 3.21 <i>Z. elegans</i> in KS with various concentrations of kinetin and IBA	82
Figure 3.22 <i>Z. angustifolia</i> in KS with 5 $\mu$ M kinetin + 0.5 $\mu$ M IBA adding $\text{AgNO}_3$ 2, 4, 8 and 16 mg/l	86
Figure 3.23 <i>Z. haageana</i> in KS with 5 $\mu$ M kinetin + 0.5 $\mu$ M IBA adding $\text{AgNO}_3$ 2, 4, 8 and 16 mg/l	86
Figure 3.24 <i>Z. elegans</i> in KS with kinetin 5 $\mu$ M + IBA 0.5 $\mu$ M adding $\text{AgNO}_3$ 2, 4, 8 and 16 mg/l	86
Figure 3.25 Callus induced on MS medium supplemented with 0.1 $\mu$ M TDZ and 1 $\mu$ M TDZ	87

## **LIST OF FIGURES (continued)**

	<b>Page</b>
Figure3.26 Callus induced on MS medium supplemented with 17.6 $\mu\text{M}$ BA and 5.4 $\mu\text{M}$ NAA	88
Figure3.27 Callus induced on MS medium supplemented with 9 $\mu\text{M}$ 2, 4-D and 9.9 $\mu\text{M}$ IBA	88
Figure3.28 Organogenesis in <i>Z. elegans</i> cv. 'Dreamland'	89

## LIST OF ABBREVIATIONS

2,4-D	=	2,4-Dichlorophenoxyacetic acid
BA	=	N <sup>6</sup> benzylaminopurine
CV	=	Coefficient of variation
DAPI	=	4, 6-diamidino-2-phenylindole
DNA	=	Deoxyribonucleic acid
dNTP	=	Deoxyribonucleotide triphosphate
FCM	=	Flow Cytometry
FK	=	Fukuda & Komamine (1980)
KS	=	Kohlenbach & Schmidt (1975)
IBA	=	Idole butyric acid
IS	=	Internal standard
MS	=	Murashige and Skoog (1962)
NAA	=	1-Naphthaleneacetic acid
PCR	=	Polymerase Chain Reaction
pg	=	picogram
PI	=	Propidium iodide
PVP	=	Polyvinylpyrrolidone
RAPD	=	Random Amplified Polymorphic DNA
TDZ	=	Thidiazuron
TE	=	Trachea elementary