

BIBLIOGRAPHY

- Ahrens,M., Ankenbauer,T., Schroder,D., Hollnagel,A., Mayer,H., and Gross,G. (1993). Expression of human bone morphogenetic proteins-2 or -4 in murine mesenchymal progenitor C3H10T1/2 cells induces differentiation into distinct mesenchymal cell lineages. *DNA Cell Biol.* **12**, 871-880.
- Alden,T.D., Pittman,D.D., Hankins,G.R., Beres,E.J., Engh,J.A., Das,S., Hudson,S.B., Kerns,K.M., Kallmes,D.F., and Helm,G.A. (1999). In vivo endochondral bone formation using a bone morphogenetic protein 2 adenoviral vector. *Hum. Gene Ther.* **10**, 2245-2253.
- Anderson,K.M., Cheung,P.H., and Kelly,M.D. (1997). Rapid generation of homologous internal standards and evaluation of data for quantitation of messenger RNA by competitive polymerase chain reaction. *J. Pharmacol. Toxicol. Methods* **38**, 133-140.
- Aoki,H., Fujii,M., Imamura,T., Yagi,K., Takehara,K., Kato,M., and Miyazono,K. (2001). Synergistic effects of different bone morphogenetic protein type I receptors on alkaline phosphatase induction. *Journal of Cell Science* **114**, 1483-1489.
- Ashique,A.M., Fu,K., and Richman,J.M. (2002). Endogenous bone morphogenetic proteins regulate outgrowth and epithelial survival during avian lip fusion. *Development* **129**, 4647-4660.
- Ashton,B.A., Allen,T.D., Howlett,C.R., Eaglesom,C.C., Hattori,A., and Owen,M. (1980). Formation of bone and cartilage by marrow stromal cells in diffusion chambers in vivo. *Clin. Orthop.* **294**-307.
- Bahamonde,M.E. and Lyons,K.M. (2001). BMP3: to be or not to be a BMP. *J. Bone Joint Surg. Am.* **83-A Suppl 1**, S56-S62.
- Baur,S.T., Mai,J.J., and Dymecki,S.M. (2000). Combinatorial signaling through BMP receptor IB and GDF5: shaping of the distal mouse limb and the genetics of distal limb diversity. *Development (Cambridge, England)* **127**, 605-619.
- Bentz,H., Nathan,R.M., Rosen,D.M., Armstrong,R.M., Thompson,A.Y., Segarini,P.R., Mathews,M.C., Dasch,J.R., Piez,K.A., and Seyedin,S.M. (1989). Purification and characterization of a unique osteoinductive factor from bovine bone. *J. Biol. Chem.* **264**, 20805-20810.

- Bessho,K., Kusumoto,K., Fujimura,K., Konishi,Y., Ogawa,Y., Tani,Y., and Iizuka,T. (1999). Comparison of recombinant and purified human bone morphogenetic protein. *Br. J. Oral Maxillofac. Surg.* *37*, 2–5.
- Bodensteiner,K.J., Clay,C.M., Moeller,C.L., and Sawyer,H.R. (1999). Molecular cloning of the ovine Growth/Differentiation factor-9 gene and expression of growth/differentiation factor-9 in ovine and bovine ovaries. *Biol. Reprod.* *60*, 381–386.
- Bostrom,M.P., Lane,J.M., Berberian,W.S., Missri,A.A., Tomin,E., Weiland,A., Doty,S.B., Glaser,D., and Rosen,V.M. (1995). Immunolocalization and expression of bone morphogenetic proteins 2 and 4 in fracture healing. *J. Orthop. Res.* *13*, 357–367.
- Brommage,R. and DeLuca,H.F. (1985). Evidence that 1,25-dihydroxyvitamin D₃ is the physiologically active metabolite of vitamin D₃. *Endocr. Rev.* *6*, 491–511.
- Bruder,S.P., Fink,D.J., and Caplan,A.I. (1994). Mesenchymal stem cells in bone development, bone repair, and skeletal regeneration therapy. *J. Cell Biochem.* *56*, 283–294.
- Bucay,N., Sarosi,I., Dunstan,C.R., Morony,S., Tarpley,J., Capparelli,C., Scully,S., Tan,H.L., Xu,W., Lacey,D.L., Boyle,W.J., and Simonet,W.S. (1998). osteoprotegerin-deficient mice develop early onset osteoporosis and arterial calcification. *Genes Dev.* *12*, 1260–1268.
- Canalis,E. and Delany,A.M. (2002). Mechanisms of glucocorticoid action in bone. *Ann. N. Y. Acad. Sci.* *966*, 73–81.
- Canalis,E., Economides,A.N., and Gazzero,E. (2003). Bone morphogenetic proteins, their antagonists, and the skeleton. *Endocr. Rev.* *24*, 218–235.
- Carlevaro,M.F., Cermelli,S., Cancedda,R., and Descalzi,C.F. (2000). Vascular endothelial growth factor (VEGF) in cartilage neovascularization and chondrocyte differentiation: auto-paracrine role during endochondral bone formation. *J. Cell Sci.* *113* (Pt 1), 59–69.
- Carrington,J.L., Roberts,A.B., Flanders,K.C., Roche,N.S., and Reddi,A.H. (1988). Accumulation, localization, and compartmentation of transforming growth factor beta during endochondral bone development. *J. Cell Biol.* *107*, 1969–1975.
- Celeste,A.J., Iannazzi,J.A., Taylor,R.C., Hewick,R.M., Rosen,V., Wang,E.A., and Wozney,J.M. (1990). Identification of transforming growth factor beta family

- members present in bone-inductive protein purified from bovine bone. Proc. Natl. Acad. Sci. U. S. A **87**, 9843–9847.
- Celeste, A. J., Song, J. J., Cox, K., Rosen, V., and Wozney, J. M. Bone morphogenetic protein-9, a new member of the TGF- β superfamily. J. Bone Miner. Res. Suppl 1, S136. 1994.
- Ref Type: Abstract
- Chang,S.C., Hoang,B., Thomas,J.T., Vukicevic,S., Luyten,F.P., Ryba,N.J., Kozak,C.A., Reddi,A.H., and Moos,M., Jr. (1994). Cartilage-derived morphogenetic proteins. New members of the transforming growth factor-beta superfamily predominantly expressed in long bones during human embryonic development. J. Biol. Chem. **269**, 28227–28234.
- Chay,S.H., Rabie,A.B., and Itthagaran,A. (2000). Ultrastructural identification of cells involved in the healing of intramembranous bone grafts in both the presence and absence of demineralised intramembranous bone matrix. Aust. Orthod. J. **16**, 88–97.
- Cheng,H., Jiang,W., Phillips,F.M., Haydon,R.C., Peng,Y., Zhou,L., Luu,H.H., An,N., Breyer,B., Vanichakarn,P., Szatkowski,J.P., Park,J.Y., and He,T.C. (2003). Osteogenic activity of the fourteen types of human bone morphogenetic proteins (BMPs). J. Bone Joint Surg. Am. **85-A**, 1544–1552.
- Cheng,S.L., Yang,J.W., Rifas,L., Zhang,S.F., and Avioli,L.V. (1994). Differentiation of human bone marrow osteogenic stromal cells in vitro: induction of the osteoblast phenotype by dexamethasone. Endocrinology **134**, 277–286.
- Chuen,F.S., Chuk,C.Y., Ping,W.Y., Nar,W.W., Kim,H.L., and Ming,C.K. (2004). Immunohistochemical characterization of cells in adult human patellar tendons. J. Histochem. Cytochem. **52**, 1151–1157.
- Chung,U.I., Lanske,B., Lee,K., Li,E., and Kronenberg,H. (1998). The parathyroid hormone/parathyroid hormone-related peptide receptor coordinates endochondral bone development by directly controlling chondrocyte differentiation. Proc. Natl. Acad. Sci. U. S. A **95**, 13030–13035.
- Clarke,T.R., Hoshiya,Y., Yi,S.E., Liu,X., Lyons,K.M., and Donahoe,P.K. (2001). Mullerian Inhibiting Substance Signaling Uses a Bone Morphogenetic Protein (BMP)-Like Pathway Mediated by ALK2 and Induces Smad6 Expression. Mol Endocrinol **15**, 946–959.

- Clemens,T.L., Garrett,K.P., Zhou,X.Y., Pike,J.W., Haussler,M.R., and Dempster,D.W. (1988). Immunocytochemical localization of the 1,25-dihydroxyvitamin D₃ receptor in target cells. *Endocrinology* **122**, 1224–1230.
- Cohn,D.V. and Wong,G.L. (1979). Isolated bone cells. In *Skeletal research*, D.J.Simmons and A.S.Kunin, eds. (New York: Academic Press), pp. 3–20.
- Cosman,F., Nieves,J., Woelfert,L., Formica,C., Gordon,S., Shen,V., and Lindsay,R. (2001). Parathyroid hormone added to established hormone therapy: effects on vertebral fracture and maintenance of bone mass after parathyroid hormone withdrawal. *J. Bone Miner. Res.* **16**, 925–931.
- Daluiski,A., Engstrand,T., Bahamonde,M.E., Gamer,L.W., Agius,E., Stevenson,S.L., Cox,K., Rosen,V., and Lyons,K.M. (2001). Bone morphogenetic protein-3 is a negative regulator of bone density. *Nat. Genet.* **27**, 84–88.
- Dean,C., Ito,M., Makarenkova,H.P., Faber,S.C., and Lang,R.A. (2004). Bmp7 regulates branching morphogenesis of the lacrimal gland by promoting mesenchymal proliferation and condensation. *Development* **131**, 4155–4165.
- Derynck,R., Jarrett,J.A., Chen,E.Y., Eaton,D.H., Bell,J.R., Assoian,R.K., Roberts,A.B., Sporn,M.B., and Goeddel,D.V. (1985). Human transforming growth factor-beta complementary DNA sequence and expression in normal and transformed cells. *Nature* **316**, 701–705.
- Domenicucci,C., Goldberg,H.A., Hofmann,T., Isenman,D., Wasi,S., and Sodek,J. (1988). Characterization of porcine osteonectin extracted from foetal calvariae. *Biochem. J.* **253**, 139–151.
- Dube,J.L., Wang,P., Elvin,J., Lyons,K.M., Celeste,A.J., and Matzuk,M.M. (1998). The bone morphogenetic protein 15 gene is X-linked and expressed in oocytes. *Mol. Endocrinol.* **12**, 1809–1817.
- Ducy,P. and Karsenty,G. (1998). Genetic control of cell differentiation in the skeleton. *Curr. Opin. Cell Biol.* **10**, 614–619.
- Ducy,P., Zhang,R., Geoffroy,V., Ridall,A.L., and Karsenty,G. (1997). Osf2/Cbfα1: a transcriptional activator of osteoblast differentiation. *Cell* **89**, 747–754.
- Dudl,R.J., Ensinck,J.W., Baylink,D., Chesnut,C.H., III, Sherrard,D., Nelp,W.B., and Palmieri,G.M. (1973). Evaluation of intravenous calcium as therapy for osteoporosis. *Am. J. Med.* **55**, 631–637.

- Duneas,N., Crooks,J., and Ripamonti,U. (1998). Transforming growth factor-beta 1: induction of bone morphogenetic protein genes expression during endochondral bone formation in the baboon, and synergistic interaction with osteogenic protein-1 (BMP-7). *Growth Factors* 15, 259-277.
- Eckery,D.C., Whale,L.J., Lawrence,S.B., Wylde,K.A., McNatty,K.P., and Juengel,J.L. (2002). Expression of mRNA encoding growth differentiation factor 9 and bone morphogenetic protein 15 during follicular formation and growth in a marsupial, the brushtail possum (*Trichosurus vulpecula*). *Mol Cell Endocrinol* 192, 115-126.
- Eisman,J.A. (2001). Pharmacogenetics of the vitamin D receptor and osteoporosis. *Drug Metab Dispos.* 29, 505-512.
- Engstrand,T., Daluiski,A., Bahamonde,M.E., Melhus,H., and Lyons,K.M. (2000). Transient production of bone morphogenetic protein 2 by allogeneic transplanted transduced cells induces bone formation. *Hum. Gene Ther.* 11, 205-211.
- Erenpreisa,J. and Roach,H.I. (1996). Epigenetic selection as a possible component of transdifferentiation. Further study of the commitment of hypertrophic chondrocytes to become osteocytes. *Mech. Ageing Dev.* 87, 165-182.
- Fang,J. and Hall,B.K. (1997). Chondrogenic cell differentiation from membrane bone periosteal. *Anat. Embryol. (Berl)* 196, 349-362.
- Ferguson,C.M., Schwarz,E.M., Puzas,J.E., Zuscik,M.J., Drissi,H., and O'Keefe,R.J. (2004). Transforming growth factor-beta1 induced alteration of skeletal morphogenesis in vivo. *J. Orthop. Res.* 22, 687-696.
- Finelli,A.L., Xie,T., Bossie,C.A., Blackman,R.K., and Padgett,R.W. (1995). The tolkin gene is a tolloid/BMP-1 homologue that is essential for *Drosophila* development. *Genetics* 141, 271-281.
- Fisher,L.W. and Termine,J.D. (1985). Noncollagenous proteins influencing the local mechanisms of calcification. *Clin. Orthop.* 362-385.
- Franceschi,R.T. and Iyer,B.S. (1992). Relationship between collagen synthesis and expression of the osteoblast phenotype in MC3T3-E1 cells. *J. Bone Miner. Res.* 7, 235-246.
- Franceschi,R.T., Wang,D., Krebsbach,P.H., and Rutherford,R.B. (2000). Gene therapy for bone formation: in vitro and in vivo osteogenic activity of an adenovirus expressing BMP7. *J. Cell Biochem.* 78, 476-486.

- Friedenstein,A.J. (1980). Stromal mechanisms of bone marrow: cloning in vitro and retransplantation in vivo. *Haematol. Blood Transfus.* **25**, 19–29.
- Fu,S.C., Wong,Y.P., Chan,B.P., Pau,H.M., Cheuk,Y.C., Lee,K.M., and Chan,K.M. (2003). The roles of bone morphogenetic protein (BMP) 12 in stimulating the proliferation and matrix production of human patellar tendon fibroblasts. *Life Sci.* **72**, 2965–2974.
- Furuya,K., Nifuji,A., Rosen,V., and Noda,M. (1999). Effects of GDF7/BMP12 on proliferation and alkaline phosphatase expression in rat osteoblastic osteosarcoma ROS 17/2.8 cells. *J. Cell Biochem.* **72**, 177–180.
- Gajavelli,S., Wood,P.M., Pennica,D., Whittemore,S.R., and Tsoufas,P. (2004). BMP signaling initiates a neural crest differentiation program in embryonic rat CNS stem cells. *Exp. Neurol.* **188**, 205–223.
- Galloway,S.M., McNatty,K.P., Cambridge,L.M., Laitinen,M.P., Juengel,J.L., Jokiranta,T.S., McLaren,R.J., Luiro,K., Dodds,K.G., Montgomery,G.W., Beattie,A.E., Davis,G.H., and Ritvos,O. (2000). Mutations in an oocyte-derived growth factor gene (BMP15) cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nat. Genet.* **25**, 279–283.
- Gitelman,S.E., Kobrin,M.S., Ye,J.Q., Lopez,A.R., Lee,A., and Derynck,R. (1994). Recombinant Vgr-1/BMP-6-expressing tumors induce fibrosis and endochondral bone formation in vivo. *J. Cell Biol.* **126**, 1595–1609.
- Goldberg,H.A., Domenicucci,C., Pringle,G.A., and Sodek,J. (1988a). Mineral-binding proteoglycans of fetal porcine calvarial bone. *J. Biol. Chem.* **263**, 12092–12101.
- Goldberg,H.A., Maeno,M., Domenicucci,C., Qi,Z.N., and Sodek,J. (1988b). Identification of small collagenous proteins with properties of procollagen alpha 1 (I) pN-propeptide in fetal porcine calvarial bone. *Coll. Relat Res.* **8**, 187–197.
- Goumans,M.J., Valdimarsdottir,G., Itoh,S., Rosendahl,A., Sideras,P., and ten Dijke,P. (2002). Balancing the activation state of the endothelium via two distinct TGF- β type I receptors. *EMBO J.* **21**, 1743–1753.
- Graf,D., Nethsinghe,S., Palmer,D.B., Fisher,A.G., and Merkenschlager,M. (2002). The developmentally regulated expression of Twisted gastrulation reveals a role for bone morphogenetic proteins in the control of T cell development. *J. Exp. Med.* **196**, 163–171.

- Gugala,Z., Olmsted-Davis,E.A., Gannon,F.H., Lindsey,R.W., and Davis,A.R. (2003). Osteoinduction by ex vivo adenovirus-mediated BMP2 delivery is independent of cell type. *Gene Ther.* **10**, 1289–1296.
- Gundle,R., Joyner,C.J., and Triffitt,J.T. (1995). Human bone tissue formation in diffusion chamber culture in vivo by bone-derived cells and marrow stromal fibroblastic cells. *Bone* **16**, 597–601.
- Guterman,I.A., Boman,T.E., Wang,G.J., and Balian,G. (1988). Bone induction in intramuscular implants by demineralized bone matrix: sequential changes of collagen synthesis. *Coll. Relat Res.* **8**, 419–431.
- Gygi,S.P., Rochon,Y., Franzia,B.R., and Aebersold,R. (1999). Correlation between protein and mRNA abundance in yeast. *Mol. Cell Biol.* **19**, 1720–1730.
- Hager-Theodorides,A.L., Outram,S.V., Shah,D.K., Sacedon,R., Shrimpton,R.E., Vicente,A., Varas,A., and Crompton,T. (2002). Bone morphogenetic protein 2/4 signalling regulates early thymocyte differentiation. *J. Immunol.* **169**, 5496–5504.
- Hanahan,D. (1983). Studies on transformation of Escherichia coli with plasmids. *J. Mol. Biol.* **166**, 557–580.
- Hanamura,H., Higuchi,Y., Nakagawa,M., Iwata,H., Nogami,H., and Urist,M.R. (1980). Solubilized bone morphogenetic protein (BMP) from mouse osteosarcoma and rat demineralized bone matrix. *Clin. Orthop.* **281**–290.
- Hardesty,R.A. and Marsh,J.L. (1990). Craniofacial onlay bone grafting: a prospective evaluation of graft morphology, orientation, and embryonic origin. *Plast. Reconstr. Surg.* **85**, 5–14.
- Heldin,C.H., Miyazono,K., and ten Dijke,P. (1997). TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* **390**, 465–471.
- Helm,G.A., Alden,T.D., Beres,E.J., Hudson,S.B., Das,S., Engh,J.A., Pittman,D.D., Kerns,K.M., and Kallmes,D.F. (2000a). Use of bone morphogenetic protein-9 gene therapy to induce spinal arthrodesis in the rodent. *J. Neurosurg.* **92**, 191–196.
- Helm,G.A., Alden,T.D., Sheehan,J.P., and Kallmes,D. (2000b). Bone morphogenetic proteins and bone morphogenetic protein gene therapy in neurological surgery: a review. *Neurosurgery* **46**, 1213–1222.
- Herring,G.M. (1968). The chemical structure of tendon, cartilage, dentin and bone matrix. *Clin. Orthop.* **60**, 261–299.

- Hill,P.A., Reynolds,J.J., and Meikle,M.C. (1994). The purification and partial characterization of bone resorptive polypeptides from bovine bone matrix. *Biochim. Biophys. Acta* **1201**, 193–202.
- Hiraki,Y., Inoue,H., Shigeno,C., Sanma,Y., Bentz,H., Rosen,D.M., Asada,A., and Suzuki,F. (1991). Bone morphogenetic proteins (BMP-2 and BMP-3) promote growth and expression of the differentiated phenotype of rabbit chondrocytes and osteoblastic MC3T3-E1 cells in vitro. *J. Bone Miner. Res.* **6**, 1373–1385.
- Hogan,B.L. (1996). Bone morphogenetic proteins in development. *Curr. Opin. Genet. Dev.* **6**, 432–438.
- Hou,L.T., Liu,C.M., Chen,Y.J., Wong,M.Y., Chen,J., and Chen,K.C. (2000). Gel-filtrated fractions of alveolar bone extract contain factors promoting cell attachment and a mitogenic effect on periodontal ligament fibroblasts. *Proc. Natl. Sci. Counc. Repub. China B* **24**, 33–40.
- Hughes,F.J., Collyer,J., Stanfield,M., and Goodman,S.A. (1995). The effects of bone morphogenetic protein-2, -4, and -6 on differentiation of rat osteoblast cells in vitro. *Endocrinology* **136**, 2671–2677.
- Inoda,H., Yamamoto,G., and Hattori,T. (2004). Histological investigation of osteoinductive properties of rh-BMP2 in a rat calvarial bone defect model. *J. Craniomaxillofac. Surg.* **32**, 365–369.
- Isaacs,N.W. (1995). Cystine knots. *Curr. Opin. Struct. Biol.* **5**, 391–395.
- Iwasaki,S., Hattori,A., Sato,M., Tsujimoto,M., and Kohno,M. (1996). Characterization of the bone morphogenetic protein-2 as a neurotrophic factor. Induction of neuronal differentiation of PC12 cells in the absence of mitogen-activated protein kinase activation. *J. Biol. Chem.* **271**, 17360–17365.
- Jane,J.A., Jr., Dunford,B.A., Kron,A., Pittman,D.D., Sasaki,T., Li,J.Z., Li,H., Alden,T.D., Dayoub,H., Hankins,G.R., Kallmes,D.F., and Helm,G.A. (2002a). Ectopic osteogenesis using adenoviral bone morphogenetic protein (BMP)-4 and BMP-6 gene transfer. *Mol. Ther.* **6**, 464–470.
- Jane,J.A., Jr., Dunford,B.A., Kron,A., Pittman,D.D., Sasaki,T., Li,J.Z., Li,H., Alden,T.D., Dayoub,H., Hankins,G.R., Kallmes,D.F., and Helm,G.A. (2002b). Ectopic osteogenesis using adenoviral bone morphogenetic protein (BMP)-4 and BMP-6 gene transfer. *Mol Ther.* **6**, 464–470.

- Jochum,W., David,J.P., Elliott,C., Wutz,A., Plenk,H., Jr., Matsuo,K., and Wagner,E.F. (2000). Increased bone formation and osteosclerosis in mice overexpressing the transcription factor Fra-1. *Nat. Med.* **6**, 980–984.
- Juang,R.H., Chang,Y.D., Sung,H.Y., and Su,J.C. (1984). Oven-drying method for polyacrylamide gel slab packed in cellophane sandwich. *Anal. Biochem.* **141**, 348–350.
- Kanatani,M., Sugimoto,T., Kaji,H., Kobayashi,T., Nishiyama,K., Fukase,M., Kumegawa,M., and Chihara,K. (1995). Stimulatory effect of bone morphogenetic protein-2 on osteoclast-like cell formation and bone-resorbing activity. *J. Bone Miner. Res.* **10**, 1681–1690.
- Karsenty,G. (1999). The genetic transformation of bone biology. *Genes Dev.* **13**, 3037–3051.
- Katagiri,T., Yamaguchi,A., Komaki,M., Abe,E., Takahashi,N., Ikeda,T., Rosen,V., Wozney,J.M., Fujisawa-Sehara,A., and Suda,T. (1994). Bone morphogenetic protein-2 converts the differentiation pathway of C2C12 myoblasts into the osteoblast lineage. *J. Cell Biol.* **127**, 1755–1766.
- Katz,R.W. and Reddi,A.H. (1988). Dissociative extraction and partial purification of osteogenin, a bone inductive protein, from rat tooth matrix by heparin affinity chromatography. *Biochem. Biophys. Res. Commun.* **157**, 1253–1257.
- Kawabata,M., Imamura,T., and Miyazono,K. (1998). Signal transduction by bone morphogenetic proteins. *Cytokine Growth Factor Rev.* **9**, 49–61.
- Kim,H.J., Rice,D.P., Kettunen,P.J., and Thesleff,I. (1998). FGF-, BMP- and Shh-mediated signalling pathways in the regulation of cranial suture morphogenesis and calvarial bone development. *Development* **125**, 1241–1251.
- Kim,N., Odgren,P.R., Kim,D.K., Marks,S.C., Jr., and Choi,Y. (2000). Diverse roles of the tumor necrosis factor family member TRANCE in skeletal physiology revealed by TRANCE deficiency and partial rescue by a lymphocyte-expressed TRANCE transgene. *Proc. Natl. Acad. Sci. U. S. A* **97**, 10905–10910.
- Kim,R.Y., Robertson,E.J., and Solloway,M.J. (2001). Bmp6 and Bmp7 are required for cushion formation and septation in the developing mouse heart. *Dev. Biol.* **235**, 449–466.
- Kingsley,D.M. (1994). What do BMPs do in mammals? Clues from the mouse short-ear mutation. *Trends Genet.* **10**, 16–21.

- Kingsley,D.M., Bland,A.E., Grubber,J.M., Marker,P.C., Russell,L.B., Copeland,N.G., and Jenkins,N.A. (1992). The mouse short ear skeletal morphogenesis locus is associated with defects in a bone morphogenetic member of the TGF beta superfamily. *Cell* **71**, 399–410.
- Kirsch,T., Nickel,J., and Sebald,W. (2000). BMP-2 antagonists emerge from alterations in the low-affinity binding epitope for receptor BMPR-II. *EMBO J.* **19**, 3314–3324.
- Komori,T., Yagi,H., Nomura,S., Yamaguchi,A., Sasaki,K., Deguchi,K., Shimizu,Y., Bronson,R.T., Gao,Y.H., Inada,M., Sato,M., Okamoto,R., Kitamura,Y., Yoshiki,S., and Kishimoto,T. (1997). Targeted disruption of Cbfa1 results in a complete lack of bone formation owing to maturational arrest of osteoblasts. *Cell* **89**, 755–764.
- Kumagai,T., Lee,I., Ono,Y., Maeno,M., and Takagi,M. (1998). Ultrastructural localization and biochemical characterization of vitronectin in developing rat bone. *Histochem. J.* **30**, 111–119.
- Kuznetsov,S.A., Friedenstein,A.J., and Robey,P.G. (1997). Factors required for bone marrow stromal fibroblast colony formation in vitro. *Br. J. Haematol.* **97**, 561–570.
- Lacey,D.L., Timms,E., Tan,H.L., Kelley,M.J., Dunstan,C.R., Burgess,T., Elliott,R., Colombero,A., Elliott,G., Scully,S., Hsu,H., Sullivan,J., Hawkins,N., Davy,E., Capparelli,C., Eli,A., Qian,Y.X., Kaufman,S., Sarosi,I., Shalhoub,V., Senaldi,G., Guo,J., Delaney,J., and Boyle,W.J. (1998). Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* **93**, 165–176.
- Laemmli,U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Laitinen,M., Vuojolainen,K., Jaatinen,R., Ketola,I., Aaltonen,J., Lehtonen,E., Heikinheimo,M., and Rittos,O. (1998). A novel growth differentiation factor-9 (GDF-9) related factor is co-expressed with GDF-9 in mouse oocytes during folliculogenesis. *Mech. Dev.* **78**, 135–140.
- Lecoeur,L. and Ouhayoun,J.P. (1997). In vitro induction of osteogenic differentiation from non-osteogenic mesenchymal cells. *Biomaterials* **18**, 989–993.

- Lee,I., Ono,Y., Lee,A., Omiya,K., Moriya,Y., and Takagi,M. (1998). Immunocytochemical localization and biochemical characterization of large proteoglycans in developing rat bone. *J. Oral Sci.* **40**, 77-87.
- Liem,K.F., Jr., Tremml,G., Roelink,H., and Jessell,T.M. (1995). Dorsal differentiation of neural plate cells induced by BMP-mediated signals from epidermal ectoderm. *Cell* **82**, 969-979.
- Long,F., Schipani,E., Asahara,H., Kronenberg,H., and Montminy,M. (2001). The CREB family of activators is required for endochondral bone development. *Development* **128**, 541-550.
- Luyten,F.P., Cunningham,N.S., Ma,S., Muthukumaran,N., Hammonds,R.G., Nevins,W.B., Woods,W.I., and Reddi,A.H. (1989). Purification and partial amino acid sequence of osteogenin, a protein initiating bone differentiation. *J. Biol. Chem.* **264**, 13377-13380.
- Luyten,F.P., Cunningham,N.S., Vukicevic,S., Paralkar,V., Ripamonti,U., and Reddi,A.H. (1992). Advances in osteogenin and related bone morphogenetic proteins in bone induction and repair. *Acta Orthop. Belg.* **58 Suppl 1**, 263-267.
- Macias,D., Ganan,Y., Sampath,T.K., Piedra,M.E., Ros,M.A., and Hurle,J.M. (1997). Role of BMP-2 and OP-1 (BMP-7) in programmed cell death and skeletogenesis during chick limb development. *Development* **124**, 1109-1117.
- Maeno,M., Oikawa,E., Taguchi,M., Suzuki,N., Shiraishi,Y., Hayashi,A., Yahagi,N., Katayama,I., and Otsuka,K. (1993). Characteristics of a 28-kDa collagenous protein extracted with guanidine from EDTA-demineralized rabbit alveolar bone. *J. Nihon Univ Sch Dent.* **35**, 258-266.
- Manolagas,S.C. (2000). Birth and death of bone cells: basic regulatory mechanisms and implications for the pathogenesis and treatment of osteoporosis. *Endocr. Rev.* **21**, 115-137.
- Marden,L.J., Hollinger,J.O., Chaudhari,A., Turek,T., Schaub,R.G., and Ron,E. (1994). Recombinant human bone morphogenetic protein-2 is superior to demineralized bone matrix in repairing craniotomy defects in rats. *J. Biomed. Mater. Res.* **28**, 1127-1138.
- Mason,A.J., Hayflick,J.S., Ling,N., Esch,F., Ueno,N., Ying,S.Y., Guillemin,R., Niall,H., and Seburg,P.H. (1985). Complementary DNA sequences of ovarian follicular

- fluid inhibin show precursor structure and homology with transforming growth factor-beta. *Nature* **318**, 659–663.
- Massague,J. (1996). TGFbeta signaling: receptors, transducers, and Mad proteins. *Cell* **85**, 947–950.
- McGrath,S.A., Esquela,A.F., and Lee,S.J. (1995). Oocyte-specific expression of growth/differentiation factor-9. *Mol Endocrinol* **9**, 131–136.
- McKibbin,B. (1978). The biology of fracture healing in long bones. *J. Bone Joint Surg. Br.* **60-B**, 150–162.
- Miller,A.F., Harvey,S.A., Thies,R.S., and Olson,M.S. (2000). Bone morphogenetic protein-9. An autocrine/paracrine cytokine in the liver. *J. Biol. Chem.* **275**, 17937–17945.
- Miyazono,K., ten Dijke,P., and Heldin,C.H. (2000). TGF-beta signaling by Smad proteins. *Adv. Immunol.* **75**, 115–157.
- Moskalewski,S., Osiecka,A., and Malejczyk,J. (1988). Comparison of bone formed intramuscularly after transplantation of scapular and calvarial osteoblasts. *Bone* **9**, 101–106.
- Nah,H.D., Pacifici,M., Gerstenfeld,L.C., Adams,S.L., and Kirsch,T. (2000). Transient chondrogenic phase in the intramembranous pathway during normal skeletal development. *J. Bone Miner. Res.* **15**, 522–533.
- Nakahara,H., Bruder,S.P., Goldberg,V.M., and Caplan,A.I. (1990). In vivo osteochondrogenic potential of cultured cells derived from the periosteum. *Clin. Orthop.* **223**–232.
- Nakashima,K., Zhou,X., Kunkel,G., Zhang,Z., Deng,J.M., Behringer,R.R., and de Crombrugghe,B. (2002). The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* **108**, 17–29.
- Nonner,D., Barrett,E.F., Kaplan,P., and Barrett,J.N. (2001). Bone morphogenetic proteins (BMP6 and BMP7) enhance the protective effect of neurotrophins on cultured septal cholinergic neurons during hypoglycemia. *J. Neurochem.* **77**, 691–699.
- Otsuka,E., Yamaguchi,A., Hirose,S., and Hagiwara,H. (1999). Characterization of osteoblastic differentiation of stromal cell line ST2 that is induced by ascorbic acid. *Am. J. Physiol* **277**, C132–C138.

- Otsuka,F., Yao,Z., Lee,T., Yamamoto,S., Erickson,G.F., and Shimasaki,S. (2000). Bone morphogenetic protein-15. Identification of target cells and biological functions. *J. Biol. Chem.* **275**, 39523–39528.
- Otto,F., Thornell,A.P., Crompton,T., Denzel,A., Gilmour,K.C., Rosewell,I.R., Stamp,G.W., Beddington,R.S., Mundlos,S., Olsen,B.R., Selby,P.B., and Owen,M.J. (1997). Cbfa1, a candidate gene for cleidocranial dysplasia syndrome, is essential for osteoblast differentiation and bone development. *Cell* **89**, 765–771.
- Ozkaynak,E., Rueger,D.C., Drier,E.A., Corbett,C., Ridge,R.J., Sampath,T.K., and Oppermann,H. (1990). OP-1 cDNA encodes an osteogenic protein in the TGF-beta family. *EMBO J.* **9**, 2085–2093.
- Ozkaynak,E., Schnegelsberg,P.N., Jin,D.F., Clifford,G.M., Warren,F.D., Drier,E.A., and Oppermann,H. (1992). Osteogenic protein-2. A new member of the transforming growth factor-beta superfamily expressed early in embryogenesis. *J. Biol. Chem.* **267**, 25220–25227.
- Paralkar,V.M., Nandedkar,A.K., Pointer,R.H., Kleinman,H.K., and Reddi,A.H. (1990). Interaction of osteogenin, a heparin binding bone morphogenetic protein, with type IV collagen. *J. Biol. Chem.* **265**, 17281–17284.
- Peretto,P., Cummings,D., Modena,C., Behrens,M., Venkatraman,G., Fasolo,A., and Margolis,F.L. (2002). BMP mRNA and protein expression in the developing mouse olfactory system. *J. Comp Neurol.* **451**, 267–278.
- Prober,J.M., Trainor,G.L., Dam,R.J., Hobbs,F.W., Robertson,C.W., Zagursky,R.J., Cocuzza,A.J., Jensen,M.A., and Baumeister,K. (1987). A system for rapid DNA sequencing with fluorescent chain-terminating dideoxynucleotides. *Science* **238**, 336–341.
- Prockop,D.J. (1997). Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science* **276**, 71–74.
- Rabie,A.B. (1997). Molecular and biochemical advances in the field of clinical bone induction. In *Bone formation and repair*, A.B.Rabie and M.R.Urist, eds. (Amsterdam; Elsevier), pp. 89–100.
- Rabie,A.B., Dan,Z., and Samman,N. (1996). Ultrastructural identification of cells involved in the healing of intramembranous and endochondral bones. *Int. J. Oral Maxillofac. Surg.* **25**, 383–388.

- Reddi,A.H. (1981). Cell biology and biochemistry of endochondral bone development. *Coll. Relat Res.* **1**, 209–226.
- Reddi,A.H. (1983). Extracellular bone matrix dependent local induction of cartilage and bone. *J. Rheumatol. Suppl* **11**, 67–69.
- Reddi,A.H. and Cunningham,N.S. (1990). Bone induction by osteogenin and bone morphogenetic proteins. *Biomaterials* **11**, 33–34.
- Reddi,A.H. and Huggins,C. (1972). Biochemical sequences in the transformation of normal fibroblasts in adolescent rats. *Proc. Natl. Acad. Sci. U. S. A* **69**, 1601–1605.
- Reddi,A.H., Muthukumaran,N., Ma,S., Carrington,J.L., Luyten,F.P., Paralkar,V.M., and Cunningham,N.S. (1989). Initiation of bone development by osteogenin and promotion by growth factors. *Connect. Tissue Res.* **20**, 303–312.
- Reissmann,E., Ernsberger,U., Francis-West,P.H., Rueger,D., Brickell,P.M., and Rohrer,H. (1996). Involvement of bone morphogenetic protein-4 and bone morphogenetic protein-7 in the differentiation of the adrenergic phenotype in developing sympathetic neurons. *Development* **122**, 2079–2088.
- Ripamonti,U., Crooks,J., Matsaba,T., and Tasker,J. (2000). Induction of endochondral bone formation by recombinant human transforming growth factor-beta2 in the baboon (*Papio ursinus*). *Growth Factors* **17**, 269–285.
- Ripamonti,U., Duneas,N., Van Den,H.B., Bosch,C., and Crooks,J. (1997). Recombinant transforming growth factor-beta1 induces endochondral bone in the baboon and synergizes with recombinant osteogenic protein-1 (bone morphogenetic protein-7) to initiate rapid bone formation. *J. Bone Miner. Res.* **12**, 1584–1595.
- Ripamonti,U., Ma,S., Cunningham,N.S., Yeates,L., and Reddi,A.H. (1992). Initiation of bone regeneration in adult baboons by osteogenin, a bone morphogenetic protein. *Matrix* **12**, 369–380.
- Ripamonti,U., Ramoshebi,L.N., Matsaba,T., Tasker,J., Crooks,J., and Teare,J. (2001). Bone induction by BMPs/OPs and related family members in primates. *J. Bone Joint Surg. Am.* **83-A Suppl 1**, S116–S127.
- Risteli,L. and Risteli,J. (1993). Biochemical markers of bone metabolism. *Ann. Med.* **25**, 385–393.
- Roach,H.I. (1992). Trans-differentiation of hypertrophic chondrocytes into cells capable of producing a mineralized bone matrix. *Bone Miner.* **19**, 1–20.

- Roach,H.I. and Erenpreisa,J. (1996). The phenotypic switch from chondrocytes to bone-forming cells involves asymmetric cell division and apoptosis. *Connect. Tissue Res.* **35**, 85-91.
- Roach,H.I., Erenpreisa,J., and Aigner,T. (1995). Osteogenic differentiation of hypertrophic chondrocytes involves asymmetric cell divisions and apoptosis. *J. Cell Biol.* **131**, 483-494.
- Rodan,G.A. and Martin,T.J. (1981). Role of osteoblasts in hormonal control of bone resorption--a hypothesis. *Calcif. Tissue Int.* **33**, 349-351.
- Ruppert,R., Hoffmann,E., and Sebald,W. (1996). Human bone morphogenetic protein 2 contains a heparin-binding site which modifies its biological activity. *Eur. J. Biochem.* **237**, 295-302.
- Sampath,T.K., Coughlin,J.E., Whetstone,R.M., Banach,D., Corbett,C., Ridge,R.J., Ozkaynak,E., Oppermann,H., and Rueger,D.C. (1990). Bovine osteogenic protein is composed of dimers of OP-1 and BMP-2A, two members of the transforming growth factor-beta superfamily. *J. Biol. Chem.* **265**, 13198-13205.
- Sampath,T.K., DeSimone,D.P., and Reddi,A.H. (1982). Extracellular bone matrix-derived growth factor. *Exp. Cell Res.* **142**, 460-464.
- Sampath,T.K., Maliakal,J.C., Hauschka,P.V., Jones,W.K., Sasak,H., Tucker,R.F., White,K.H., Coughlin,J.E., Tucker,M.M., Pang,R.H., and . (1992). Recombinant human osteogenic protein-1 (hOP-1) induces new bone formation in vivo with a specific activity comparable with natural bovine osteogenic protein and stimulates osteoblast proliferation and differentiation in vitro. *J. Biol. Chem.* **267**, 20352-20362.
- Sampath,T.K., Muthukumaran,N., and Reddi,A.H. (1987). Isolation of osteogenin, an extracellular matrix-associated, bone-inductive protein, by heparin affinity chromatography. *Proc. Natl. Acad. Sci. U. S. A* **84**, 7109-7113.
- Scammell,B.E. and Roach,H.I. (1996). A new role for the chondrocyte in fracture repair: endochondral ossification includes direct bone formation by former chondrocytes. *J. Bone Miner. Res.* **11**, 737-745.
- Schwartz,Z., Weesner,T., van Dijk,S., Cochran,D.L., Mellonig,J.T., Lohmann,C.H., Carnes,D.L., Goldstein,M., Dean,D.D., and Boyan,B.D. (2000). Ability of deproteinized cancellous bovine bone to induce new bone formation. *J. Periodontol.* **71**, 1258-1269.

- Scott,C.K., Bain,S.D., and Hightower,J.A. (1994). Intramembranous bone matrix is osteoinductive. *Anat. Rec.* **238**, 23–30.
- Scott,C.K. and Hightower,J.A. (1991). The matrix of endochondral bone differs from the matrix of intramembranous bone. *Calcif. Tissue Int.* **49**, 349–354.
- Shimasaki,S., Moore,R.K., Otsuka,F., and Erickson,G.F. (2004). The bone morphogenetic protein system in mammalian reproduction. *Endocr. Rev.* **25**, 72–101.
- Shore,E.M., Cook,A.L., Hahn,G.V., Kaplan,F.S., Wozney,J.M., Wagner,M.J., and Wells,D.E. (1995). BMP-1 sublocalization on human chromosome 8. Molecular anatomy and orthopaedic implications. *Clin. Orthop.* **199**–209.
- Simonet,W.S., Lacey,D.L., Dunstan,C.R., Kelley,M., Chang,M.S., Luthy,R., Nguyen,H.Q., Wooden,S., Bennett,L., Boone,T., Shimamoto,G., DeRose,M., Elliott,R., Colombero,A., Tan,H.L., Trail,G., Sullivan,J., Davy,E., Bucay,N., Renshaw-Gegg,L., Hughes,T.M., Hill,D., Pattison,W., Campbell,P., Boyle,W.J., and . (1997). Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell* **89**, 309–319.
- Smith,J.D. and Abramson,M. (1974). Membranous vs endochondrial bone autografts. *Arch. Otolaryngol.* **99**, 203–205.
- Smith,L.M., Sanders,J.Z., Kaiser,R.J., Hughes,P., Dodd,C., Connell,C.R., Heiner,C., Kent,S.B., and Hood,L.E. (1986). Fluorescence detection in automated DNA sequence analysis. *Nature* **321**, 674–679.
- Somi,S., Buffing,A.A., Moorman,A.F., and Van Den Hoff,M.J. (2004). Dynamic patterns of expression of BMP isoforms 2, 4, 5, 6, and 7 during chicken heart development. *Anat. Rec. A Discov. Mol Cell Evol. Biol.* **279**, 636–651.
- Song,J.J., Celeste,A.J., Kong,F.M., Jirtle,R.L., Rosen,V., and Thies,R.S. (1995). Bone morphogenetic protein-9 binds to liver cells and stimulates proliferation. *Endocrinology* **136**, 4293–4297.
- Spelsberg,T.C., Subramaniam,M., Riggs,B.L., and Khosla,S. (1999). The actions and interactions of sex steroids and growth factors/cytokines on the skeleton. *Mol. Endocrinol.* **13**, 819–828.
- Strawich,E. and Glimcher,M.J. (1983). Differences in the extent and heterogeneity of lysyl hydroxylation in embryonic chick cranial and long bone collagens. *J. Biol. Chem.* **258**, 555–562.

- Suda,N. (1997). Parathyroid hormone-related protein (PTHrP) as a regulating factor of endochondral bone formation. *Oral Dis.* *3*, 229–231.
- Sugimoto,T., Kanatani,M., Kaji,H., Yamaguchi,T., Fukase,M., and Chihara,K. (1993). Second messenger signaling of PTH- and PTHRP-stimulated osteoclast-like cell formation from hemopoietic blast cells. *Am. J. Physiol* *265*, E367–E373.
- Sullivan,W.G. and Szwajkun,P. (1991). Membranous versus endochondral bone. *Plast. Reconstr. Surg.* *87*, 1145.
- Suzuki,F. (1996). Roles of cartilage matrix proteins, chondromodulin-I and -II, in endochondral bone formation: a review. *Connect. Tissue Res.* *35*, 303–307.
- Suzuki,K. (1990). [Purification and characterization of a growth factor from human bone matrix]. *Nippon Seikeigeka Gakkai Zasshi* *64*, 928–936.
- Suzuki,N., Labosky,P.A., Furuta,Y., Hargett,L., Dunn,R., Fogo,A.B., Takahara,K., Peters,D.M., Greenspan,D.S., and Hogan,B.L. (1996). Failure of ventral body wall closure in mouse embryos lacking a procollagen C-proteinase encoded by Bmp1, a mammalian gene related to *Drosophila tolloid*. *Development* *122*, 3587–3595.
- Syftestad,G.T., Triffitt,J.T., Urist,M.R., and Caplan,A.I. (1984). An osteo-inductive bone matrix extract stimulates the in vitro conversion of mesenchyme into chondrocytes. *Calcif. Tissue Int.* *36*, 625–627.
- Takaoka,K., Yoshikawa,H., Shimizu,N., Ono,K., Amitani,K., and Nakata,Y. (1982). Partial purification of bone-inducing substances from a murine osteosarcoma. *Clin. Orthop.* *265*–270.
- Termine,J.D., Belcourt,A.B., Conn,K.M., and Kleinman,H.K. (1981). Mineral and collagen-binding proteins of fetal calf bone. *J. Biol. Chem.* *256*, 10403–10408.
- Termine,J.D., Belcourt,A.B., Miyamoto,M.S., and Conn,K.M. (1980). Properties of dissociatively extracted fetal tooth matrix proteins. II. Separation and purification of fetal bovine dentin phosphoprotein. *J. Biol. Chem.* *255*, 9769–9772.
- Thielemann,F.W., Schmidt,K., and Koslowski,L. (1982). Osteoinduction. Part II: Purification of the osteoinductive activities of bone matrix. *Arch. Orthop. Trauma Surg.* *100*, 73–78.
- Thomson,J.A., Itskovitz-Eldor,J., Shapiro,S.S., Waknitz,M.A., Swiergiel,J.J., Marshall,V.S., and Jones,J.M. (1998). Embryonic stem cell lines derived from human blastocysts. *Science* *282*, 1145–1147.

- Uchida,T. (1994). [Purification and characterization of growth factor extracted from bone]. Kokubyo Gakkai Zasshi *61*, 44–55.
- Urist,M.R. (1965). Bone: formation by autoinduction. Science *150*, 893–899.
- Urist,M.R., Huo,Y.K., Brownell,A.G., Hohl,W.M., Buyske,J., Lietze,A., Tempst,P., Hunkapiller,M., and DeLange,R.J. (1984). Purification of bovine bone morphogenetic protein by hydroxyapatite chromatography. Proc. Natl. Acad. Sci. U. S. A *81*, 371–375.
- Urist,M.R., Sato,K., Brownell,A.G., Malinin,T.I., Lietze,A., Huo,Y.K., Prolo,D.J., Oklund,S., Finerman,G.A., and DeLange,R.J. (1983). Human bone morphogenetic protein (hBMP). Proc. Soc. Exp. Biol. Med. *173*, 194–199.
- Vidal,O., Lindberg,M.K., Hollberg,K., Baylink,D.J., Andersson,G., Lubahn,D.B., Mohan,S., Gustafsson,J.A., and Ohlsson,C. (2000). Estrogen receptor specificity in the regulation of skeletal growth and maturation in male mice. Proc. Natl. Acad. Sci. U. S. A *97*, 5474–5479.
- Visser,J.A., Olaso,R., Verhoef-Post,M., Kramer,P., Themmen,A.P.N., and Ingraham,H.A. (2001). The Serine/Threonine Transmembrane Receptor ALK2 Mediates Mullerian Inhibiting Substance Signaling. Mol Endocrinol *15*, 936–945.
- Volek-Smith,H. and Urist,M.R. (1996). Recombinant human bone morphogenetic protein (rhBMP) induced heterotopic bone development in vivo and in vitro. Proc. Soc. Exp. Biol. Med. *211*, 265–272.
- Vukicevic,S., Latin,V., Chen,P., Batorsky,R., Reddi,A.H., and Sampath,T.K. (1994). Localization of osteogenic protein-1 (bone morphogenetic protein-7) during human embryonic development: high affinity binding to basement membranes. Biochem. Biophys. Res. Commun. *198*, 693–700.
- Wagner,D.S. and Mullins,M.C. (2002). Modulation of BMP activity in dorsal–ventral pattern formation by the chordin and noggin antagonists. Dev. Biol. *245*, 109–123.
- Waite,K.A. and Eng,C. (2003). From developmental disorder to heritable cancer: it's all in the BMP/TGF-beta family. Nat. Rev. Genet. *4*, 763–773.
- Wang,E.A., Rosen,V., Cordes,P., Hewick,R.M., Kriz,M.J., Luxenberg,D.P., Sibley,B.S., and Wozney,J.M. (1988). Purification and characterization of other distinct bone-inducing factors. Proc. Natl. Acad. Sci. U. S. A *85*, 9484–9488.
- Wang,E.A., Rosen,V., D'Alessandro,J.S., Bauduy,M., Cordes,P., Harada,T., Israel,D.I., Hewick,R.M., Kerns,K.M., LaPan,P., and . (1990). Recombinant human bone

- morphogenetic protein induces bone formation. Proc. Natl. Acad. Sci. U. S. A **87**, 2220–2224.
- Waterborg,J.H. and Matthews,H.R. (1996). The Lowry Method for Protein Quantitation. In The Protein Protocols Handbook, J.M.Walker, ed. (Totowa: Humana Press Inc.), pp. 7–9.
- Weber,K. and Osborn,M. (1969). The reliability of molecular weight determinations by dodecyl sulfate–polyacrylamide gel electrophoresis. J. Biol. Chem. **244**, 4406–4412.
- Wendel,M., Sommarin,Y., and Heinegard,D. (1998). Bone matrix proteins: isolation and characterization of a novel cell-binding keratan sulfate proteoglycan (osteoadherin) from bovine bone. J. Cell Biol. **141**, 839–847.
- Weryha,G. and Leclerc,J. (1995). Paracrine regulation of bone remodeling. Horm. Res. **43**, 69–75.
- Wharton,K.A., Thomsen,G.H., and Gelbart,W.M. (1991). Drosophila 60A gene, another transforming growth factor beta family member, is closely related to human bone morphogenetic proteins. Proc. Natl. Acad. Sci. U. S. A **88**, 9214–9218.
- Windahl,S.H., Vidal,O., Andersson,G., Gustafsson,J.A., and Ohlsson,C. (1999). Increased cortical bone mineral content but unchanged trabecular bone mineral density in female ERbeta(−/−) mice. J. Clin. Invest **104**, 895–901.
- Wong,R.W. and Rabie,A.B. (1999). A quantitative assessment of the healing of intramembranous and endochondral autogenous bone grafts. Eur. J. Orthod. **21**, 119–126.
- Wordinger,R.J., Agarwal,R., Talati,M., Fuller,J., Lambert,W., and Clark,A.F. (2002). Expression of bone morphogenetic proteins (BMP), BMP receptors, and BMP associated proteins in human trabecular meshwork and optic nerve head cells and tissues. Mol. Vis. **8**, 241–250.
- Wozney,J.M. (1989). Bone morphogenetic proteins. Prog. Growth Factor Res. **1**, 267–280.
- Wozney,J.M. (1992). The bone morphogenetic protein family and osteogenesis. Mol. Reprod. Dev. **32**, 160–167.
- Wozney,J.M. and Rosen,V. (1998). Bone morphogenetic protein and bone morphogenetic protein gene family in bone formation and repair. Clin. Orthop. **26**–37.

- Wozney,J.M., Rosen,V., Celeste,A.J., Mitsock,L.M., Whitters,M.J., Kriz,R.W., Hewick,R.M., and Wang,E.A. (1988). Novel regulators of bone formation: molecular clones and activities. *Science* **242**, 1528–1534.
- Wright,V., Peng,H., Usas,A., Young,B., Gearhart,B., Cummins,J., and Huard,J. (2002). BMP4-expressing muscle-derived stem cells differentiate into osteogenic lineage and improve bone healing in immunocompetent mice. *Mol Ther.* **6**, 169–178.
- Xiao,G., Gopalakrishnan,R., Jiang,D., Reith,E., Benson,M.D., and Franceschi,R.T. (2002). Bone morphogenetic proteins, extracellular matrix, and mitogen- activated protein kinase signaling pathways are required for osteoblast- specific gene expression and differentiation in MC3T3-E1 cells. *J. Bone Miner. Res.* **17**, 101–110.
- Yamaguchi,A., Komori,T., and Suda,T. (2000). Regulation of osteoblast differentiation mediated by bone morphogenetic proteins, hedgehogs, and Cbfa1. *Endocr. Rev.* **21**, 393–411.
- Yamashita,H., ten Dijke,P., Heldin,C.H., and Miyazono,K. (1996). Bone morphogenetic protein receptors. *Bone* **19**, 569–574.
- Yan,C., Wang,P., DeMayo,J., DeMayo,F.J., Elvin,J.A., Carino,C., Prasad,S.V., Skinner,S.S., Dunbar,B.S., Dube,J.L., Celeste,A.J., and Matzuk,M.M. (2001). Synergistic roles of bone morphogenetic protein 15 and growth differentiation factor 9 in ovarian function. *Mol Endocrinol* **15**, 854–866.
- Yasuda,H., Shima,N., Nakagawa,N., Yamaguchi,K., Kinosaki,M., Mochizuki,S., Tomoyasu,A., Yano,K., Goto,M., Murakami,A., Tsuda,E., Morinaga,T., Higashio,K., Udagawa,N., Takahashi,N., and Suda,T. (1998). Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc. Natl. Acad. Sci. U. S. A* **95**, 3597–3602.
- Ying,Y., Liu,X.M., Marble,A., Lawson,K.A., and Zhao,G.Q. (2000). Requirement of Bmp8b for the generation of primordial germ cells in the mouse. *Mol Endocrinol* **14**, 1053–1063.
- Yu,Y.M., Sklar,M.M., Nissley,S.P., and Reddi,A.H. (1993). Changes in the expression of insulin-like growth factor II/mannose-6-phosphate receptor during endochondral bone development. *Biochem. Biophys. Res. Commun.* **195**, 516–524.

- Zhao,G.Q. (2003). Consequences of knocking out BMP signaling in the mouse. *Genesis.* *35*, 43–56.
- Zhao,G.Q., Deng,K., Labosky,P.A., Liaw,L., and Hogan,B.L. (1996). The gene encoding bone morphogenetic protein 8B is required for the initiation and maintenance of spermatogenesis in the mouse. *Genes Dev.* *10*, 1657–1669.
- Zhao,G.Q. and Hogan,B.L. (1996). Evidence that mouse Bmp8a (Op2) and Bmp8b are duplicated genes that play a role in spermatogenesis and placental development. *Mech. Dev.* *57*, 159–168.
- Zhao,H., Ettala,O., and Vaananen,H.K. (2002). Intracellular membrane trafficking pathways in bone-resorbing osteoclasts revealed by cloning and subcellular localization studies of small GTP-binding rab proteins. *Biochem. Biophys. Res. Commun.* *293*, 1060–1065.
- Zhu,W., Rawlins,B.A., Boachie-Adjei,O., Myers,E.R., Arimizu,J., Choi,E., Lieberman,J.R., Crystal,R.G., and Hidaka,C. (2004). Combined bone morphogenetic protein-2 and -7 gene transfer enhances osteoblastic differentiation and spine fusion in a rodent model. *J. Bone Miner. Res.* *19*, 2021–2032.
- Zins,J.E. and Whitaker,L.A. (1983). Membranous versus endochondral bone: implications for craniofacial reconstruction. *Plast. Reconstr. Surg.* *72*, 778–785.
- Zoricic,S., Maric,I., Bobinac,D., and Vukicevic,S. (2003). Expression of bone morphogenetic proteins and cartilage-derived morphogenetic proteins during osteophyte formation in humans. *J. Anat.* *202*, 269–277.