

BIBLIOGRAPHY

- Alder, M.N., Dames, S., Gaudet, J. and Mango, S.E. 2003. Gene silencing in *Caenorhabditis elegans* by transitive RNA interference. *RNA* 9: 25-32.
- Algar E., Khromykh T., Smith S., Blackburn D., Bryson G. and Smith P. 1996. A WT1 antisense oligonucleotide inhibits proliferation and induces apoptosis in myeloid leukemia cell lines. *Oncogene* 12: 1005-14.
- Amin, K.M., Litzky, L.A., Smythe, W.R., Mooney, A.M., Morris, J.M., Mews, D.J.Y., Pass, H.I., Kari, C., Rodeck, U., Rauscher, F.J., Kaiser, I.R. and Albelda, S.M. 1995. Wilms' tumor (WT1) susceptibility gene products are selectively expressed in malignant mesothelioma. *Am J Pathol.* 146: 344-56.
- Anondalakshmi, R., Marathe, R., Ge, X., Herr, Jr., J.M., Mau, C., Mallory, A., Pruss, G., Bowman, L. and Vance, V.B. 2000. A calmodulin-related protein that suppress posttranscriptional gene silencing in plants. *Science* 290: 142-4.
- Bardeesy, N., Zabel, B., Schmitt, K. and Pelletier, J. 1994. WT1 mutations associated with incomplete Denys-Drash syndrome define a domain predicted to behave in a dominant-negative fashion. *Genomics* 21: 663-4.
- Bernstein, E., Caudy, A.A., Hammond, S.M. and Hannon, G.J. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409: 363-6.
- Bickmore, W.A., Oghene, K., Little, M.H., Seawright, A., van Heyningen, V. and Hastie, N.D. 1992. Modulating DNA binding specificity by alternative splicing of the Wilms'tumor wt1 gene transcript. *Science* 257: 235-7.
- Boik, John. 1996. Cancer and natural medicine : a textbook of basic science and clinical research. Oregon Medical Press, Minnesota, USA.
- Bridge, A.J., Pebernard, S., Ducraux, A., Nicoulaz, A.L., and Iggo, R. 2003. Induction of an interferon response by RNAi vectors in mammalian cells. *Nat. Genet.* 8: 8.
- Brinton, L., Lacey, J., Devesa, S.S., 2002. Epidemiology of breast cancer. In: Konegan, W.L., Spratt, J.S. (Eds.), *Cancer of the Breast*, (5th edR) WB Saunders, Philadelphia, p.111-32.

- Bruening, W., Gros, P., Sato, T., Stanimir, J., Nakamura, Y., Housman, D. and Pelletier, J. 1993. Analysis of the 11p13 Wilms tumor suppressor gene (WT1) in ovarian tumors. *Cancer Invest.* 11: 393-9.
- Bruening, W. And Pelletier, J. 1996. A non-AUG translation initiation event generates novel WT1 isoforms. *J Biol Chem.* 271: 8646-54.
- Buell, P. 1973 Changing incidence of breast cancer in Japanese-American women. *J Natl Cancer Inst.* 51: 1479-83.
- Burwell, E.A., McCarty, G.P., Simpson, L.A., Thomson, K.A. and Loeb, D.M. 2006. Isoforms of Wilms' tumor suppressor gene (WT1) have distinct effects on mammary epithelial cells. *Oncogene* 11; [Epub ahead of print].
- Call, K.M., Glaser, T., Ito, C.Y., Buckler, A.J., Pelletier, J., Haber, D.A., Rose, E.A., Kral, A., Yeger, H., Lewis, W.H., Jones, C. and Housman, D.E. 1990. Isolation and characterization of zinc finger polypeptide gene at the chromosome 11 Wilms' tumor locus. *Cell* 60: 509-20.
- Chen, Z. 2001. The possible role and application of WT1 in human leukemia. *Int. J. Hematol.* 73: 39-46.
- Chicas, A. and Macino, G. 2001. Characteristics of post-transcriptional gene silencing. *EMBO Rep.* 2: 992-6.
- Cogoni, C. 2001. Homology-dependent gene silencing mechanisms in fungi. *Annu. Rev. Microbial.* 55: 381-406.
- Coppes, M.J., Campbell, C.E. and Williams, B.R.G. 1993. The role of WT1 in Wilms tumorigenesis. *FASEB J.* 7: 886-95.
- Czauderna, F., Fechtner, M., Dames, S., Agun, H., Klippel, A., Pronk, G.J., Giese, K. and Kaufmann, J. 2003. Structural variations and stabilizing modifications of synthetic siRNAs in mammalian cells. *Nucleic Acid Res.* 31: 2705-16.
- Dalmay, T., Hamilton, A., Rudd, S., Angell, S., and aulcombe, D.C. 2000. An RNA-dependent RNA polymerase gene in *Arabidopsis* is required for posttranscriptional gene silencing mediated by a transgene but not by a virus. *Cell* 101: 543-53.

- Dalmay, T., Horsefield, R., Braunstein, T.H. and Baulcombe, D.C. 2001. SDE3 encodes and RNA helicase required for post-transcriptional gene silencing in *Arabidopsis*. *EMBO J.* 20: 2069-78.
- Dawson, P.J., Wolman, S.R., Tait, L., Heppner, G.H. and Miller, F.R. 1996. MCF10AT: a model for the evolution of cancer from proliferative breast disease. *Am J Pathol.* 148: 313-9.
- Dey, B.R., Sukhatme, V.P., Roberts, A.B., Sporn, M.B., Rauscher, F.J.III and Kim, S.J. 1994. Repression of the transforming growth factor - β 1 gene by the Wilms' tumor suppressor WT1 gene product. *Mol Endocrinol.* 8: 595-602.
- Drummond, I.A., Madden, S.L., Rohwer-Nutter, P., Bell, G.I., Sukhatme, V.P., Rauscher, F.J.III. 1992. Repression of the insulin-like growth factor II gene by the Wilms tumor suppressor WT1. *Science* 257: 674-8.
- Dupont, W.D., Parl, F.F., Hartmann, W.H., Brinton, L.A., Winfield, A.C., Worrell, J.A., Schuyler, P.A. and Plummer, W.D. 1993. Breast cancer risk associated with proliferative breast disease and a typical hyperplasia. *Cancer* 71: 1258-65.
- Dzitoyeva, S., Dimitrijevic, N. and Manev, H., 2001. Intra-abdominal injection of double-stranded RNA into anesthetized adult *Drosophila* trigger RNA interference in the central nervous system. *Mol. Psychiatry.* 6: 665-70.
- Easton, D.F., Ford, D. and Bishop, D.T. 1995. Breast and ovarian cancer incidence in BRCA1 mutation carriers. Breast Cancer Linkage Consortium. *Am J Hum Genet.* 56: 265-71.
- Elbashir, S.M., Lendeckel, W. and Tuschl, T. 2001a. RNA interference is mediated by 21- and 22-nucleotide RNAs. The authors demonstrate that synthetic siRNA can initiate RNAi in *Drosophila* embryo extracts. *Genes Dev.* 15: 188-200.
- Elbashir, S.M., Harborth, J., Lendeckel, W., Yalcin, A., Weber, K. and Tuschl, T. 2001b. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 411: 494-8.
- Elbashir, S.M., Harborth, J., Weber, K. and Tuschl, T. 2002. Analysis of gene function in somatic mammalian cells using small interfering RNAs. *Methods* 26: 199-213.
- Englert, C., Hou, X., Maheswaran, S., Bennett, P., Ngwu, C., Re, G.G., Garvin, A.J., Rosner, M.R. and Haber, D.A. 1995. WT1 suppresses synthesis of the epidermal growth factor receptor and induces apoptosis. *EMBO J.* 14: 4662-75.

- Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E. and Mello, C.C. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391: 806-11.
- Freshney, R.I., 1994. Culture of Animal Cells: A Manual of Basic Technique. 3rd Ed. Wiley-Liss. New York. USA
- Gashler, A. L., Bonthon, D. T., Madden, S. L., Rauscher, F. J., III, Collins, T. and Sukhatme, V. P. 1992. Human platelet-derived growth factor A chain is transcriptionally repressed by the Wilms tumor suppressor WT1. *Proc. Natl. Acad. Sci. USA*. 89: 10984-8.
- Gessler, M., Poustka, A., Cavenee, W., Neve, R. L., Orkin, S. H. and Bruns, G. A. 1990. Homozygous deletion in Wilms tumours of a zinc-finger gene identified by chromosome jumping. *Nature* 34: 774-8.
- Gessler, M., Konig, A. and Bruns, G.A. 1992. The genomic organization and expression of the WT1 gene. *Genomics* 12: 807-13.
- Graidist, P., Phongdara, A. and Fujise, K. 2004. Antiapoptotic protein partners fortilin and MCL1 independently protect cells from 5-Fluorouracil-induced cytotoxicity. *J Biol Chem* 279: 40868-75.
- Haber, D.A., Park, S., Maheswaran, S., Englert, C., Re, G.G., Hazen-Martin, D.J., Sens, D.A. and Garvin, A.J. 1993. WT1-mediated growth suppression of Wilms tumor cells expressing a WT1 splicing variant. *Science* 262: 2057-9.
- Hamilton, T.B., Barilla, K.C. and Ronaniuk, P.J. 1995. High affinity binding site for the Wilms' tumor suppressor protein WT1. *Nucleic Acids Res.* 23: 277-84.
- Hammilton, A.J. and Baulcombe, D.C. 1999. A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science* 286: 950-2.
- Hammond, S.M., Bernstein, E., Beach, D. and Hannon, G.J. 2000. An RNA directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* 404:293 – 296.
- Hannon, G.J. 2002 . RNA interference. *Nature* 418: 244-51.
- Harris J.R., Lippman M.E., Veronesi, U. and Eillett, w. 1992. Breast cancer. *N Engl J Med.* 327: 473-80.
- Hasuwa, H., Kaseda, K., Einarsdottir, T., Okabe, M. 2002. Small interfering RNA and gene silencing in transgenic mice and rats. *FEBS Lett.* 532: 227-30.

- Hewitt, S.M., Hamada, S., McDonnell, T.J., Rauscher, F.J.III and Saunders, G.F. 1995. Regulation of the proto-oncogene bcl-2 and c-myc by the Wilms' tumor suppressor gene WT1. *Cancer Res.* 55: 5386-9.
- Hsiao, W.C., Young, K.C., Lin, S.L. and Lin, P.W. 2004. Estrogen receptor-alpha polymorphism in a Taiwanese clinical breast cancer population: a case control study. *Breast Cancer Res.* 6: 180-6.
- Hutvagner, G. and Zamore, P.D. 2002. A microRNA in a multiple-turnover RNAi enzyme complex. *Science* 297: 2056-60.
- Inoue, K., Sugiyama, H., Ogawa, H., Nakagawa, M., Yamagami, T., Miwa, H., Kita, K., Hiraoka, A., Masaoka, T., Nasu, K., Kyo, T., Dohy, H., Nakauchi, H., Ishidate, T., Akiyama, T. and Kishimoto, T., 1994. WT1 as a new prognostic factor and a new marker for the detection of minimal residual disease in acute leukemia. *Blood* 84: 3071-9.
- Inoue, K., Tamaki, H., Ogawa, H., Oka, Y., Soma, T., Tatekawa, T., Oji, Y., Tsuboi, A., Kim, E. H., Kawakami, M., Akiyama, T., Kishimoto, T. and Sugiyama, H. 1998. Wilms' tumor gene (*WT1*) competes with differentiation-inducing signal in hematopoietic progenitor cells. *Blood* 91: 2969-76.
- Jensen, S., Gassama, M.P. and Heidemann, T. 1999. Taming of transposable elements by homology-dependent gene silencing. *Nat. Genet.* 21: 209-12.
- Johnstone, R.W., See, R.H., Sells, S.F., Wang, J., Muthukumar, S., Englert, C., Haber, D.A., Licht, J.D., Sugrue, S.P., Roberts, T., Rangnekar, V.M. and Shi, Y. 1996. A novel repressor, par-4, modulates transcription and growth suppressor function of the Wilms' tumor suppressor WT1. *Mol Cell Biol.* 19: 6945-56.
- Junqueira, L., Carneiro, J. and Kelley, R. 1995. The female reproductive system. In: Basic Histology, 8 th ed, pp. 443-6.
- Kalidas, S. and Smith, D.P. 2002. Novel genomic cDNA hybrids produce effective RNA interference in adult *Drosophila*. *Neuron* 33: 177-84.
- Kamath, R.S., Martinez-Campos, M., Zipperlen, P., Fraser, A.G. and Ahringer, J. 2001. Effectiveness of specific RNA-mediated interference through ingested double-stranded RNA in *Caenorhabditis elegans*. *Genome Biol.* 2: 1-10.

- Kennerdell, J.R. and Carthew, R.W. 1998. Use of dsRNA-mediated genetic interference to demonstrate that *frizzled* and *frizzled 2* act in the wingless pathway. *Cell* 95: 1017-26.
- Ketting, R.F., Haverkamp, T.H., van Luenen, H.G. and Plasterk, R.H. 1999. Mut-7 of *C. elegans*, required for transposon silencing and RNA interference, is a homolog of Werner syndrome helicase and RNaseD. *Cell* 99: 133-41.
- King-Underwood, L., Renshaw, J. and Pritchard-Jones, K. 1996. Mutations in the Wilms' tumor gene WT1 in leukemias. *Blood* 87: 2171-9.
- Kumar, M. and Carmichael, G.G. 1998. Antisense RNA: function and fate of duplex RNA in cells of higher eukaryote. *Microbiol. Mol. Biol. Rev.* 62: 1415-34.
- Kumar V., Cotran, R. and Robbins, S. 1997. Female genital system and breast. In: Basic Pathology, 6 th ed WB Saunders Co, Philadelphia. pp. 623-35.
- Kurreck, J. 2003. Antisense technologies: improvement through novel chemical modification. *FEBS*. 270: 1628-44.
- Lee, S.S., Lee, R.Y., Fraser, A.G., Kamath, R.S., Ahringer, J. and Ruvkun, G. 2003. A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat. Genet.* 33: 40-8.
- Loeb, D.M., Evron, E., Patel, C.B., Sharma, P.M., Niranjan, B., Buluwela, L., Weitzman, S.A., Korz, D. and Sukumar, S. 2001. Wilms' tumor suppressor gene (WT1) is expressed in primary breast tumor despite tumor specific promoter methylation. *Cancer Res.* 61: 921-5.
- London, S.J., Cannolly, J.L., Schnitt, S.J. and Colditz, G.A. 1992. A prospective study of benign breast disease and the risk of breast cancer. *Jama* 267: 941-4.
- Lucy, A.P., Guo, H.S., Li, W.X. and Ding, S.W. 2000. Suppression of post-transcriptional gene silencing by a plant viral protein localized in the nucleus. *EMBO J.* 19: 1672-80.
- McManus, M.T. and Sharp, P.A. 2002. Gene silencing in mammals by small interfering RNAs. *Nat Rev Genet* 3: 737-47.
- Madden, S.L., Cook, D.M., Morris, J.F., Gashler, A., Sukhatme, V.P. and Rauscher, F.J. III. 1991. Transcriptional repression mediated by the WT Wilms' tumor gene product. *Science* 253: 1550-2.

- Madden, S.L., Cook, D.M. and Rauscher, F.J.III. 1993. A structure-function analysis of transcriptional repression mediated by the WT1, Wilms' tumor suppressor protein. *Oncogene* 8: 1713-20.
- Maheswaran, S., Park, S., Bernard, A., Morris, J.F., Rausche, F.J., Hill, D.F. and Haber, D.A. 1993. Physical and functional interaction between WT1 and p53 proteins. *Proc Natl Acad Sci USA*. 90: 5100-4.
- Maheswaran, S., Englert, C., Bennett, P., Heinrich, G., Haber, D.A. 1995. The WT1 gene product stabilizes p53 and inhibits p53-mediated apoptosis. *Gene Dev.* 9: 2143-56.
- Malinsky, S., Bucheton, A. and Busseau, I. 2000. New insights on homology-dependent silencing of I factor activity by transgenes containing ORF1 in *Drosophila melanogaster*. *Genetics* 156: 1147-55.
- Malkin, D., Li, F.P., Strong, L.C., Fraumeni, J.F. Jr., Nelson, C.E., Kim, D.H. Kassel, J., Gryka, M.A., Bischoff, F.Z. and Tainsky, M.A. 1990. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* 250: 1233-8.
- Martinez, J., Patkaniowska, A., Urlaub, H., Luhrmann, R. and Tuschl, T. 2002. Single-stranded antisense siRNAs guide target RNA cleavage in RNAi. *Cell* 110: 563-74.
- Menke, A.L., Shvarts, A., Riteco, N., Van Ham, R.C., Van der Eb, A.J. and Jochemsen, A.G. 1997. Wilms' tumor 1-KTS isoforms induce p53-independent apoptosis that can be partially rescued by expression of the epidermal growth factor receptor or the insulin receptor. *Cancer Res.* 57: 1353-63.
- Menssen, H.D., Renkl, H.L., Rodeck, U. Maaurer, J., Notter, M., Schwartz, S., Reinhardt, R. and thieli, E. 1995. Presence of Wilms'tumor gene (WT1) transcripts and the WT1 nuclear protein in the majority of human acute leukemias. *Leukemia* 9: 1060-7.
- Miyoshi, Y., Ando, A., Egawa, C., Taguchi, T., Tamaki, Y., Tamaki, H., Sugiyama, H. and Noguchi, S. 2002. High expression of Wilms' tumor suppressor gene predicts poor prognosis in breast cancer patients. *Clin Cancer Res.* 8: 1167-71.
- Morris, J.F., Madden, S.L., Tournay, O.E., Cook, D.M., Sukhatme, V.P. and Rauscher, F.J. 1991. Characterization of the zinc finger protein encoded by the WT1 Wilms' tumor locus. *Oncogene* 6: 2339-48.

- Musgrove, E., Hui, R., Sweeney, K., Watts, C. and Sutherland, R. 1996. Cyclins and breast cancer. *J. Mamm. Gland Biol. Neoplasia* 1: 153–62.
- Naama, R., Sharon, M., Rive, S., Shirley, A., Wade, V.W., Edward, M.C., Adrian, V.L. and Haim, W. 2005. The WT1 wilms'tumor suppressor gene product interacts with estrogen receptor-alpha and regulates IGF-I receptor gene transcription in breast cancer cells. *J Mol Endo.* 35: 135-44.
- Nakagama, H., Heinrich, G., Pelletier, J. and Housman, D.E. 1995. Sequence and structural requirements for high affinity DNA binding by the WT1 gene product. *Mol Cell Biol.* 15: 1489-98.
- Napoli, C., Lemieux, C. and Jorgensen, R.A. 1990. Introduciton fo a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes in trans. *Plant Cell.* 2: 279-89.
- Nykanen, A., Haley, B., and Zamore, P.D. 2001. ATP requirements and small interfering RNA structure in the RNA interference pathway. *Cell* 107: 309-21.
- Oelgeschlager, M., Larrain, J., Geissert, D. and De Robertis, E.M. 2000. The evolutionarily consered BMP-binding protein Twisted gastrulation promotes BMP signaling. *Nature* 405: 757-63.
- Page, D.L. Dupont, W.D. Rogers, L.W. and Rados, M.S. 1985. Atypical hyperplastic lesions of the female breast. A long-term follow-up study. *Cancer* 55: 2698-708.
- Park, S., Schalling, M., Bernard, A., Maheswaran, S., Shipley, G.C., Roberts, D., Flectcher, J., Shipman, R., Rheinwald, J., Demetri, G., Griffin, J., Minden, M., Housmanm and Haber, D.A. 1993. The Wilms'tumor gene WT1 is expressed in murine mesoderm-derived tissues and mutated in a human mesothelioma. *Nat Genet.* 4: 415-20.
- Park, W., Li, J., Song, R., Messing, J. and Chen, X. 2002. CARPEL FACTORY, a dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in *Arabidopsis thaliana*. *Curr. Biol.* 12: 1484-95.
- Parkin, D.M., Bray, F.I. and Devesa, S.S. 2001. Cancer burden in the year 2000. The global picture. *Eur J Cancer.* 8: 54-66.

- Parrish, S., Fleenor, J., Xu, S., Mello, C. and Fire, A. 2000. Functional anatomy of a dsRNA trigger: differential requirement for the two trigger strands in RNA interference. *Mol Cell*, 6: 1077-87.
- Pelletier, J., Bruening, W., Kashtan, C.E., Mauer, S.M., Manivel, J.C., Striegel, J.E., Houghton, D.C., Junien, C., Habib, R. and Fouser, L. 1991. Germline mutations in the Wilms' tumor suppressor gene are associated with abnormal urogenital development in Denys-Drash syndrome. *Cell* 67: 437-47.
- Pritchard-Jones, K., Fleming, S., Davidson, D., Bickmore, W., Porteous, D., Gosden, C., Bard, J., Buckler, A., Pelletier, J., and Housman, D. 1990. The candidate Wilms' tumor gene is involved in genitourinary development. *Nature* 346: 194-7.
- Ratcliff, F.G., MacFarlane, S.A. and Baulcombe, D.C. 1999. Gene silencing without DNA: RNA-mediated cross-protection between viruses. *Plant Cell*. 11: 1207-16.
- Rauscher, F.J.III, Morris, J.F., Tournay, O.E., Cook, D.M. and Curren,T. 1990. Binding of the Wilms'tumor locus zinc finger protein to the Egr-1 consensus sequence. *Science* 25: 1259-62.
- Resnicoff, M., Burgaud, J.L., Rotman, H.L., Abraham, D. and Baserga, R. 1995. Correlation between apoptosis, tumorigenesis, and the level of insulin growth factor I receptor. *Cancer Res.* 55: 3739-41.
- Rodeck, U., Bossler, A., Kari, C., Humpray, C.W., Gyorfi, T., Maurer, J., Thiel, E., Messen, H.D. 1994. Expression of WT1 Wilms' tumor suppresor gene by normal and malignant melanocytes. *Int J Cancer*. 59: 78-82.
- Rupprecht, H.D., Drummond, I.A., Madden, S.L., Rauscher, F.J.III and Sukhatme, V.P. 1994. The Wilms'tumor suppressor gene WT1 is negatively regulated. *J Biol Chem*. 269: 6198-206.
- Ryan, G., Steele-Perkins, V., Morris, J.F., Rauscher, F.J. III and Dressler, G.R. 1995. Repression of Pax2 by WT1 during normal kidney development. *Development* 121: 867-75.
- Scharnhorst, V., Van der Eb, A.J. and Jochemsen, A.G. 2001. WT1 protein: function in growth and differentiation. *Gene* 273: 141-61.
- Schedl, A. and Hastie, N. 1998. Multiple roles for the Wilms'tumor suppressor gene, WT1 in genitourinary development. *Mol Cell Endocrinol*. 140: 65-9.

- Sells, S.F., Han, S.S., Muthukkumar, S., Maddiwar, N., Johnstone, R., Boghaert, E., Gillis, D., Liu, G., Nair, P., Monnig, S., Collini, P., Mattson, M.P., Sukhatme, V.P., Zimme, S.G., Wood, D.P. Jr., McRoberts, J.W., Shi, Y. and Rangnekar, V.M. 1997. Expression and function of the leucine zipper protein Par-4 in apoptosis. *Mol Cell Biol.* 17: 3823-32.
- Sharma, P.M., Bowman, M., Madden, S.L., Rauscher, G.J. and Sukumar, S. 1994. RNA editing in the Wilms' tumor susceptibility gene. *Genes Dev.* 8: 720-31.
- Shinagawa, T. and Ishii, S. 2003. Generation of Ski-knockdown mice by expressing a long double-strand RNA from an RNA polymerase II promoter. *Genes Dev.* 17: 1340-5.
- Smith, S.I., Down, M., Boyd, A.W. and Li, C.L. 2000. Expression of the Wilms' tumor suppressor gene, WT1, reduces the tumorigenicity of the leukemic cell line M1 in C.B-17 scid/scid mice. *Cancer Res.* 60: 808-14.
- Silberstein, G.B., Van Horn, K., Strickland, P., Robert, C.T. Jr. and Daniel, C.W. 1997. Altered expression of WT1 Wilms' tumor suppressor gene in human breast cancer. *Proc Natl Acad Sci USA.* 94: 8132-7.
- Sriplung, H., Sontipong, S., Martin, N., Wiangnon, S., Vootiprux, V., Cheirsilpa, A., Kanchanabat, C., and Khuhaprema, 2003. Cancer in Thailand, Vol III. Ministry of Public Health and Ministry of University Affair.
- Stein, P., Svoboda, P. and Schultz, R.M. 2003. Transgenic RNAi in mouse oocytes: a simple and fast approach to study gene function. *Dev. Biol.* 256: 187-93.
- Svoboda, P., Stein, P., Hayashi, H. and Schultz, R.M. 2000. Selective reduction of dormant maternal mRNAs in mouse oocytes by RNA interference. *Development* 127: 4147-56.
- Tabara, H., Sarkissian, M., Kelly, W.G., Fleenor, J., Grishok, A., Timmons, L., Fire, A. and Mello, C.C. 1999. The rde-1 gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* 99: 123-32.
- Timmons, L., Court, D.L. and Fire, A. 2001. Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene* 263: 103-13.
- Tuschl, T. 2001. RNA interference and small interfering RNAs. *Chembiochem* 2: 239-45.

- Van der Krol, A.R., Mur, L., Beld, M., Mol, J.N.M. and Stuitje, A.R. 1990. Flavonoid genes in petunia: Addition of a limited number of gene copies may lead to a suppression of gene expression. *Plant Cell.* 2: 291-9.
- Vance, V. and Vaucheret, H. 2001. RNA silencing in plants-Defense and counterdefense. *Science* 292: 2277-88.
- Voinnet, O., Lederer, C. and Baulcombe, D.C. 2000. A viral movement protein prevents spread of the gene silencing signal in *Nicotiana benthamiana*. *Cell* 103: 157-67.
- Wang, Z.Y., Madden, S.L., Deuel, T.F. and Rauscher, J.F.III. 1992. The Wilms' tumor suppressor gene product, WT1, repress transcription of the platelet-derived growth factor A chain gene. *J Biol Chem.* 267: 21999-2002.
- Wang, Z.Y., Qiu, Q.Q., Enger, K.T. and Deuel, T.F. 1993. A second transcriptionally active DNA-binding site for the Wilms tumor gene product, WT1. *Proc Natl Acad Sci USA.* 90: 8896-900.
- Wang, Z., Qiu, Q., Haung, J., Gurrieri, M. and Deuel, T.F. 1995. Products of alternative splice transcripts of the Wilms' tumor suppressor gene, WT1, have altered DNA binding specificity and regulate transcription in different ways. *Oncogene* 10: 415-22.
- Wei, Q., Lipardi, C. and Paterson, B.M. 2003. Analysis of the 3'-hydroxyl group in *Drosophila* siRNA function. *Methods* 30: 337-47.
- Werner, H., Roberts, C.T.Jr. and LeRoith, D. 1993. The regulation of IGF-I receptor gene expression by positive and negative zinc-finger transcription factors. *Adv Exp Med Biol.* 343: 91-103.
- WHO: *World Health Report 2003* Geneva: World Health Organization; 2003.
- Wooster, R., Neuhausen, S.L. mangion, J., Quirk, Y., Ford, D., Collins, N., Nguyen, K., Seal, S., Tran, T. and Averill, D. 1994. Localization of breast cancer susceptibility gene, BRCA2 to chromosome 13q12-13. *Science* 265: 2088-90.
- Yamagami, T., Sugiyama, H., Inoue, K., Ogawa, H., Tatekawa, T., Hirata, M., Kudoh, T., Akiyama, T., Murakami, A. and Maekawa, T. 1996. Growth inhibition of human leukemic cells by WT1 (Wilms' tumor gene) antisense oligodeoxynucleotides: implications for the involvement of WT1 in leukemogenesis. *Blood.* 87: 2878-84.

- Yang, D., Lu, H. and Erickson, J.W. 2000. Evidence that processed small dsRNAs may mediate sequence-specific mRNA degradation during RNAi in *Drosophila* embryos. *Curr. Biol.* 10: 1191-200.
- Yong, B. and Heath, J. 2000. Female reproductive system. In: Wheater functional histology : a text and color atlas, 4 th ed , Harcourt Publishers, Grafos, SA pp. 368-77.
- Zamore, P.D., Tuschl, T., Sharp, P.A. and Bartel, D.P. 2000. RNAi: double stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* 101: 25-33.
- Zapata-Benavides, P., Tuna, M., Lopez-Berestein, G. and Tari, A.M. 2002. Down regulation of Wilms' tumor 1 protein inhibits breast cancer proliferation. *Biochem Biophys Res Commun.* 295: 784-90.