

Contents

	Page
Contents	vii
List of Tables	ix
List of Figures	xii
Chapter	
1. Introduction	1
1.1 Chitin-Chitosan	2
1.2 Ethanol	8
1.3 Membrane process	9
1.4 Pervaporation process	10
1.5 Advantages of membrane process	12
1.6 Synthetic membranes	12
1.7 Review of Literatures	14
1.8 Research objectives	22
2. Theory	23
2.1 Pervaporation process	23
2.2 Response Surface Methodology	28
3. Experiment	31
3.1 Materials	31
3.2 Equipments	31
3.3 Method	32
3.3.1 Membrane Preparation	32
3.3.2 Membrane Characterization	34
3.3.3 Pervaporation Experiment	35

Contents (Continued)

	Page
4. Results and Discussion	39
4.1 Membrane Characterization	39
4.1.1 Swelling ratio	39
4.1.2 Tensile strength	43
4.1.3 Sorption selectivity	45
4.1.4 Pervaporation performance	47
4.2 Pervaporation studies	55
4.2.1 Feed concentration	55
4.2.2 Feed temperature	58
5. Conclusions	63
6. References	65
7. Appendixes	71
8. Vitae	115

List of Tables

Table	Page
1.1 Comparison of Mark-Houwink constants	5
1.2 Important membrane separation processes	11
1.3 Examples of Applications and Alternative Separation Processes	11
2.1 Coded variables of independent variables	29
3.1 Maximum and minimum values of each variable	32
3.2 Coded and Natural variables of each independent variable for membrane preparation	33
3.3 Coded and Natural variables of each independent variable for pervaporation study	38
4.1 Comparison of pervaporation dehydration of ethanol using various membranes	61
A1 Results from viscosity measurement of chitosan solution at 25 °C	72
C1 Refractive index of water-ethanol mixtures at 20 °C	77
D1 Swelling ratio of each experiment	80
D2 Sorption selectivity of each experiment	81
D3 The efficiency of chitosan membrane for dehydration of ethanol using pervaporation process (Flux)	82
D4 The efficiency of chitosan membrane for dehydration of ethanol using pervaporation process (Seperaton factor)	83
D5 Effect of feed concentrations (ethanol) and feed temperature on flux using pervaporation process	84
D6 Effect of feed concentrations (ethanol) and feed temperature on separation factor using pervaporation process	85
D7a, b and c Computer output from essential linear regression for fitting a model of swelling ratio	86
D8 Effect of crosslinking time and membrane formation temperature on swelling ratio	88

List of Tables (Continued)

Table	Page
D9 Effect of crosslinking time and concentration of sulfuric acid on swelling ratio	89
D10 Effect of membrane formation temperature and concentration of sulfuric acid on swelling ratio	90
D11a, b and c Computer output from essential linear regression for fitting a model of tensile strength	91
D12 Effect of membrane formation temperature and concentration of sulfuric acid on tensile strength	93
D13 Effect of membrane formation temperature and crosslinking time on tensile strength	94
D14 Effect of crosslinking time and concentration of sulfuric acid on tensile strength	95
D15a, b and c Computer output from essential linear regression for fitting a model of sorption selectivity	96
D16 Effect of crosslinking time and concentration of sulfuric acid on sorption selectivity	98
D17 Effect of crosslinking time and membrane formation temperature on sorption selectivity	99
D18 Effect of concentration of sulfuric acid and membrane formation temperature on sorption selectivity	100
D19a, b and c Computer output from essential linear regression for fitting a model of separation factor	101
D20 Effect of crosslinking time and concentration of sulfuric acid on separation factor	103
D21 Effect of crosslinking time and membrane formation temperature on separation factor	104

List of Tables (Continued)

Table	Page
D22 Effect of concentration of sulfuric acid and membrane formation temperature on separation factor	105
D23a, b and c Computer output from essential linear regression for fitting a model of flux	106
D24 Effect of crosslinking time and concentration of sulfuric acid on flux	108
D25 Effect of crosslinking time and membrane formation temperature on flux	109
D26 Effect of concentration of sulfuric acid and membrane formation temperature on flux	110
D27 a, b and c Computer output from essential linear regression for fitting a model of separation factor	111
D28 a, b and c Computer output from essential linear regression for fitting a model of flux	113

List of Figures

Figure	Page
1.1 Structural examples of chitin and chitosan	3
1.2 Principle of membrane process	10
1.3 Schematic drawing of porous and non-porous membranes	13
1.4 Schematic drawing of symmetric and asymmetric membrane cross-section	14
2.1 Schematic diagram of solution-diffusion model	24
3.1 Schematic diagram of sorption apparatus	35
3.2 Schematic diagram of cross-section of permeation cell	36
3.3 Schematic diagram of pervaporation setup	37
4.1 Chemical structures of ionic crosslinked chitosan membrane	40
4.2 Effect of concentration of sulfuric acid on swelling ratio at different crosslinking times and membrane formation temperatures	41
4.3 Effect of crosslinking time on swelling ratio at different concentrations of sulfuric acid and membrane formation temperatures	41
4.4 Effect of membrane formation temperature on swelling ratio at different concentrations of sulfuric acid and crosslinking times	42
4.5 Effect of concentration of sulfuric acid on tensile strength at different crosslinking times and membrane formation temperatures	43
4.6 Effect of crosslinking time on tensile strength at different concentrations of sulfuric acid and membrane formation temperatures	44
4.7 Effect of membrane formation temperature on tensile strength at different concentrations of sulfuric acid and crosslinking times	44
4.8 Effect of concentration of sulfuric acid on sorption selectivity of water at different crosslinking times and membrane formation temperatures	45

List of Figures (Continued)

Figure	Page
4.9 Effect of crosslinking time on sorption selectivity of water at different concentrations of sulfuric acid and membrane formation temperatures	46
4.10 Effect of membrane formation temperature on sorption selectivity of water at different concentrations of sulfuric acid and crosslinking times	46
4.11 Effect of concentration of sulfuric acid on total flux for dehydration of 87 %w/w ethanol at 60 °C through membrane prepared at different crosslinking times and membrane formation temperatures	47
4.12 Effect of concentration of sulfuric acid on separation factor for dehydration of 87 %w/w ethanol at 60 °C through membrane prepared at different crosslinking times and membrane formation temperatures	48
4.13 Effect of crosslinking time on total flux for dehydration of 87 %w/w ethanol at 60 °C through membrane prepared at different concentrations of sulfuric acid and membrane formation temperatures	49
4.14 Effect of crosslinking time on separation factor for dehydration of 87 %w/w ethanol at 60 °C through membrane prepared at different concentrations of sulfuric acid and membrane formation temperatures	49
4.15 Effect of membrane formation temperature on total flux for dehydration of 87 %w/w ethanol at 60 °C through membrane prepared at different concentrations of sulfuric acid and crosslinking times	50
4.16 Effect of membrane formation temperature on separation factor for dehydration of 87 %w/w ethanol at 60 °C through membrane prepared at different concentrations of sulfuric acid and crosslinking times	50

List of Figures (Continued)

Figure	Page
4.17 Effect of (a) concentration of sulfuric acid and crosslinking time, (b) crosslinking time and heating temperature, and (c) concentration of sulfuric acid and heating temperature, on separation factor for dehydration of 87 %w/w ethanol at 60 °C	52
4.18 Effect of (a) concentration of sulfuric acid and crosslinking time, (b) crosslinking time and heating temperature, and (c) concentration of sulfuric acid and heating temperature, on flux for dehydration of 87 %w/w ethanol at 60 °C	54
4.19 Effect of feed concentration on total flux for dehydration of ethanol at downstream pressure of 2.54 mm of Hg and different feed temperatures	55
4.20 Effect of feed concentration on separation factor for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed temperatures	56
4.21 Effect of feed concentration on ethanol flux for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed temperatures	56
4.22 Effect of feed concentration on water flux for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed temperatures	57
4.23 Effect of feed temperature on total flux for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed concentrations	58
4.24 Effect of feed temperature on separation factor for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed concentrations	58

List of Figures (Continued)

Figure	Page
4.25 Effect of feed temperature on ethanol flux for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed concentrations	59
4.26 Effect of feed temperature on water flux for dehydration of ethanol at downstream pressure 2.54 mm of Hg and different feed concentrations	59
4.27 Effect of feed concentration and feed temperature on separation factor for dehydration of ethanol at downstream pressure 2.54 mm of Hg	60
4.28 Effect of feed concentration and feed temperature on flux for dehydration of ethanol at downstream pressure 2.54 mm of Hg	61
A1 Relation between η_{sp}/C and concentration of chitosan solution	73
B1 IR spectrum of chitosan membrane	75
B2 Calibration curve of Degree of Deacetylation	76
C1 Refractive index of water-ethanol mixtures	79
D1 Effect of crosslinking time and membrane formation temperature on swelling ratio	88
D2 Effect of crosslinking time and concentration of sulfuric acid on swelling ratio	89
D3 Effect of membrane formation temperature and concentration of sulfuric acid on swelling ratio	90
D4 Effect of membrane formation temperature and concentration of sulfuric acid on tensile strength	93
D5 Effect of membrane formation temperature and crosslinking time on tensile strength	94

List of Figures (Continued)

Figure	Page
D6 Effect of crosslinking time and concentration of sulfuric acid on tensile strength	95
D7 Effect of crosslinking time and concentration of sulfuric acid on sorption selectivity	98
D8 Effect of crosslinking time and membrane formation temperature on sorption selectivity	99
D9 Effect of concentration of sulfuric acid and membrane formation temperature on sorption selectivity	100
D10 Effect of crosslinking time and concentration of sulfuric acid on separation factor	103
D11 Effect of crosslinking time and membrane formation temperature on separation factor	104
D12 Effect of concentration of sulfuric acid and membrane formation temperature on separation factor	105
D13 Effect of crosslinking time and concentration of sulfuric acid on flux	108
D14 Effect of crosslinking time and membrane formation temperature on flux	109
D15 Effect of concentration of sulfuric acid and membrane formation temperature on flux	110
D16 Effect of feed concentration and feed temperature on separation factor	112
D17 Effect of feed concentration and feed temperature on flux	114