

Contents

| | Page |
|--|-------------|
| บทคัดย่อ | (3) |
| Abstract | (5) |
| Acknowledgement | (7) |
| Contents | (8) |
| List of Tables | (11) |
| List of Figures | (12) |
| Chapter | |
| 1 Introduction | 1 |
| Introduction and review of literature | 1 |
| 1 General information of <i>Candida albicans</i> | 1 |
| 2 Pathogenicity of <i>C. albicans</i> | 3 |
| 2.1 Adherence of <i>Candida</i> | 3 |
| 2.2 Persorption | 4 |
| 2.3 Dimorphism and germ-tube formation | 4 |
| 2.4 High frequency switching | 5 |
| 2.5 Interference with phagocytosis | 6 |
| 2.6 Interference of immune defense | 6 |
| 2.7 Interference with complement | 7 |
| 2.8 Synergism with bacteria | 7 |
| 2.9 Enzymes of <i>Candida</i> | 7 |
| 2.10 Toxic substances produced by <i>C. albicans</i> | 8 |
| 2.11 Nitrosamines | 8 |
| 2.12 Acid metabolites of <i>Candida</i> | 8 |
| 3 Oral candidiasis and HIV infection | 9 |
| 3.1 Pathogenicity of <i>C. albicans</i> in HIV infection | 10 |
| 3.2 Clinical manifestations | 13 |

Contents (continued)

| | Page |
|---|-------------|
| 4 Diversity of <i>C. albicans</i> | 14 |
| 5 Techniques for typing of <i>C. albicans</i> | 15 |
| 5.1 Phenotyping | 15 |
| 5.2 Genotyping | 18 |
| 5.2.1 Restriction endonuclease analysis (REA) | 18 |
| 5.2.2 Randomly amplified polymorphic DNA (RAPD) | 19 |
| 5.2.3 Pulsed-field gel electrophoresis (PFGE) | 21 |
| Rationale for the study | 22 |
| Aims of the study | 24 |
| 2 Materials and Methods | 25 |
| 1 <i>C. albicans</i> strains | 25 |
| 2 DNA extraction | 25 |
| 3 DNA concentration measurement | 26 |
| 4 Primers | 26 |
| 5 PCR assays | 27 |
| 6 Gel electrophoresis and gel images | 27 |
| 7 Biotyping | 28 |
| 8 Antifungal susceptibility testing | 28 |
| 9 Genetic analysis | 29 |
| 10 Relationship analysis | 32 |
| 3 Results | 33 |
| Optimization of the PCR assays | 33 |
| Genetic analysis | 35 |
| 1 Clonal variability of <i>C. albicans</i> | 35 |
| 2 Genetic relatedness | 37 |

Contents (continued)

| | Page |
|---|-------------|
| 2.1 Genetic relatedness of <i>C. albicans</i> in HIV-infected individuals | 37 |
| 2.2 Genetic relatedness in HIV-uninfected subjects | 41 |
| Phenotypic analysis | 42 |
| 1 Biotyping | 42 |
| 2 Ketoconazole susceptibility test | 44 |
| Relationship between genotype and biotype, antifungal susceptibility | 45 |
| 1 Genetic relatedness of <i>C. albicans</i> for the assessment of genotype-phenotype relationship | 45 |
| 2 Genotype and biotype | 46 |
| 3 Genotype and ketoconazole susceptibility | 46 |
| 4 Statistical analysis of assessment of association between <i>C. albicans</i> genotype, biotype, and ketoconazole susceptibility | 49 |
| 4 Discussion | 51 |
| 5 Conclusion | 59 |
| Bibliography | 60 |
| Appendix | 74 |
| Vitae | 83 |

List of Tables

| Table | Page |
|---|-------------|
| 1 Potential virulent factors of <i>Candida</i> species | 4 |
| 2 Random primers tested in pilot studies for RAPD analysis of <i>C. albicans</i> isolates | 27 |
| 3 The summary of the concentrations of 7 primers, MgCl ₂ and the temperature profiles used in PCR analyses | 30 |
| 4 Isotypes from 189 strains situated in the clusters shown in the dendrogram | 40 |
| 5 Details of <i>C. albicans</i> strains tightly related at S _{ABS} of 0.80 or greater | 41 |
| 6 The API biotype profiles and their percentage distribution of 106 <i>C. albicans</i> strains from HIV-infected patients | 43 |
| 7 Identical biotypes of <i>C. albicans</i> , but with varying RAPD genotypes, found in the same patient | 44 |
| 8 Details of ketoconazole susceptibility testing in 94 strains | 45 |
| 9 Statistical analysis by Chi-square test | 49 |
| 10 Odds ratio analysis in biotype and ketoconazole susceptibility of 94 <i>C. albicans</i> strains isolated from HIV-infected individuals | 50 |
| 11 Data on the source of <i>C. albicans</i> evaluated and respective number of clones and clusters as per the source of collection | 74 |
| 12 Patient code details and the number of <i>C. albicans</i> strain(s) isolated from HIV-free subjects | 75 |
| 13 Details of clusters with genotypes, biotypes and ketoconazole susceptibility test of <i>C. albicans</i> strains studied | 78 |

List of Figures

| Figure | Page |
|--|------|
| 1 Schematic diagram illustrating the possible theoretical considerations which may contribute to the selection of <i>Candida</i> strains | 12 |
| 2 Diagram demonstrating restriction endonuclease analysis (REA) procedure | 20 |
| 3 Diagram demonstrating randomly amplified polymorphic DNA (RAPD) procedure | 21 |
| 4 Optimization of PCR assays : primer 1 | 34 |
| 5 Optimization of PCR assays : primer PA03 | 34 |
| 6 Optimization of PCR assays : primer RSD 11, RSD12 | 35 |
| 7 Different genotypes of <i>C. albicans</i> derived from twelve HIV-infected individuals | 36 |
| 8 Dendrogram of <i>C. albicans</i> from HIV-infected individuals | 39 |
| 9 Dendrogram of <i>C. albicans</i> from non-HIV subjects | 40 |
| 10 The details of genotyping by dendrogram, biotyping and MICs | 47 |
| 11 Diagram showing genotype and biotype relationship | 48 |
| 12 Diagram showing genotype and ketoconazole susceptibility Relationship | 48 |
| 13 Dendrogram of 112 <i>C. albicans</i> subsequently compared with phenotypes | 77 |