

CONTENTS

	PAGE
บทคัดย่อ	(3)
ABSTRACT	(5)
ACKNOWLEDGEMENT	(7)
CONTENTS	(8)
LIST OF TABLES	(11)
LIST OF FIGURES	(12)
CHAPTER	
1 INTRODUCTION	1
1.1 Background and Rationale	1
1.2 Objectives of the Research	3
1.3 Hypothesis	4
1.4 Scope and Limitation of the Research	4
1.5 Expecting Outcomes	5
2 LITERATURE REVIEW	6
2.1 n-3 Polyunsaturated Fatty Acids (n-3 PUFAs)	6
2.1.1 Source of n-3 PUFAs	6
2.1.2 Health Benefits of n-3 PUFAs	8
2.1.3 Intake of n-3 PUFAs	10
2.2 Transesterification	11
2.2.1 Distinction of Enzymatic and Chemical Transesterification	12
2.2.2 Chemistry of Chemical Transesterification	13
2.2.3 Mechanism of Chemical Transesterification	13
2.2.4 Applications of Chemical Transesterification in Food Lipids	15
2.3 Microencapsulation	16
2.3.1 Corn Syrup Solids as Encapsulant	17
2.3.2 Emulsification	19
2.3.3 Spray Drying	23
2.4 Changes in physicochemical properties of spray dried powder	25
2.4.1 Structural Collapse and Crystallization	25
2.4.2 Maillard Reaction	27
	(8)

CONTENTS (Continued)

	PAGE
2.5 Lipid Oxidation	30
2.5.1 Lipid Oxidation in Liquid Emulsion	30
2.5.2 Lipid Oxidation in Dried Emulsion	33
3 RESEARCH BRIEFING	36
4 CHEMICAL TRANSESTERIFICATION OF TUNA OIL TO ENRICH n-3 POLYUNSATURATED FATTY ACIDS	38
4.1 Abstract	38
4.2 Introduction	38
4.3 Materials and Methods	39
4.4 Results and Discussion	42
4.5 Conclusion	51
4.6 Concision of Further Study	51
5 EMULSIFICATION OF TUNA OIL IN TWO-LAYERED INTERFACIAL MEMBRANES	52
5.1 Abstract	52
5.2 Introduction	52
5.3 Materials and Methods	54
5.4 Results and Discussion	58
5.5 Conclusion	71
5.6 Concision of Further Study	71
6 INCREASING THE OXIDATIVE STABILITY OF LIQUID AND DRIED TUNA OIL-IN WATER EMULSIONS	72
6.1 Abstract	72
6.2 Introduction	72
6.3 Materials and Methods	74
6.4 Results and Discussion	77
6.5 Conclusion	86
6.6 Concision of Further Study	86

CONTENTS (Continued)

	PAGE
7 CHARACTERIZATION OF SPRAY-DRIED TUNA OIL EMULSIFIED IN TWO- LAYERED INTERFACIAL MEMBRANES	87
7.1 Abstract	87
7.2 Introduction	87
7.3 Materials and Methods	88
7.4 Results and Discussion	93
7.5 Conclusion	102
7.6 Concision of Further Study	102
8 EFFECT OF STORAGE ENVIRONMENTS ON STABILITY OF SPRAY-DRIED TUNA OIL EMULSION	103
8.1 Abstract	103
8.2 Introduction	103
8.3 Materials and Methods	105
8.4 Results and Discussion	109
8.5 Conclusion	121
9 RESEARCH SUMMARY	122
9.1 Summary of Research Investigation	122
9.2 Suggestion for Future Research	125
REFERENCES	127
VITAE	144

LIST OF TABLES

TABLE	PAGE
1	Amount of EPA and DHA in fish, crustacean and fish oils. 7
2	Reaction conditions for transesterification of sodium methoxide (NaOCH ₃). 14
3	Glass transition temperature of maltodextrins/corn syrup solids as a function of water activity. 27
4	Lipid composition (% peak area) of refined tuna oil, n-3 fatty acids methyl ester (n-3 FAME) and transesterified tuna oil. 44
5	Fatty acid composition (mg/g oil) in refined tuna oil, n-3 fatty acids methyl ester (n-3 FAME) and transesterified tuna oil. 46
6	Linear regression coefficient of full factorial design-effects for response variables (TG recovery, EPA and DHA) on transesterification of tuna oil. 48
7	Influence of corn syrup solids concentration on electrical charge (ζ -potential), apparent viscosity (η_{APP}) and relative viscosity (η_{REL}) of secondary emulsions consisting of 5 wt% tuna oil, 1 wt% lecithin, 0.2 wt% chitosan, 100 mM acetate buffer (pH 3.0). 58
8	Influence of NaCl (0 or 200 mM) and corn syrup solids, CSS (0 or 20 wt%) on mean particle diameter and electrical charge (ζ -potential) of emulsions consisting of 5 wt% tuna oil, 1 wt% lecithin, 100 mM acetate buffer (pH 3.0), and either 0 wt% chitosan (primary emulsion) or 0.2 wt% chitosan (secondary emulsion). 61
9	Dependence of mean particle diameter (μm) of primary and secondary emulsions on number of freeze-thaw cycles (-18°C for 22 h/ 30°C for 2 h) in the absence and presence of 20 wt% corn syrup solids (CSS). 67
10	Effect of inlet temperature on properties of spray dried tuna oil emulsion. 95
11	Effect of inlet temperature on color of spray dried tuna oil emulsion. 98

LIST OF FIGURES

FIGURE		PAGE
1	Proposed reaction mechanisms for chemical transesterification: a), carbonyl addition and b), Claisen condensation.	15
2	Instability of emulsion.	19
3	Molecular structural of major phospholipids; R = fatty acid residue.	20
4	A schematic picture of the structure and the conformation of cationic starch/ anionic surfactant complexes in dilute aqueous solutions.	21
5	Structures of chitin and chitosan.	23
6	Spray drying system.	24
7	Schematic presentation of a droplet during drying in the crust formation period; Q = heat, b = crust thickness, r-b = radius of the liquid droplet, dm/dt = mass transfer.	25
8	Interactions between Maillard reaction and lipid oxidation pathways in nonenzymatic browning development.	29
9	Generalized scheme for autoxidation of lipids.	31
10	Interfacial phenomena to explain the action of antioxidants in bulk oil and oil-in-water emulsion systems.	33
11	Relative rate of reaction as a function of water activity: a), nonenzymatic browning. b), lipid oxidation.	34
12	Effect of reaction time on remaining of TG. Transesterification was carried out at 80°C for 24 h, the mole ratio of tuna oil to n-3 FAME was 1:4 and NaOCH ₃ amount was 1.0% wt% of reactants.	43
13	Effect of reaction time on incorporation of EPA and DHA into tuna oil with transesterification. Transesterification was carried out at 80°C for 24 h, the mole ratio of tuna oil to n-3 FAME was 1:4 and NaOCH ₃ amount was 1.0% wt% of reactants.	45
14	Effect of reaction conditions on TG remaining of tuna oil after transesterification with the mole ratios of tuna oil to n-3 FAME were 1:2 (a), 1:3 (b) and 1:4(c) at 60, 70 and 80°C.	47

LIST OF FIGURES (Continued)

FIGURE		PAGE
15	Effect of reaction condition on incorporation of EPA into TG of tuna oil after transesterification. The mole ratios of tuna oil to n-3 FAME were 1:2 (a), 1:3 (b) and 1:4(c) at 60, 70 and 80°C.	49
16	Effect of reaction condition on incorporation of DHA into TG of tuna oil after transesterification. The mole ratios of tuna oil to n-3 FAME were 1:2 (a), 1:3 (b) and 1:4(c) at 60, 70 and 80°C.	50
17	Influence of pH on electrical charge (ζ -potential) of emulsion droplets in primary (5 wt% tuna oil, 1 wt% lecithin, 100 mM acetate buffer, pH 3.0) and secondary emulsions (5 wt% tuna oil, 1 wt% lecithin, 0.2 wt% chitosan, 100 mM acetate, pH 3.0) in the absence and presence of 20 wt% corn syrup solids (CSS).	62
18	Influence of pH on mean diameter of emulsion droplets in primary (5 wt% tuna oil, 1 wt% lecithin, 100 mM acetate, pH 3.0) and secondary emulsions (5 wt% tuna oil, 1 wt% lecithin, 0.2 wt% chitosan, 100 mM acetate, pH 3.0) in the absence and presence of 20 wt% corn syrup solids (CSS).	63
19	Dependence of microstructure on isothermal treatment temperature (30 and 90°C for 30 min) for: a) primary emulsion; b) primary emulsion in the presence of 20 wt% corn syrup solids; and c) secondary emulsions.	66
20	Microstructure of reconstituted emulsions for: a) primary emulsion; b) and c) for secondary emulsions in the absence and presence of 20 wt% corn syrup solids, respectively.	69
21	Time dependence of the mean particle diameter and obscuration of powdered secondary emulsions containing 20 wt% corn syrup solids that were dispersed into an aqueous solution in the measurement cell of a light scattering instrument.	70