

## REFERENCES

- เยาวดี คุปตะพันธ์, 2546, สารเร่งเนื้อแดง: ปัจจัยของคนบริโภคหมู, วารสารอาหาร, สถาบันกั้นกว่า  
และพัฒนาผลิตภัณฑ์อาหาร, มหาวิทยาลัยเกษตรศาสตร์, 33(2), เมษายน-มิถุนายน 2546,  
หน้า103-104.
- วีໄล ตระกูลไօสດ, รุ่งรัตน์ ปิยันนท์รัศศรี, วัฒนาพร แพทบี้ประสีทธิ์, วงศธร งานเอนก, จันคนา  
บุรณะไօสດ และ ดาวลักษ์ ศรีพงษ์, 2544, การวิเคราะห์หาปริมาณชา酇ูทามอลที่ตกค้างใน  
เนื้อสุกร ด้วยวิธีไฮเพอร์ฟอร์แมนซ์ลิกวิดไฮดรมาโทกราฟี : การศึกษาเบื้องต้น, วารสาร  
มหาวิทยาลัยศิลปากร, 21-22(1) หน้า 235-247.
- กระทรวงสาธารณสุข, ประกาศกระทรวงสาธารณสุข (ฉบับที่ 269) พ.ศ. 2546 เรื่อง มาตรฐาน  
อาหารที่มีการปนเปื้อนสารเคมีอุ่นเบื้องต้น โภณิสต์.
- สำนักงานคณะกรรมการอาหารและยา, กระทรวงสาธารณสุข, ประกาศสำนักงานคณะกรรมการ  
อาหารและยา เรื่องหลักเกณฑ์ เงื่อนไข และวิธีการตรวจวิเคราะห์การปนเปื้อนสารเคมีบาง  
ชนิดในอาหาร, 2546.
- Aerts, M.M.I., Hogenboom, A.C. and Brinkman, U.S.Th. 1995. Analytical strategies  
for the screening of veterinary drugs and their residues in edible products.  
*Journal of Chromatography B:Biomedical Applications*, 667: 1-40.
- Ardrey, R.E. 2003. Liquid chromatography mass spectrometry: An introduction.  
London: John Wiley & Sons, pp 6-29.
- Barker, S.A., Long, A.R. and Short, C.R. 1989. Isolation of drug residues by solid  
phase dispersion. *Journal of chromatography*, 475: 353-361.
- Barker<sup>a</sup>, S.A. 2000. Applications of matrix solid-phase dispersion in food analysis.  
*Journal of chromatography A*, 880: 63-68.

- Barker<sup>b</sup>, S. A. 2000. Matrix solid-phase dispersion: A review. *Journal of chromatography A*, **885**: 115-127.
- Barnett, N. W., Hindson, B.J. and Lewis, S.W. 1999. Determination of ranitidine and salbutamol by flow injection analysis chemiluminescence detection. *Analytica Chimica Acta*, **384**: 151-158.
- Bergés, R., Segura, F., Torra, X. and Ventura, R. 1999. Analytical methodology for enantiomers of salbutamol in human urine for application in doping control. *Journal of Chromatography B*, **723**: 173-184.
- Bilbao-Garay, J., Hoyo-Jimenez, J.F., Lopez-Jimenez ,M., Viruela-Sebastian, M., Perianes-Matesanz, J., Munoz-Moreno, P. and Ruiz-Galiana, J. 1997. Clenbuterol poisoning. Clinical and analytical data on an outbreak in Mostoles, Madrid. *Revista Clinica Espanola*, **197**:92-25.
- Bland, R.E., Tanner, R.J.N., Chern, W.H., Lang, J.R. and Powell, J.R. 1990. Determination of albuterol concentrations in human plasma using solid-phase extraction and high-performance liquid chromatography with fluorescence detection. *Journal of Pharmaceutical & Biomedical Analysis*, **8**: 591-596.
- Boulton, D.W. and Fawcett, J.P. 1995. Determination of salbutamol enantiomers on human plasma and urine by chiral high-performance liquid chromatography. *Journal of Chromatography B: Biomedical Application*, **672**: 103-109.
- Boyd, D. and O'Keeffe, M. 1994. Matrix solid-phase dispersion as a multiresidue extraction technique for  $\beta$ -agonists in bovine liver tissue. *The Analyst*, **119**: 1467-1470.
- Boyd, D., O'Keeffe, M. and Smyth, S.M. 1995. Matrix solid-phase dispersion linked to solid phase extraction for  $\beta$ -agonists in liver samples: An update. *Analytical Proceedings Including Analytical Communications*, **32**: 301-303.

- Boyd, D., O'Keeffe, M. and Smyth, M. R. 1996. Methods for the determination of ( $\beta$ -agonists in biological matrices: A review. *The Analyst*, 121: 1R-10R.
- Brambilla, G., Loizzo, A., Fontana, L., Strozzi, M., Guarino, A. and Soprano, V. 1997. Food poisoning following consumption of clenbuterol-treated veal in Italy. *Journal of the American Medical Association*, 278:635.
- Phenomenex 2005. Catalogue 2005/2006.
- Chan, T.Y.K. 1998. Food poisoning following consumption of clenbuterol-treated livestock. *Hong Kong Practitioner*, 20: 366-369.
- Chen, Y., Wang, W., Duan, J., Chen, H. and Chen, G. 2005. Separation and determination of clenbuterol, cimaterol and salbutamol by capillary electrophoresis with amperometric detection. *Electroanalysis*, 17: 706-712.
- Chin-En, T. and Fusao, K. 1994. Liquid chromatographic determination of salbutamol and clenbuterol residues in swine serum and muscle. *Microbios*, 80: 251-8.
- Couper, F.J. and Drummer, O.H. 1996. Gas chromatography-mass spectrometry determination of  $\beta_2$ -agonists in postmortem blood: application in forensic medicine. *Journal of Chromatography B:Biomedical Applications*, 685: 265-272.
- Crouch, M.D. and Baker, S.A. 1997. Analysis of toxic waste in tissues from aquatic species: Applications of matrix solid-phase dispersion (Review). *Journal of Chromatography A*, 774: 287-309.
- Damasceno, L., Ventura, R., Ortúñoz, J. and Segura, J. 2000. Derivatization procedures for the detection of  $\beta_2$ -agonists by gas chromatographic/ mass spectrometric analysis. *Journal of Mass Spectrometry*, 35: 1285-1294.

Department of Medical Science, Ministry of Public Health 1998. *Annual Report of Department of Medical Sciences 1998*, p 55.

Department of Medical Science, Ministry of Public Health 2004. *Annual Report of Department of Medical Sciences 2004*, pp 15-17.

Dol, I. and Knochen, M. 2004. Flow-injection spectrophotometric determination of salbutamol with 4-aminoantipirine. *Talanta*, 64: 1233-1236.

Dunshea, F.R., D'Souza, D.N., Pethick, D.W., Harper G.S. and Warner, R.D. 2005. Effects of dietary factors and other metabolic modifiers on quality and nutritional value of meat (Review), *Meat Science*, 71: 8-38.

Council Directive 96/22/EC of 29 April 1996 concerning the prohibition on the use in stockfarming of certain substances having a hormonal or thyrostatic action and of  $\beta$ -agonists, and repealing Directives 81/602/EEC, 88/146/EEC and 88/299/EC, *Official Journal of the European Communities*, L125, pp 3-9.

Commission Decision 2002/657/EC of 12 August 2002 implementing Council Directive 96/23/EC concerning the performance of analytical methods and the interpretation of results, *Official Journal of the European Communities*, L221, pp 8-36.

Emm, T., Lesko, L.J. and Leslie, J. 1988. Determination of albuterol in human serum by reversed-phase high-performance liquid chromatography with electrochemical detection. *Journal of Chromatography*, 277: 311-317.

Esquisabel, A., Hernández, R.M., Gascón, A.R., Igartua, M., Calvo, B. and Pedez, J.L. 1997. Determination of salbutamol enantiomers by high-performance capillary electrophoresis and its application to dissolution assays. *Journal of Pharmaceutical and Biomedical Analysis*, 16: 357-366.

EURACHEM Guide. 1998. The Fitness for Purpose of Analytical Method: A Laboratory Guide to Method Validation and Related Topics (1<sup>st</sup> ed), 61 pp.

Euro-diagnostica,  $\beta$ -agonist EIA. 2003. A microtiter plate based competitive enzyme immunoassay for analysis and screening of urine, faeces, feed, bile, tissue, plasma, hair and choroid/retina samples samples, 5061BAG1p[17] 05.03.

Felix, F.S., Quintino, M. S.M., Carvalho, A.Z., Coelho, L. H.G., Lago, C.L. and Angnes, L. 2006. Determination of salbutamol in syrups by capillary electrophoresis with contactless conductivity detection (CE-C<sup>4</sup>D). *Journal of Pharmaceutical and Biomedical Analysis*, **40**: 1288-1292.

Food and Drug Administration, Ministry of Public Health 2002. *Annual Report of Food and Drug Administration 2002*, p 17.

Food and Drug Administration, Ministry of Public Health 2003. *Annual Report of Food and Drug Administration 2003*, p 45.

Hamilton, R.J. and Swell, P.A. 1982. Introduction to high performance liquid chromatography (2<sup>nd</sup> ed.), London: Chapman and Hall, 248 pp.

Harris, D.C. 1995. Quantitative chemical analysis (4<sup>th</sup> ed.), New York: W.H. Freeman and Company, pp 139-142.

Hernández-Carrasquilla, M. 2000. Gas chromatography-mass spectrometry analysis of  $\beta_2$ -agonists in bovine retina. *Analytica Chimica Acta*, **408**: 285-290.

Huclová, J., Šatinský, D., Sklenářová, H. and Karlíček, R. 2003. Determination of salbutamol using on-line solid phase extraction and sequential injection analysis. Comparison of chemiluminescence and fluorescence detection. *Analytical and Bioanalytical Chemistry*, **376**: 448-454.

- Hutchings, M.J. 1983. Determination of salbutamol in plasma by high-performance liquid chromatography with fluorescence detection. *Journal of Chromatography*, 277: 423-426.
- Jacobson, G.A., Chong, F.V. and Davies, N.W. 2003. LC-MS method for the determination of albuterol enantiomers in human plasma using manual solid-phase extraction and a non-deuterated internal standard. *Journal of Pharmaceutica and Biomedical Analysis*, 31: 1237-1243.
- Kaewklapanyachareon, L. 2001. Determination of salbutamol residue in pork tissue, pork liver and kidney by HPLC. *Bulletin of the Department of Medical Science*, 3: 215-221.
- Kataoka, H. 2003. New trends in sample preparation for clinical and pharmaceutical analysis. *Trends in Analytical Chemistry*, 22: 232-244.
- Koh, Y.M., Saleh, M.I. and Tan, S.C. 2003. Selective extraction of salbutamol from human plasma with the use of phenylboronic acid. *Journal of Chromatography B*, 987: 257-267.
- Koole, A., Bosman, J., Franke, J.P. and Zeeuw, R.A. 1999. Multiresidue analysis of  $\beta_2$ -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection. *Journal of Chromatography B*, 726: 149-156.
- Loss II, J.R., Orzechowski, R.F. and Hock, R.S. 2000. Measurement of albuterol in guinea pig serum by high performance liquid chromatography with fluorescence detection. *Biomedical Chromatography*, 14: 1-5.
- Maistro, S., Chiesa, E., Angeletti, R. and Brambilla, G. 1995. Beta blockers to prevent clenbuterol poisoning. *Lancet*, 346: 180.

- Malkki-Laine, L. and Hartikainen, E. 1996. Electrokinetic behaviour of salbutamol and its decomposition products and determination of salbutamol by micellar electrokinetic capillary chromatography. *Journal of Chromatography A*, 724: 297-306.
- Martínez-Navarro, J. F. 1990. Food poisoning related to consumption of illicit beta-agonist in liver. *Lancet*, 336: 1311.
- Miller, J.N. and Miller J.C. 2000. Statistics and Chemometrics for Analytical Chemistry (4<sup>th</sup> ed.), London: Dorset Press, 271 pp.
- Miller, L.G. and Greenblatt, D.J. 1986. Determination of albuterol in human plasma by high-performance liquid chromatography with fluorescence detection. *Journal of Chromatography*, 381: 205-208.
- Mills III, T. and Roberson, J.C. 1987. Instrument Data for Drug Analysis (2<sup>nd</sup> ed.), Vol. 2 and Vol.3, New York: Elsevier Science Publisher, pp 172-173 and pp 2058-2059.
- Mitra, S., 2003, Sample Preparation Techniques in Analytical Chemistry, Vol.162, New Jersey: John Wiley & Sons, pp 25-35.
- Moffat, A.C., Jackson, J.V., Moss, M.S. and Widdop, B. 1986. Clarke's Isolation and Identifications of drugs in Pharmaceuticals, body fluids, and postmortem (2<sup>nd</sup> ed.), London: The Pharmaceutical Press, pp 370-371 and 963-964.
- Molony, A., Allen, P., Joseph, R. and Tarrant, V. 1991. Influence of Beta-Adrenergic Agonists and Similar Compounds on Growth. in Growth Regulation in Farm Animals Advances in Meat Research, Vol.7, London: Elsevier Applied Science, pp 455-513.
- Neue, U.D. 1997. HPLC Columns. Theory, Technology, and Practice, New York: Wiley-VCH, 393 pp.

O'Keeffe, M.J., O'Keeffe, M. and Glennon, J.D. 1999. Supercritical fluid extraction (SFE) as a multi-residue extraction procedure for  $\beta$ -agonists in bovine liver tissue. *The Analyst*, **124**: 1355-1360.

O'Neil, M.J., Smith, A., Heckelman, P.E., Obenchain Jr., J.R., Gallipeau, J.A.R., D'Arecca, M.A. and Budavaris, S. 2001. The MERCK Index, An Encyclopedia of Chemicals, Drugs, and Biologicals (13<sup>th</sup> ed.), MERCK Research Laboratories Division of MERCK & Co., Inc., pp 42 and 167.

Ouyang, J., Duan, J.L., Baeyens, W.R.G. and Delanghe, J.R. 2005. A simple method for the study of salbutamol pharmacokinetics by ion chromatography with direct conductivity detection. *Talanta*, **65**: 1-6.

Parriott, D. 1993, A Practical Guide to HPLC Detection, California: Academic Press, pp 111-143.

Plum, D.C. 1999. Veterinary drug handbook (3<sup>rd</sup> ed.), New York, Pharma Vet Publishing, pp 14-15.

Polettini, A. 1996, Bioanalysis of  $\beta_2$ -agonists by hyphenated chromatographic and mass spectrometric techniques (Review). *Journal of Chromatography B*, **687**: 27-42.

Poole, C.F. and Schuette S.A. 1984. Contemporary Practice of Chromatography, Amsterdam: Elsevier Science Publisher, 708 pp.

R Development Core Team 2006. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.

- Ramos, F., Santos, C., Silva, A. and Silveira, M.I.N. 1998.  $\beta_2$ -Adrenergic agonists residues: simultaneous methyl- and butylboronic derivatization for confirmatory analysis by gas chromatography-mass spectrometry. *Journal of Chromatography B*, **716**: 366-370.
- Ramos, F., Banobre, M.C., Castilho, M.d.C and Silverra, M.I.N. 1999. Solid phase extraction (SPE) for multiresidue analysis of  $\beta_2$ -agonists in bovine urine. *Journal of Liquid Chromatography and Related Technologies*, **22**: 2307-2320.
- Ramos, F.J. 2000.  $\beta_2$ -Agonist extraction procedure for chromatographic analysis (Review). *Journal of Chromatography A*, **880**: 69-83.
- Reig, M., Batlle, N., Navarro, J.L. and Toldra, F. 2005. Stability of  $\beta$ -agonist methyl boronic derivatives before gas chromatography-mass spectrometry analysis. *Analytica Chimica Acta*, **529**: 293-297.
- Rouessac, F. and Rouessac, A. 2000. Chemical Analysis: Modern Instrumental methods and Techniques, London: John Wiley & Sons, 445 pp.
- Sagar, K.A., Hua, C., Kelly, M.T. and Smyth, M.R. 1992. Analysis of salbutamol in human plasma by high-performance liquid chromatography with electrochemical detection using a micro-electrochemical flow cell. *Electroanalysis*, **4**: 481-486.
- Saleh, M.I., Koh, Y.M., Tan, S.C. and Aishah, A.L. 2000. Clean-up detection and determination of salbutamol in human urine and serum. *The Analyst*, **125**: 1569-1572.
- Salleras, L., Domínguez, A., Mata, E., Taberner, J. L., Moro I. and Salvà P. 1995. Epidemiologic study of an outbreak of clenbuterol poisoning in Catalonia, Spain. *Public Health Reports*, **110**: 338-342.

Šatinský, D., Karlíček, R. and Svoboda, A. 2002. Using on-line solid phase extraction for flow-injection spectrophotometric determination of salbutamol. *Analytica Chimica Acta*, **455**: 103-109.

Schmeer, K., Sauter, T. and Schmid, J. 1997. Rapid pharmacokinetic screening of salbutamol in plasma samples by column-switching high-performance liquid chromatography-electrospray mass spectrometry. *Journal of Chromatography A*, **777**: 67-72.

Shepherd and M.J. 1991. Analysis of Veterinary Drug Residues in Edible Animal Products, in: Creaser, C.S. and Purchase, V., Food Contaminants - Sources and Surveillance, Cambridge, The Royal Society of Chemistry, pp 109-176.

Shishani, E., Chai, S. C., Jamokha, S., Aznar, G. and Hoffman, M. K. 2003. Determination of ractopamine in animal tissues by liquid chromatography-fluorescence and liquid chromatography/tandem mass spectrometry. *Analytica Chimica Acta*, **483**: 137-145.

Simpson, N.J.K. 2000. Solid-phase extraction: Principles, Techniques, and Applications, New York: Marcel Dekker, 514 pp.

Snyder, L.R. and Kirkland, J.J. 1979. Introduction to Modern Liquid Chromatography (2<sup>nd</sup> ed.), New York: John Wiley & Sons, 863 pp.

Snyder, L.R., Kirkland, J.J. and Glajch, J.L. 1997. Practical HPLC method development (2<sup>nd</sup> ed.), New York: John Wiley & Sons, 765 pp.

Strata X user's Guide, Phenomenex USA, p 6.

Tan, Y.K. and Soldin, S.J. 1984. Determination of salbutamol in human serum by reversed-phase high-performance liquid chromatography with amperometric detection. *Journal of Chromatography*, **277**: 311-317.

The World Anti-doping Agency 2004. The World Anti-Doping Code, the 2004 prohibited list, 18 pp.

U.S. Department of Health and Human Services, Food and Drug Administration, Center for Veterinary Medicine (CVM) 1999. Guidance for Industry, Validation of analytical procedures: Definition and terminology, Final guidance, 8 pp.

U.S. Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Center for Veterinary Medicine (CVM) 2001. Guidance for Industry, Bioanalytical Method Validation, 22 pp.

Ventura, R., Damasceno, L., Farré, M., Cardoso, J. and Segura, J. 2000. Analytical methodology for the detection of  $\beta$ -agonists in urine by gas chromatography-mass spectrometry for application in doping control. *Analytica Chimica Acta*, 418: 79-92.

Vlis, E., Mazereeuw, M., Tjaden, U.R., Irth, H. and Greef, J. 1995. Combined liquid-liquid electroextraction-isotachophoresis for loadability enhancement in capillary zone electrophoresis-mass spectrometry. *Journal of Chromatography A*, 712: 227-234.

Vyncht, G.V., Preece, S., Gaspar, P., Maghuin-Rogister, G. and DePauw, E. 1996. Gas and liquid chromatography coupled to tandem mass spectrometry for the multiresidue analysis of  $\beta$ -agonists in biological matrices. *Journal of Chromatography A*, 750: 43-49.

Waters Corporation 2003. Symmetry<sup>®</sup>: Applications notebook. p 19.

- Zhang, J., Xu, Y., Di, X. and Wu, M. 2006. Quantitation of salbutamol in human urine by liquid chromatography-electrospray ionization mass spectrometry. *Journal of Chromatography B*, **831**: 328-332.
- Zhang, X.Z., Gan, Y.R. and Zhao, F.N. 2004. Determination of salbutamol in human plasma and urine by high-performance liquid chromatography with a coulometric electrode array system. *Journal of Chromatographic Science*, **42**:263-267.
- Zhou, T., Hu, Q., Yu, H. and Fang, Y. 2001. Separation and determination of ( $\beta$ -agonists in serum by capillary zone electrophoresis with amperometric detection. *Analytica Chimica Acta*, **441**: 23-28.