

# CONTENTS

	<b>Page</b>
CONTENTS	viii
LIST OF TABLES	xi
LIST OF FIGURES	xiii
CHAPTER	
1. <a href="#">INTRODUCTION AND LITERATURE REVIEW</a>	1
<a href="#">1.1 Overview</a>	1
<a href="#">1.2 Arsenic contaminated in the environment</a>	1
<a href="#">1.2.1 Arsenic in soil</a>	3
<a href="#">1.2.2 Arsenic in plants</a>	4
<a href="#">1.2.3 Arsenic in water</a>	5
<a href="#">1.3 The analytical techniques for arsenic analysis</a>	5
<a href="#">1.4 Health Effects of arsenic on human</a>	8
<a href="#">1.5 Source and value of exposure</a>	9
<a href="#">1.6 Arsenic contamination in Ronphibun Sub-district</a>	10
<a href="#">1.6.1 Background information of arsenic contamination</a>	10
<a href="#">1.6.2 Distribution of arsenic in Ronphibun Sub-district</a>	12
<a href="#">1.6.3 Risk assessment studies</a>	13
<a href="#">1.7 Objectives</a>	14
<a href="#">1.8 Anticipation outcome</a>	14
2. <a href="#">METHODOLOGY</a>	15
<a href="#">2.1 Chemical and material</a>	15
<a href="#">2.2 Instruments</a>	15
<a href="#">2.3 Hydride Generation Atomic Absorption Spectrophotometry<sup>17</sup> (HGAAS)</a>	
<a href="#">2.4 Optimization of the analytical method</a>	18
<a href="#">2.4.1 Carrier gas (Ar) flow rate</a>	18

## CONTENTS (CONTINUED)

	<b>Page</b>	
2.4.2	<a href="#">Effect of NaBH<sub>4</sub> concentration</a>	18
2.4.3	<a href="#">Effect of HCl concentration</a>	18
2.4.4	Effect of KI / Ascorbic acid concentration	18
2.4.5	Effect of reduction time	19
2.4.6	Optimum of atomization temperature	19
2.5	Sampling site and sample preparation	19
2.5.1	Soil sampling and preparation	22
2.5.2	Plant sampling and preparation	22
2.6	Comparison of the extraction methods	24
2.6.1	Autoclave extraction method	24
2.6.2	Hot plate extraction method	24
2.7	Samples Extraction	24
2.7.1	Soil extraction	24
2.7.2	Plant extraction	24
2.8	Determination of total arsenic in soils and edible plants	25
2.9	Standard addition method	26
2.10	Method of validation	26
2.10.1	Detection limit (DL)	26
2.10.2	Precision	27
2.10.3	Accuracy	27
2.10.4	Recovery	27
2.10.5	Linear dynamic range	28
3	RESULTS AND DISSCUSSION	29
3.1	Optimization of parameters used for FIAS 100-AAAnalyst 800	29
3.1.1	Effect of carrier gas flow rate (Ar)	29
3.1.2	Effect of NaBH <sub>4</sub> concentration	29
3.1.3	Effect of HCl concentration	30
3.1.4	Effect of Potassium iodide / Ascorbic concentration	31

## CONTENTS (CONTINUED)

	<b>Page</b>
3.1.5 Effect of reduction time	32
3.1.6 Effect of atomization temperature	32
3.2 Comparison of the method used for extraction	33
3.3 Standard addition	35
3.4 Method of validation	36
3.4.1 Detection limit (DL)	36
3.4.2 Precision	36
3.4.3 Accuracy	36
3.4.4 Percent Recovery	36
3.4.5 Linear dynamic range	37
3.5 Total amount of arsenic in soil and edible plant samples	37
3.5.1 Arsenic level in soil	37
3.5.2 Arsenic level in plants	40
3.6 Relationship between arsenic contents of soil and plant	44
3.7 Risk assessment study	48
4 <u>CONCLUSION</u>	51
<u>REFERENCES</u>	53
<u>APPENDICES</u>	62
<u>A</u>	63
<u>B</u>	66
<u>C</u>	68
<u>D</u>	79
<u>E</u>	80
<u>VITAE</u>	81

## LIST OF TABLES

<b>Table</b>	<b>Page</b>
<u>1- 1</u> <u>Some arsenic compounds and their chemical structure which are found in the environment</u>	2
<u>1- 2</u> <u>Arsenic contents in the soil of various countries</u>	4
<u>1- 3</u> <u>Analytical method used with arsenic determination</u>	7
<u>1- 4</u> <u>Estimated daily intake of arsenic for each individual</u>	10
<u>1- 5</u> <u>Arsenic concentration from various environmental media in Ronphibun Sub-district</u>	12
<u>2- 1</u> <u>The Village names of the sampling area in Ronphibun Sub-district, Nakhorn Si Thammarat</u>	20
<u>3- 1</u> <u>Arsenic concentration in Certified Reference Material (CRM) PACS-2</u>	36
<u>3- 2</u> <u>Bioconcentration factor value (BCF) of each plant growing on Ronphibun Sub-district Nakhorn Si Thammarat</u>	46
<u>3- 3</u> <u>Risk index of people in Ronphibun Sub-district consuming arsenic contaminated food</u>	50
<u>A- 1</u> <u>Stock solution and all chemical reagents list</u>	63
<u>A- 2</u> <u>The parameters used for Perkin Elmer model 5000</u>	63
<u>A- 3</u> <u>The parameters used for Perkin Elmer FIAS -AAanlyst 800</u>	64
<u>A- 4</u> <u>Name and GPS position of each sampling site</u>	65
<u>C- 1</u> <u>Optimization of carrier gas (argon)</u>	68
<u>C- 2</u> <u>The effect of NaBH<sub>4</sub> concentration on the peak area of arsine</u>	68
<u>C- 3</u> <u>The effect of HCl concentration (%v/v) on the peak height and peak area of arsine</u>	68
<u>C- 4</u> <u>The effect of KI/ Ascorbic acid reagent using as reducing agent on the peak area of arsine</u>	69
<u>C- 5</u> <u>The effect of reduction time on the peak area of arsine</u>	69
<u>C- 6</u> <u>The effect of optimization temperature on the peak area of arsine</u>	69
<u>C- 7</u> <u>Comparison of the method used for extraction</u>	70
<u>C- 8</u> <u>The result of the standard addition in soil (B<sub>79/1</sub> M<sub>9</sub>)</u>	70
<u>C- 9</u> <u>The result of the standard addition in <i>Curcuma Longa.</i> plant (B<sub>79/1</sub> M<sub>9</sub>)</u>	70

## LIST OF TABLES (CONTINUED)

<b>Table</b>	<b>Page</b>
<u>C- 10</u> <u>The detection limit for arsenic measurement using hydride generation of AAS Perkin Model 5000</u>	71
<u>C- 11</u> <u>The detection limit for arsenic measurement using hydride generation of FIAS 100-AAAnalyst 800</u>	71
<u>C- 12</u> <u>The % RSD calculation from soil sample (M<sub>14</sub>B<sub>204</sub>)</u>	72
<u>C- 13</u> <u>The % RSD calculation from <i>Carica</i> sp. (M<sub>2</sub>B<sub>10</sub>)</u>	72
<u>C- 14</u> <u>Percent recovery</u>	73
<u>C- 15</u> <u>Linear dynamic range for AAAnalyst 800 and Perkin Elmer Model 5000</u>	73
<u>C- 16</u> <u>Arsenic concentration in soil samples at the Ronphibun Sub-district</u>	74
C- 17 Arsenic concentration in edible plants	73
C- 18 The bioconcentration factor of edible plant sample	76
<u>D- 1</u> <u>Total population in Ronphibun Sub-district, Nakhorn Si Thammarat</u>	79

## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
<u>1- 1 Biological transformations of arsenic</u>	3
<u>1- 2 Illustration of skin cancer caused by arsenic</u>	9
<u>1- 3 Map of Ronphibun Sub-district, Nakhorn Si Thammarat Province</u>	11
<u>2- 1 Perkin Elmer model 5000 coupled with Flow injection system</u>	16
<u>2- 2 Perkin Elmer model AAnalyst 800 coupled with Flow injection analysis system 100 (FIAS100)</u>	16
<u>2- 3 The sampling location</u>	21
<u>2- 4 Compositing sampling protocol</u>	22
<u>2- 5 Edible plants collected from Villages No. 1, 2, 8, 9, 11, 13 and 14 grown in Ronphibun Sub-district</u>	23
<u>2- 6 Samples extraction (A: soil extraction. and B: plant extraction)</u>	25
<u>3- 1 The effect of carrier gas (argon) flow rate on the peak area of arsine generated from FIAS -AAnalyst 800 system</u>	29
<u>3- 2 The effect of NaBH<sub>4</sub> concentration on the peak area of arsine generated from FIAS 100- AAnalyst 800 system</u>	30
<u>3- 3 The effect of HCl concentration (%v/v) on the peak height and peak area of arsine generated from FIAS 100- AAnalyst 800 system</u>	31
<u>3- 4 The effect of KI/ Ascorbic acid reagent using as reducing agent on the peak area of arsine generated from FIAS 100- AAnalyst 800 system</u>	31
<u>3- 5 The effect of reduction time on the peak area of arsine generated from FIAS 100-AAnalyst 800 system</u>	32
<u>3- 6 The effect of optimization temperature on the peak area of arsine generated from FIAS 100-AAnalyst 800 system</u>	33
<u>3- 7 Peak are generated from extractants of soil samples for extraction using autoclave and hot plate</u>	34
<u>3- 8 Comparison standard calibration curve and standard addition curve method for soil sample</u>	35
<u>3- 9 Comparison standard calibration curve and standard addition curve method for plant sample</u>	35

## LIST OF FIGURES (CONTINUED)

<b>Figure</b>	<b>Page</b>
<u>3- 10 Peak are generated from FIAS 100 -AAlyst 800 system (A)</u>	37
<u>3- 11 Distribution of arsenic concentration in soil samples collected from Villages No. 1, 2, 8, 9, 11, 12, 13 and 14 in Ronphibun Sub-district, Nakhorn Si Thammarat</u>	39
<u>3- 12 Box and outlier plot presents <math>Q_1</math>, <math>Q_2</math> and <math>Q_3</math> of arsenic level in soil of each Village (Moo) in Ronphibun Sub-district Nakhorn Si Thammarat</u>	40
<u>3- 13 Box plot of arsenic concentration presented in plants that have edible root</u>	41
<u>3- 14 Box plot of arsenic concentration presented in the plants that have edible leaves</u>	42
<u>3- 15 Box plot of arsenic concentration presented in the plants that have edible fruit</u>	43
<u>3- 16 The relationship between soil arsenic and accumulated in different part of plants growing in Ronphibun Sub-district Nakhorn Si Thammarat</u>	45
3-17 Box plot of arsenic concentrations of arsenic accumulated in edible part of plants which are grown in both high and low contaminated area	45