

## CONTENTS

	<b>Page</b>
บทคัดย่อ	iii
ABSTRACT	v
ACKNOWLEDGEMENT	vii
CONTENTS	viii
LIST OF TABLES	xi
LIST OF FIGURES	xiii
CHAPTER	
1. INTRODUCTION	1
2. LITERATURE REVIEW	3
2.1 Botanical description of <i>Senna alata</i>	3
2.2 Thai traditional use of <i>S. alata</i>	4
2.2.1 Used as laxative	4
2.2.2 Treatment of ringworm or <i>Tinea versicolor</i>	5
2.3 Ethnomedical uses	5
2.4 Chemical constituents of <i>S. alata</i>	6
2.5 Determination of hydroxyanthracene derivatives	8
2.6 Biological activities of <i>S. alata</i> extracts	9
2.7 Biological activity of anthraquinones	13
2.7.1 Aloe-emodin	13
2.7.2 Rhein	15
2.7.3 Emodin	16
2.7.4 Chrysophanol	18
2.8 Biosynthesis of anthraquinones	18
2.9 Standardized of herbal extract	21
3. MATERIALS AND METHODS	27
3.1 Materials	27
3.1.1 Plant material	27
3.1.2 Chemicals and reagents	27



## CONTENTS (Continued)

	<b>Page</b>
4.4.2 Determination of the moisture content	55
4.4.3 Determination of the total ash content	56
4.4.4 Determination of the microbial contamination	57
4.5 Solubility of the anthraquinone high-yielding <i>S. alata</i> leaf extract	59
4.6 Partition coefficient	59
4.7 Stability tests	60
4.7.1 Effect of light on the stability of the extract	60
4.7.2 Effect of temperature on the stability of the extract	62
4.7.3 Effect of accelerated condition on the stability of the extract	64
4.7.4 Effect of pH on the stability of the extract	65
5. CONCLUSIONS	68
BIBLIOGRAPHY	70
VITAE	83

## LIST OF TABLES

Table	Page
2-1 Chemical constituents of <i>S. alata</i>	6
3-1 General instrumentation equipment	29
3-2 Linear ranges and correlation coefficients of the calibration curves	34
3-3 Solubility criteria of the extract in various solvents	39
4-1 Anthraquinone content in <i>S. alata</i> leaf extracts extracted under reflux with various concentrations of hydrochloric acid in methanol	43
4-2 Anthraquinone content in <i>S. alata</i> leaf extracts extracted under reflux with various concentrations of ferric chloride in 5 %v/v hydrochloric acid in methanol	46
4-3 Anthraquinone content in <i>S. alata</i> leaf extracts extracted under reflux with various concentrations of water in 5 %v/v hydrochloric acid and 5 %w/v ferric chloride in methanol	48
4-4 Anthraquinone content in <i>S. alata</i> leaf extracts isolated by two different chromatographic methods	50
4-5 Antifungal activity of anthraquinone high-yielding <i>S. alata</i> leaf extract and standard anthraquinones	52
4-6 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts	55
4-7 Moisture content of the anthraquinone high-yielding <i>S. alata</i> leaf extract	56
4-8 Total ash content of the anthraquinone high-yielding <i>S. alata</i> leaf extract	57
4-9 Determination of microbial contamination in the anthraquinone high-yielding <i>S. alata</i> leaf extracts	57
4-10 Solubility of the anthraquinone high-yielding <i>S. alata</i> leaf extract	59
4-11 Partition coefficient values of anthraquinone in anthraquinone high-yielding <i>S. alata</i> leaf extract	60
4-12 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts stored under light and protected from light conditions	61
4-13 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts stored under 4 °C and 30 °C	63

## LIST OF TABLES (Continued)

Table	Page
4-14 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts stored under accelerated condition	64
4-15 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts in the solution at pH 5.5, 7.0, and 8.0	66

## LIST OF FIGURES

Figure	Page
2-1 <i>Senna alata</i> (L.) Roxb.	3
2-2 Chemical structures of anthraquinones	12
2-3 Biogenesis pathways of anthraquinones	20
3-1 Silica gel vacuum column chromatography	31
3-2 Anion exchange chromatography	33
3-3 Calibration curves of standards; rhein (A), aloë-emodin (B), emodin (C), and chrysophanol (D)	35
4-1 HPLC chromatograms of standard anthraquinones (A), methanolic extract (B), and the 5 %v/v HCl in methanol extract (C)	44
4-2 Anthraquinone content in <i>S. alata</i> leaf extracts extracted under reflux with various concentrations of hydrochloric acid in methanol	45
4-3 Hydrolysis of emodin-8-glucoside in acid condition	45
4-4 Oxidation of sennoside A	46
4-5 Anthraquinone content in <i>S. alata</i> leaf extracts extracted under reflux with various concentrations of ferric chloride in 5 %v/v hydrochloric acid in methanol	47
4-6 Anthraquinone content in <i>S. alata</i> leaf extracts extracted under reflux with various concentrations of water in 5 %v/v hydrochloric acid and 5 %w/v ferric chloride in methanol	48
4-7 HPLC chromatograms of <i>S. alata</i> leaf extract isolated by silica gel vacuum column chromatography (A) and anion exchange chromatography (B)	50
4-8 Anthraquinone content in <i>S. alata</i> leaf extracts isolated by two different chromatographic methods	51
4-9 Anthraquinone high-yielding <i>S. alata</i> leaf extract	53
4-10 HPLC chromatograms of standard; aloë-emodin, rhein, emodin, and chrysophanol (A) and anthraquinone high-yielding <i>S. alata</i> leaf extract (B)	54
4-11 Determination of microbial contaminations in the anthraquinone high-yielding <i>S. alata</i> leaf extract; aerobic bacteria(A), <i>Escherichia coli</i> (B), and fungi(C)	58

## LIST OF FIGURES

Figure	Page
4-12 Physical appearance of the anthraquinone high-yielding <i>S. alata</i> leaf extracts kept in well-closed containers protected from light (A) and exposed to light (B)	61
4-13 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts stored under light and protected from light conditions	62
4-14 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts stored under 4 °C and 30 °C	63
4-15 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts stored under accelerated condition	65
4-16 Anthraquinone content of the anthraquinone high-yielding <i>S. alata</i> leaf extracts in the solution at pH 5.5, 7.0, and 8.0	66
4-17 HPLC chromatograms of the anthraquinone high-yielding <i>S. alata</i> extracts in alkali solution (pH 8.0) at the initial time (A) and after 12 weeks (B)	67