

## BIBLIOGRAPHY

- ฐิติรัตน์ ภูมิสถิตย์ และ ณมน คงแก้ว. 2543. การศึกษาการสร้างสารแอนทราควิโนนและฤทธิ์ต้านออกซิเดชันจากเนื้อเยื่อเพาะเลี้ยงขุมเห็ดเทศ. รายงานการวิจัย คณะเภสัชศาสตร์ มหาวิทยาลัยสงขลานครินทร์.
- เต็ม สมิตินันท์. 2544. ชื่อพรรณไม้แห่งประเทศไทย. กรุงเทพฯ: สวนพฤกษศาสตร์ป่าไม้ สำนักวิชาการป่าไม้ กรมป่าไม้.
- สำนักงานสาธารณสุขมูลฐานและคณะกรรมการงสาธารณสุข. 2541. สมุนไพรในงานสาธารณสุขมูลฐาน. กรุงเทพฯ: โรงพิมพ์องค์การสงเคราะห์ทหารผ่านศึก.
- Abd El-Fattah, H.; Abd El-Halim, B.; Nagaya, H; Takeya, K and Itokawa, H. 1997. Cytotoxic bianthrone C-glycosides from *Asphodelus aestivus* tubers. *Alexandria J. Pharm. Sci.* 11: 77-81.
- Abdullah, M.A.; Ali, A.M.; Marziah, M.; Lajis, N.H. and Ariff, A.B. 1998. Establishment of cell suspension cultures of *Morinda elliptica* for the production of anthraquinones. *Plant Cell, Tissue and Organ Culture.* 54: 173-182.
- Abdullah, M.A.; Ariff, A.B.; Mahmood Marziah Ali, A.M. and Lajis, N.H. 2000. Growth and anthraquinone production of *Morinda elliptica* cell suspension cultures in a stirred-tank bioreactor. *J. Agric. Food Chem.* 48: 4432-4438.
- Adedayo, O.; Anderson, W.A.; Moo-Young, M.; Snieckus, V.; Patil, P.A. and Kolawole, D.O. 2001. Phytochemistry and antibacterial activity of *Senna alata* flower. *Pharm. Bio.* 39: 408-412.

- Agarwal, J.S.; Rastogi, R.P. and Srivastava, O.P. 1976. *In vitro* toxicity of constituents of *Rumex maritimus* Linn. to ringworm fungi. *Curr. Sci.* 45: 619-20.
- Agarwal, S.K.; Singh, S.S.; Verma, S. and Kumar, S. 2000. Antifungal activity of anthraquinone derivatives from *Rheum emodi*. *J. Ethnopharmacology.* 72: 43-46.
- Alam, A.I; Mamedov, V.A.; Gubaidullin, A.T.; Kalita, D. and Tsuboi, S. 2003. Isolation and identification of 2,3,7-tri-O-methylelagic acid from *Cassia alata* leaves. *Nat. Medicines.* 57: 73.
- Ali-Emmanuel, N.; Moudachirou, M.; Akakpo, J.A. and Quetin-Leclercq, J. 2003. Treatment of bovine dermatophilosis with *Senna alata*, *Lantana camara* and *Mitracarpus scaber* leaf extracts. *J. Ethnopharmacology.* 86: 167-71.
- Arosio, B.; Gagliano, N.; Fusaro, L.M.; Parmeggiani, L.; Tagliabue, J.; Galetti, P.; De Castri, D.; Moscheni, C. and Annoni G. 2000. Aloe-emodin quinone pretreatment reduces acute liver injury induced by carbon tetrachloride. *Pharmacol. and Toxicology.* 87: 229-33.
- Assane, M.; Traore, M.; Bassene, E. and Sere A. 1993. Choleric effects of *Cassia alata* Linn in the rat. *Dakar medical.* 38: 73-7.
- Adedayo, O.; Anderson, W.A.; Moo-Young, M.; Snieckus, V.; Patil, P.A. and Kolawole, D.O. 2001. Phytochemistry and antibacterial activity of *Senna alata* flower. *Pharm. Bio.* 39: 408-412.
- Balboa, G. and Lim-Sylianco, Y. 1992. Antigenotoxic effects of drug preparations Akapulko and Ampalaya. *Philippine J. Sci.* 121: 399-411.

- Barnard, D.L.; Huffman, J.H.; Morris, J. L.; Wood, S.G.; Hughes, B.G. and Sidwell, R.W. 1992. Evaluation of the antiviral activity of anthraquinones, anthrones and anthraquinone derivatives against human cytomegalovirus. *Antiviral Res.* 17: 63-77.
- Bassetti, L.; Pijnenburg, J. and Tramper, J. 1996. Silicone-stimulated anthraquinone production and release by *Morinda citrifolia* in a two-liquid-phase system. *Biotechnol. Lett.* 18: 377-382.
- Bosch, R.; Friederich, U.; Lutz, W.K.; Brocker, E.; Bachmann, M and Schlatter, C. 1987. Investigations on DNA binding in rat liver and in Salmonella and on mutagenicity in the Ames test by emodin, a natural anthraquinone. *Mutation Res.* 188: 161-168.
- Bruggeman, I.M. and van der Hoeven, J.C. 1984. Lack of activity of the bacterial mutagen emodin in HGPRT and SCE assay with V79 Chinese hamster cells. *Mutation Res.* 138: 219-224.
- Bulgakov, V.P.; Tchernoded, G.K.; Mischenko, N.P.; Khodakovskaya, M.V.; Glazunov, V.P.; Radchenko, S.V.; Zvereva, E.V.; Fedoreyev, S.A. and Zhuravlev, Y.N. 2002. Effect of salicylic acid, methyl jasmonate, ethephon and cantharidin on anthraquinone production by *Rubia cordifolia* callus cultures transformed with the rolB and rolC genes. *J. Biotechnol.* 97: 213-221.
- Cai, Y. and Chen, Q. 1988. Biochemical study of Chinese rhubarb. XXVIII. Comparison of biological activities of the metabolites of anthraquinone derivatives. *Zhongguo Yaoke Daxue Xuebao.* 19: 282-284.
- Camacho, M.R.; Kirby, G.C.; Warhurst, D.C.; Croft, S.L and Phillipson, J.D. 2000. Oxoaporphine alkaloids and quinones from *Stephania dinklagei* and evaluation of their antiprotozoal activities. *Planta Med.* 66: 478-480.

- Castiglione, S.; Fanciulli, M.; Bruno, T.; Evangelista, M.; Del Carlo, C.; Paggi, M.G.; Chersi, A. and Floridi, A. 1993. Rhein inhibits glucose uptake in Ehrlich ascites tumor cells by alteration of membrane-associated functions. *Anti-Cancer Drugs*. 4: 407-414.
- Chang, C.; Ashendel, L.; Geahlen, L.; Mclaughlin, L. and Waters, J. 1996. Oncogene signal transduction inhibitors from medicinal plants. *In Vivo*. 10: 185-190.
- Chan, C.K.; Chang, C.; Koonchanok, M. and Geahlen, L. 1993. Selective inhibition of the growth of ras-transformed human bronchial epithelial cells by emodin, a protein-tyrosine kinase inhibitor. *Biochem. Biophys. Res. Commun*. 193: 1152-1158.
- Chang, Z.; Guo, D.; Shen, X.; Wang, S. and Zheng, J. 1998. Anthraquinone production and analysis in hairy root cultures of *Rheum palmatum* L. *Acta Pharmaceutica Sinica*. 33: 869-872.
- Charlwood, B.V. and Rhodes, M.J.C. 1990. secondary products from plant tissue culture. Oxford: Clarendon Press.
- Chen, Y.; Shen, S.; Lee, W.; Hsu, F.; Lin, H.; Ko, C. and Tseng, S. 2002. Emodin induces apoptosis in human promyeloleukemic HL-60 cells accompanied by activation of caspase 3 cascade but independent of reactive oxygen species production. *Biochem. Pharmacol*. 64: 1713-1724.
- Chiu, P.Y.; Mak, D.H.; Poon, M.K.; Ko, K.M. 2002. *In vivo* antioxidant action of a lignan-enriched extract of Schisandra fruit and an anthraquinone-containing extract of *Polygonum* root in comparison with schisandrin B and emodin. *Planta Med*. 68: 951-956.

- Choi, J.S.; Lee, H.J.; Park, K.Y.; Ha, J.O. and Kang, S.S. 1997. *In vitro* antimutagenic effects of anthraquinone aglycones and naphthopyrone glycosides from *Cassia tora*. *Planta Med.* 63: 11-14.
- Choi, J.S.; Chung, H.Y.; Jung, H.A.; Park, H.J. and Yokozawa, T. 2000. Comparative Evaluation of antioxidant potential of alaternin (2-Hydroxyemodin) and emodin. *J. Agric. Food Chem.* 48: 6347-6351.
- Chung, G.; Tsou, F.; Wang, H.; Lo, H.; Hsieh, E.; Yen, S.; Wu, T.; Chang, H.; Ho, C. and Hung, F. 1998. Rhein affects arylamine N-acetyltransferase activity in *Helicobacter pylori* from peptic ulcer patients. *J. Appl. Toxicology.* 18: 117-123.
- Chung, J.; Li, Y.; Lee, Y.; Lin, J.; Cheng, K. and Chang, W. 2003. Aloe-emodin inhibited N-acetylation and DNA adduct of 2-aminofluorene and arylamine N-acetyltransferase gene expression in mouse leukemia L 1210 cells. *Leukemia Res.* 27: 831-840.
- Clauss, W.; Domokos, G. and Leng-Peschlow, E. 1988. Effect of rhein on electrogenic chloride secretion in rabbit distal colon. *Pharmacology.* 36: 104-110.
- Crockett, C.O.; Guede-Guina, F.; Pugh, D.; Vangah-Manda, M.; Robinson, T.J.; Olubadewo, J.O and Ochillo, R.F. 1992. *Cassia alata* and the preclinical search for therapeutic agents for the treatment of opportunistic infections in AIDS patients. *Cellul. Mol. Biol.* 38: 799-802.
- Cusido, R.; Palazon, J.; Teresa Pinol, M.; Bonfill, M. and Morales, C. 1999. *Datura metel*: *in vitro* production of tropane alkaloids. *Planta Med.* 65: 144-148
- D'angelo, L. 1993. Effects of cascarosides and their metabolites on colonic intestinal muscle. *Acta Toxicologica et Therapeutica.* 14: 193-197.

- Damodaran, S. and Venkataraman, S. 1994. A study on the therapeutic efficacy of *Cassia alata* Linn. leaf extract against *Pityriasis versicolor*. *J. Ethnopharmacology*. 42: 19-23.
- Danoux, L.; Pauly, G. and Moser, P. 2002. Use of extracts of the *Cassia alata* plant. (Cognis France S.A., Fr.). PCT Int. Appl.
- Dewick, M. 2001. Medicinal natural products, a biosynthetic approach. Chichester: John Wiley & Sons Ltd.
- Didry, N.; Dubreuil, L and Pinkas, M. 1994. Activity of anthraquinonic and naphthoquinonic compounds on oral bacteria. *Pharmazie*. 49: 681-683.
- Dipti, J. 1991. Flavonoid glycosides from *Cassia alata*. *Phytochemistry*. 30: 61-63.
- Dixon, R. A. 1991. Plant cell culture: A practical approach. Washington DC: IRL press.
- Dornenburg, H. and Knorr, D. 1994. Effectiveness of plant-derived and microbial polysaccharides as elicitors for anthraquinone synthesis in *Morinda citrifolia* cultures. *J. Agric. Food Chem.* 42: 1048-1052
- Dornenburg, H. and Knorr, D. 1996. Semicontinuous processes for anthraquinone production with immobilized *Cruciata glabra* cell cultures in a three-phase system. *J. Biotechnol.* 50: 55-62
- Effraim, K. D.; Sodipo, O. A. and Jacks, T. W. 1999. Antihepatotoxic activity of aqueous extract of *Cassia alata* (Linn) leaves against carbon tetrachloride induced liver damage in rats. *Pakistan Veterinary Journal*. 19: 111-114.

- Elujoba, A.A., Ajulo, O.O. and Iweibo, G. O. 1989. Chemical and biological analyses of Nigerian *Cassia* species for laxative activity. *J. Pharmaceut. Biomed.* 7: 1453-1457.
- Evans, W.C. 2002. Trease and Evans pharmacognosy. Harcourt Publisher Limited.
- Fairbairn, J.W. and Moss, J.R. 1970. Relative purgative activities of 1,8-dihydroxyanthracene derivatives. *J. Pharm. Pharmacol.* 22: 584-593.
- Farnsworth, N.R. and Bunyapraphatsara, N. 1992. Thai medicinal plant, recommended for primary health care system. Bangkok: Prachachon.
- Floridi, A.; Gentile, P.; Bruno, T.; Castiglione, S.; Zeuli, M and Benassi, M. 1990. Growth inhibition by rhein and lonidamine of human glioma cells *in vitro*. *Anticancer Res.* 10: 1633-1636.
- Floridi, A.; Gentile, P.F.; Bruno, T.; Delpino, A.; Iacobini, C. and Benassi, M. 1992. Survival response of a human glioma cell line to hyperthermia associated with rhein. *Anti-Cancer Drugs.* 3: 407-411.
- Fowler, M. W. 1983. Commercial applications and economic aspects of mass plant cell culture. Plant biotechnology. Cambridge: Cambridge University Press.
- Frieling, T.; Rupprecht, C. and Schemann, M. 1993. Rhein stimulates electrogenic chloride secretion by activation of submucosal neurons in guinea pig colon. *Pharmacology.* 47: 70-76.
- Fuzellier, M.C.; Mortier, F. and Lectard, P. 1982. Antifungal activity of *Cassia alata* L. *Annales Pharmaceutiques Francaises.* 40: 357-363.
- Grimminger, W. and Witthohn, K. 1993. Analytics of senna drugs with regard to the toxicological discussion of anthranoids. *Pharmacology.* 47: 98-109.

- Gross M; Levy R and Toepke H. 1984. Occurrence and analysis of the mycotoxin emodin. *Nahrung*. 28: 31-44.
- Guo, H.; Chang, Z.; Yang, R.; Guo, D. and Zheng, J. 1998. Anthraquinones from hairy root cultures of *Cassia obtusifolia*. *Phytochemistry*. 49: 1623-1625.
- Gupta, D.S.; Jann, B.; Bajpai, K.S. and Sharma, S.C. 1987. Structure of a galactomannan from *Cassia alata* seed. *Carbohydr. Res.* 162: 271-276.
- Hagendoorn, M.J.M.; Jamar, D.C.L.; Meykamp, B. and Plas, L.H.W. 1997. Cell division versus secondary metabolite production in *Morinda citrifolia* cell suspensions. *J. Plant Phys.* 150: 325-330.
- Hagendoorn, M.J.M.; Plas, L.H.W. and Segers, G.J. 1994. Accumulation of anthraquinones in *Morinda citrifolia* cell suspensions. A model system for the study of the interaction between secondary and primary metabolism. *Plant Cell, Tissue and Organ Culture*. 38: 227-234.
- Harrison, J. and Garro, C. 1977. Study on anthraquinone derivatives from *Cassia alata* L. (Leguminosae). *Revista Peruana de Bioquímica*. 1: 31-32.
- Hauptmann, H. and Nazario, L. 1950. Some constituents of the leaves of *Cassia alata*. *J. Am. Chem. Soc.* 72: 1492-1495.
- Heidemann, A.; Miltenburger, H.G. and Mengs, U. 1993. The genotoxicity status of senna. *Pharmacology*. 47: 178-186.
- Heijden, R.; Verpoorte, R.; Hoekstra, S.S. and Hoge, J.H.C. 1994. Nordamnacanthal, a major anthraquinone from an *Agrobacterium rhizogenes* induced root culture of *Rubia tinctorum*. *Plant Phys. Biochem. (Paris)*. 32: 399-404.



- Hemlata Kalidhar, S.B. 1994. Alatonal, an anthraquinone derivative from *Cassia alata*. *Indian J. Chem.* 33: 92-93.
- Huang, S.; Yeh, S. and Hong, C. 1995. Effect of anthraquinone derivatives on lipid peroxidation in rat heart mitochondria: structure-activity relationship. *J. Nat. Prod.* 58: 1365-1371.
- Ibrahim, D. and Osman, H. 1995. Antimicrobial activity of *Cassia alata* from Malaysia. *J. Ethnopharmacology.* 45: 151-156.
- Iosi, F.; Santini, M.T. and Malorni, W. 1993. Membrane and cytoskeleton are intracellular targets of rhein in A431 cells. *Anticancer Res.* 13: 545-554.
- Jiang, X. and Zhen, Y. 1999. Reversal of multi drug resistance by emodin in cancer cells. *Yaoxue Xuebao.* 34: 164-167.
- Jing, X.; Ueki, N.; Cheng, J.; Imanishi, H. and Hada, T. 2002. Induction of apoptosis in hepatocellular carcinoma cell lines by emodin. *Japanese J. Cancer Res.* 93: 874-882.
- Kagedal, K.; Bironaite, D. and Ollinger, K. 1999. Anthraquinone cytotoxicity and apoptosis in primary cultures of rat hepatocytes. *Free Radical Res.* 31: 419-428.
- Kamei, H.; Koide, T.; Kojima, T.; Hashimoto, Y. and Hasegawa, M. 1998. Inhibition of cell growth in culture by quinones. *Cancer Biotherapy and Radiopharmaceuticals.* 13: 185-188.
- Kelly, T. Ross; M., Zhenkun; X. and Wei. E.F. 1994. Revision of the structure of alatinone to emodin. *Phytochemistry.* 36: 253-254.
- Khan, G. and Kapoor, V.P. 1999. Rheological studies of some potential seed galactomannans. *Trends in Carbohydrate Chemistry.* 4: 99-107.

- Khan, M.R.; Kihara, M. and Omoloso A.D. 2001. Antimicrobial activity of *Cassia alata*. *Fitoterapia*. 72: 561-564.
- Kim, L.; Koonchanok, M.; Geahlen, L.; Ashendel, L. and Chang, Jer. 1997. Structure activity relationship study of emodin derivatives based on the protein-tyrosine kinase and protein kinase C inhibitory activities and cytotoxicity. *Nat. Prod. Lett.* 10: 173-180.
- Koyama, J.; Morita, I.; Tagahara, K.; Nobukuni, Y.; Mukainaka, T.; Kuchide, M.; Tokuda, H. and Nishino, H. 2002. Chemopreventive effects of emodin and cassiamin B in mouse skin carcinogenesis. *Cancer Lett.* 182: 135-139.
- Krivobok, S.; Seigle-Murandi, F.; Steiman, R.; Marzin, D.R. and Betina, V. 1992. Mutagenicity of substituted anthraquinones in the Ames/Salmonella microsome system. *Mutation Res.* 279: 1-8.
- Kuo, P.; Lin, T. and Lin, C. 2002. The antiproliferative activity of aloe-emodin is through p53-dependent and p21-dependent apoptotic pathway in human hepatoma cell lines. *Life Sci.* 71: 1879-1892.
- Kuo, Y.; Sun, C.; Ou, J. and Tsai, W. 1997. A tumor cell growth inhibitor from *Polygonum hypoleucum* Ohwi. *Life Sci.* 61: 2335-2344.
- Kupchan, M. and Karim, A. 1976. Tumor inhibitors. 114. Aloe emodin: antileukemic principle isolated from *Rhamnus frangula* L. *Lloydia.* 39: 223-224.
- Lai, G.; Zhang, Z. and Sirica, E. 2003. Celecoxib acts in a cyclooxygenase-2-independent manner and in synergy with emodin to suppress rat cholangiocarcinoma growth in vitro through a mechanism involving enhanced Akt inactivation and increased activation of caspases-9 and -3. *Mol. Cancer Therapeutics.* 2: 265-271.

- Lee, H.Z. 2001. Protein kinase C involvement in aloe-emodin- and emodin-induced apoptosis in lung carcinoma cell. *British J. Pharmacol.* 134: 1093-1103.
- Lee, H. and Tsai, S.J. 1991. Effect of emodin on cooked-food mutagen activation. *Food and Chem. Toxicology.* 29: 765-770.
- Lewis, D.F.; Ioannides, C. and Parke, D.V. 1996. COMPACT and molecular structure in toxicity assessment: a prospective evaluation of 30 chemicals currently being tested for rodent carcinogenicity by the NCI/NTP. *Environ. Health Perspectives.* 104: 1011-1016.
- Liberman, D.F.; Schaefer, F.L.; Fink, R.C.; Ramgopal, M.; Ghosh, A.C. and Mulcahy, R. 1980. Mutagenicity of islandicin and chrysophanol in the Salmonella/microsome system. *Appl. Environ. Microbiol.* 40: 476-479.
- Lin, C.C.; Chang, C.H.; Yang, J.J.; Namba, T. and Hattori, M. 1996. Hepatoprotective effects of emodin from *Ventilago leiocarpa*. *J. Ethnopharmacology.* 52: 107-111.
- Lin, S.; Li, J.; Fujii, M. and Hou, D. 2003. Rhein inhibits TPA-induced activator protein-1 activation and cell transformation by blocking the JNK-dependent pathway. *Int. J. Oncology.* 22: 829-833.
- Liu, M.; Koya, S.; Furuta, H. and Matsuzaki, S. 1996. Growth-inhibiting activity of anthraquinones and benzoquinones against methicillin-resistant *Staphylococcus aureus* (MRSA). *J. Med.Sci.* 23: 85-93.
- Lu, M. and Chen, Q. 1989. Biochemical study of Chinese rhubarb. XXIX. Inhibitory effects of anthraquinone derivatives on P388 leukemia in mice. *Zhongguo Yaoke Daxue Xuebao.* 20: 155-157.

- Malterud, K.E.; Farbrot, T.L.; Huse, A.E. and Sund, R.B. 1993. Antioxidant and radical scavenging effects of anthraquinones and anthrones. *Pharmacology*. 47: 77-85.
- Manitto, P. and Sammes, P.G. 1981. Biosynthesis of natural products. Ellis Horwood: Chichester
- Mantrova, O.V.; Dunaeva, M.V.; Kuzovkina, I.N.; Schneider, B. and Muller-Uri, F. 1999. Effect of methyl jasmonate on anthraquinone biosynthesis in transformed madder roots. *Russian J. Plant Phys.* 46: 248-251.
- Mastura, M. and Khozirah, S. 2000. Antimicrobial activity of *Senna alata* and *Cassia javanica* subsp. *nodosa* extracts against micro-organisms related to skin infections. *J. Tropical Forest Products*. 6: 218-221.
- Masuda, T. and Ueno, Y. 1984. Microsomal transformation of emodin into a direct mutagen. *Mutation Res.* 125: 135-144.
- Matsumoto, O.; Kiyohara, H.; Yamada, A.; Arakawa, K.; Cho, M. and Ohtsuka, Y. 1985. Anti-Bacteroides fragilis components in rhubarb and senna. *Wakan Iyaku Gakkaishi*. 2: 248-249.
- Medicinal Plant Research Institute. 2002. Standard of Thai herbal medicine: *Senna alata* (L.) Roxb. Nonthaburi: Department of Medical Sciences, Ministry of Public Health.
- Medina, N.P.; Viernes, J.D. and Gundran, R. 2001. Evaluation of some medicinal plants against *Escherichia coli*. *Philippines Journal of Veterinary Medicine*. 38: 9-14.
- Mengs, U. and Heidemann, A. 1993. Genotoxicity of sennosides and rhein *in vitro* and *in vivo*. *Medical Science Res.* 21: 749-750.

- Mengs, U.; Krumbiegel, G. and Volkner, W. 1997. Lack of emodin genotoxicity in the mouse micronucleus assay. *Mutation Res.* 393: 289-293.
- Mian, M.; Azzara', A.; Benetti, D.; Ruocco, L.; Bertelli, A. and Ambrogi, F. 1987. Studies *in vitro* on the effects of rhein on the chemotaxis of human leukocytes. *Int. J. Tiss. Res.* 9: 459-463.
- Miralles, J. and Gaydou, E. M. 1986. Fatty acid and sterol composition of oils extracted from seeds of three *Cassia* (Caesalpiniaceae) species from Senegal. *Revue Francaise des Corps Gras.* 33: 381-384.
- Mischenko, N.P.; Fedoreyev, S.A.; Glazunov, V.P.; Chernoded, G.K.; Bulgakov, V.P. and Zhuravlev, Y.N. 1999. Anthraquinone production by callus cultures of *Rubia cordifolia*. *Fitoterapia.* 70: 552-557.
- Mizutani, H.; Hashimoto, O.; Nakashima, R. and Nagai, J. 1997. Anthraquinone production by cell suspension cultures of *Rubia akane* Nakai. *Bioscience Biotechnol. Biochem.* 61: 1743-1744.
- Morah, F.N.I. and Otumu, H.E. 1991. *Cassia alata* seed constituents. *Jamaican J. Sci. Technol.* 2: 14-16.
- Morita, H.; Umeda, M.; Masuda, T. and Ueno, Y. 1988. Cytotoxic and mutagenic effects of emodin on cultured mouse carcinoma FM3A cells. *Mutation Res.* 204: 329-332.
- Moriyama, H.; Iizuka, T.; Nagai, M. and Murata, Y. 2003a. HPLC quantification of kaempferol-3-O-gentiobioside in *Cassia alata*. *Fitoterapia.* 74: 425-430.

- Moriyama, H.; Iizuka, T.; Nagai, M.; Miyataka, H and Satoh, T. 2003b. Antiinflammatory activity of heat-treated *Cassia alata* leaf extract and its flavonoid glycoside. *Yakugaku Zasshi. J. Pharm. Soc. Japan.* 123: 607-611.
- Mueller, O.; Eckert, I.; Lutz, W.K. and Stopper, H. 1996. Genotoxicity of the laxative drug components emodin, aloe-emodin and danthron in mammalian cells: Topoisomerase II mediated. *Mutation Res.* 371: 165-173.
- Mueller, O.; Lutz, K. and Stopper, H. 1998. Factors affecting the genotoxic potency ranking of natural anthraquinones in mammalian cell culture systems. *Mutation Res.* 414: 125-129.
- Mueller, S.O.; Schmitt, M.; Dekant, W.; Stopper, H.; Schlatter, J.; Schreier, P. and Lutz, W.K 1999. Occurrence of emodin, chrysophanol and physcion in vegetables, herbs and liquors. Genotoxicity and anti-genotoxicity of the anthraquinones and of the whole plants. *Food Chem. Toxicol.* 37: 481-491.
- Mueller, S.O. and Stopper, H. 1999. Characterization of the genotoxicity of anthraquinones in mammalian cells. *Biochimica et Biophysica Acta.* 1428: 406-414.
- Narayanaswamy, S. 1994. Plant cell and tissue culture. New Delhi: Tata McGraw-Hill Publishing Company Limited.
- National Drug Committee. 2000. National List of Essential Drug A.D. 1999 (List of Herbal Medicine). Bangkok: Agricultural Cooperative of Thailand Press Ltd.

- Nazif, N.M.; Rady, M.R. and Seif El-Nasr, M.M. 2000. Stimulation of anthraquinone production in suspension cultures of *Cassia acutifolia* by salt stress. *Fitoterapia*. 71: 34-40.
- Nguyen, N.S.; Thai, V.N. and Hoang, D.B. 2002a. Anthraquinones from the leaves of *Cassia alata* L.,Caesalpiniaceae. *Tap Chi Hoa Hoc*. 40: 11-12.
- Nguyen, N.S.; Thai, V.N. and Hoang, D.B. 2002b. Study on the flavonoid isolated from the leaves of *Cassia alata* L, Caeslapiniaceae. *Tap Chi Hoa Hoc*. 40: 1-4.
- Ogasawara, T.; Chiba, K. and Tada, M. 1993. Production in high-yield of a naphthoquinone by a hairy root culture of *Sesamum indicum*. *Phytochemistry*. 33: 1095-1098.
- Ogunti, E.O. and Elujoba, A.A. 1993. Laxative activity of *Cassia alata*. *Fitoterapia*. 64: 437-439
- Palanichamy, S.; Bhaskar, E.A.; Bakthavathsalam, R. and Nagarajan, S. 1991. Wound healing activity of *Cassia alata*. *Fitoterapia*. 62: 153-156
- Palanichamy, S. and Nagarajan, S. 1990a. Anti-inflammatory activity of *Cassia alata* leaf extract and kaempferol 3-O-sophoroside. *Fitoterapia*. 61: 44-47.
- Palanichamy, S and Nagarajan, S. 1990b. Antifungal activity of *Cassia alata* leaf extract. *J. Ethnopharmacology*. 29: 337-340.
- Palanichamy, S.; Nagarajan, S. and Devasagayam, M. 1988. Effect of *Cassia alata* leaf extract on hyperglycemic rats. *J. Ethnopharmacology*. 22: 81-90.

- Panichayupakaranant, P. and Kaewsuwan, S. 2004. Bioassay-guided isolation of the antioxidant constituent from *Cassia alata* L. leaves. *Songklanakarin J. Sci. Technol.* 26: 103-107
- Panichayupakaranant, P. and Intaraksa, N. 2003. Distribution of hydroxyanthracene derivatives in *Cassia alata* and the factors affecting the quality of the raw material. *Songklanakarin J. Sci. Technol.* 25: 497-502.
- Pecere, T.; Gazzola, M.V.; Mucignat, C.; Parolin, C.; Vecchia, F.D.; Cavaggioni, A.; Basso, G.; Diaspro, A.; Salvato, B.; Carli, M. and Palu, G. 2000. Aloe-emodin is a new type of anticancer agent with selective activity against neuroectodermal tumors. *Cancer Res.* 60: 2800-2804.
- Pecere, T.; Sarinella, F.; Salata, C.; Gatto, B.; Bet, A.; Dalla, V.; Diaspro, A.; Carli, M.; Palumbo, M. and Palu, G. 2003. Involvement of p53 in specific anti-neuroectodermal tumor activity of aloe-emodin. *Int. J. Cancer.* 106: 836-847.
- Rady, M.R. and Nazif, N.M. 1997. Response of explant type to proliferation and anthraquinones accumulation in *Cassia acutifolia*. *Fitoterapia.* 68: 349-354.
- Rai, K.N. and Prasad, S.N. 1994. Chemical examination of the stem of *Cassia alata* Linn. *J. Indian Chem. Soc.* 71: 653-654.
- Rai, P.P. 1978. Anthracene derivatives in leaves and fruits of *Cassia alata*. *Curr. Sci.* 47: 271-272.
- Rao, S.; Sastry, R.; Rao, K. and Vimaladevi, M. 1975. Occurrence of kaempferol and aloe-emodin in the leaves of *Cassia alata*. *Curr. Sci.* 44: 736-737.



- Sahm, H. 1993. Biotechnology 2<sup>nd</sup> edition vol.1. Biological Fundamentals. Weinheim: VCH Verlagsgesellschaft mbh.
- Sakharkar, P.R. and Pati, A.T. 1998. Antimicrobial activity of *Cassia alata*. *Indian J. Pharm. Sci.* 60: 311-312.
- Sarg, T.M.; Salem, S.A.; Farrag, N.M.; Abdel-Aal, M.M. and Ateya, A.M. 1990. Phytochemical and antimicrobial investigation of *Hemerocallis fulva* L. grown in Egypt. *Int. J. Crude Drug Res.* 28: 153-156.
- Schorkhuber, M.; Richter, M.; Dutter, A.; Sontag, G. and Marian, B. 1998. Effect of anthraquinone-laxatives on the proliferation and urokinase secretion of normal, premalignant and malignant colonic epithelial cells. *European J. Cancer.* 34: 1091-1098.
- Semple, J.; Pyke, M.; Reynolds, D. and Flower, P. 2001. *In vitro* antiviral activity of the anthraquinone chrysophanic acid against poliovirus. *Antiviral Res.* 49: 169-178.
- Sen, K.; Sarkar, K.; Banerjee, N. and Farooqi, H. 1987. Structural investigation of a water-soluble galactomannan from the seeds of *Cassia alata* Linn. *Indian J. Chem.* 26: 21-25.
- Sepchr, M.F. and Ghorbanli, M. 2002. Effects of nutritional factors on the formation of anthraquinones in callus cultures of *Rheum ribes*. *Plant Cell, Tissue and Organ Culture.* 68: 171-175.
- Serrame, E.; Wu, L.S. and Sylianco, C.Y.L. 1995. Inhibition of activity of genotoxic tumor promoters by two Philippines medicinal plants. *Philippines J. Sci.* 124: 53-55.
- Shargool, P.D. 1982. *Appl. Biochem. Biotech.* 7: 239-257.

- Sheludko, Y.; Gerasimenko, I.; Kolshorn, H. and Stöckigt, J. 2002a. Isolation and structure elucidation of a new indole alkaloid from *Rauvolfia serpentina* hairy root culture: the first naturally occurring alkaloid of the raumacline group. *Planta Med.* 68: 435-439.
- Sheludko, Y.; Gerasimenko, I.; Kolshorn, H. and Stöckigt, J. 2002b. New alkaloids of the sarpagine group from *Rauvolfia serpentina* hairy root culture. *J. Nat. Prod.* 65: 1006-1010.
- Shi, Y.Q; Fukai, T.; Sakagami, H.; Kuroda, J.; Miyaoka, R.; Tamura, M.; Yoshida, N. and Nomura, T. 2001. Cytotoxic and DNA damage-inducing activities of low molecular weight phenols from rhubarb. *Anticancer Res.* 21: 2847-2853.
- Shim, J.; Shin, J.; Pai, T.; Chung, I. and Lee, H. 1999. Permeabilization of elicited suspension culture of madder (*Rubia akane* Nakai) cells for release of anthraquinones. *Biotechnol. Tech.* 13: 249-252.
- Shin, S. and Lyu, L. 2000. Effects of compounds related to signal transduction on anthraquinone biosynthesis in transformed cells of *Rubia cordifolia* var. *pratensis*. *Korean J. Pharmacognosy.* 31: 235-239.
- Siddhuraju, P.; Mohan, P.S. and Becker, K. 2002. Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. *Food Chem.* 79: 61-67.
- Singh, B.K. and Tiwari, R.D. 1943. Chemical examination of *Cassia alata*. The component acids of the fatty oil from the seeds. *Proceedings of the National Academy of Sciences, India.* 13: 111-119.
- Singh, R.B. 1998. Polyalcohols from *Cassia alata* Linn. seeds. *Asian J. Chem.* 10: 185-186.

- Smith, M. and Ali, S. 1979. Anthraquinones from the leaves of *Cassia alata* from Fiji. *New Zealand J. Sci.* 22: 123-125.
- Sodipo, O. A.; Effraim, K. D. and Emmagun, E. 1998. Effect of aqueous leaf extract of *Cassia alata* (Linn.) on some hematological indices in albino rats. *Phytotherapy Res.* 12: 431-433.
- Somchit, M.N.; Reezal, I; Nur, E. and Mutalib, A.R. 2003. *In vitro* antimicrobial activity of ethanol and water extracts of *Cassia alata*. *J. Ethnopharmacology.* 84: 1-4.
- Stark, A.A; Townsend, J.M.; Wogan, G.N.; Demain, A.L.; Manmade, A. and Ghosh, A.C. 1978. Mutagenicity and antibacterial activity of mycotoxins produced by *Penicillium islandicum* Sopp and *Penicillium rugulosum*. *J. Environ. Pathol. Toxicol.* 2: 313-324.
- Strobel, J.; Hieke, M. and Groger, D. 1991. Increased anthraquinone production in *Galium verum* cell cultures induced by polymeric adsorbents. *Plant Cell, Tissue and Organ Culture.* 24: 207-210.
- Subcommittee on the establishment of the Thai Herbal Pharmacopoeia. 1998. Thai Herbal Pharmacopoeia. vol. 1. Bangkok: Prachachon.
- Sydiskis, R.J; Owen, D.G; Lohr, J.L; Rosler, K.H and Blomster, R.N. 1991. Inactivation of enveloped viruses by anthraquinones extracted from plants. *Antimicrobial agents and Chemotherapy.* 35: 2463-2466.
- Tavares, I.A.; Mascolo, N.; Izzo, A.A. and Capasso, F. 1996. Effects of anthraquinone derivatives on PAF release by human gastrointestinal mucosa in vitro. *Phytotherapy Res.* 10: S20-S21.

- Tegos, G.; Stermitz, F.R.; Lomovskaya, O. and Lewis, K. 2002. Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials. *Antimicrobial Agents and Chemotherapy*. 46: 3133-3141.
- Thamlikitkul, V.; Bunyapraphatsara, N.; Dechatiwongse, T.; Theerapong, S.; Chantrakul, C.; Thanaveerasuwan, T.; Nimitnon, S.; Boonroj, P.; Punkrut, W. and Gingsungneon, V. 1990. Randomized controlled trial of *Cassia alata* Linn. for constipation. *J. the Medical Association of Thailand*. 73: 217-222.
- Tikkanen, L.; Matsushima, T. and Natori, S. 1983. Mutagenicity of anthraquinones in the Salmonella preincubation test. *Mutation Res*. 116: 297-304.
- Tiwari, R.D. and Yadava, O.P. 1971. Structural study of the quinone pigments from the roots of *Cassia alata*. *Planta Med*. 19: 299-305.
- Ukhun, M.E. and Ifebigh, E.O. 1988. Compositional chemistry of *Cassia alata* seeds. *Food Chem*. 30: 205-210.
- Vargas, F.; Fraile, G.; Velasquez, M.; Correia, H.; Fonseca, G.; Marin, M.; Marcano, E. and Sanchez, Y. 2002. Studies on the photostability and phototoxicity of aloe-emodin, emodin and rhein. *Pharmazie*. 57: 399-404.
- Villasenor, I.M.; Canlas, A.P.; Pascua, M.; Sabando, M.N and Soliven, L. 2002. Bioactivity studies on *Cassia alata* Linn. leaf extracts. *Phytotherapy Res*. 16: S93-S96.
- Villaroya, M. and Bernal-Santos, R. 1976. A chemical investigation of *Cassia alata* L. (fam. Leguminosae). *Asian J. Pharmacy*. 3: 10-12.

- Wang, C; Huang, Y.; Chen, L.; Lee, L. and Yang, L. 2002. Inducible nitric oxide synthase inhibitors of chinese herbs III. *Rheum palmatum*. *Planta Med.* 68: 869-874.
- Wang, H.H; Chung, J.G; Ho, C.C; Wu, L T and Chang, S.H. 1998. Aloe-emodin effects on arylamine N-acetyltransferase activity in the bacterium *Helicobacter pylori*. *Planta Med.* 64: 176-178.
- Wang, H. 1993. Antitrichomonal action of emodin in mice. *J. Ethnopharmacology.* 40: 111-16.
- Wang, H.W; Chen, T.L.; Yang, P.C. and Ueng, T.H. 2001. Induction of cytochromes P450 1A1 and 1B1 by emodin in human lung adenocarcinoma cell line CL5. *Drug Metabolism and Disposition.* 29: 1229-1235.
- Wang, W.; Chen, C.; Chen, Q. and Cai, Y. 1990. Biochemical study of Chinese rhubarb XXXIX. Inhibition of anthraquinone derivatives on anaerobic bacteria. *Zhongguo Yaoke Daxue Xuebao.* 21: 354-357.
- Wanitschke, R. 1980. Influence of rhein on electrolyte and water transfer in the isolated rat colonic mucosa. *Pharmacology.* 20: 21-26.
- Wanitschke, R and Karbach, U. 1988. Influence of rhein on rat colonic sodium, potassium-ATPase and permeability *in vitro*. *Pharmacology.* 36: 98-103.
- Westendorf, J.; Marquardt, H.; Poginsky, B.; Dominiak, M.; Schmidt, J. and Marquardt, H. 1990. Genotoxicity of naturally occurring hydroxyanthraquinones. *Mutation Res.* 240: 1-12.

- Wolfe, D.; Schmutte, C.; Westendorf, J. and Marquardt H. 1990. Hydroxyanthraquinones as tumor promoters: enhancement of malignant transformation of C3H mouse fibroblasts and growth stimulation of primary rat hepatocytes. *Cancer Res.* 50: 6540-6544.
- Woo, S.; Nan, J.; Lee, S.; Park, E.; Zhao, Y. and Sohn, D. 2002. Aloe emodin suppresses myofibroblastic differentiation of rat hepatic stellate cells in primary culture. *Pharmacol. Toxicol.* 90: 193-198.
- Wu, C.H.; Hsieh, C.L.; Song, T.Y. and Yen, G.C. 2001. Inhibitory Effects of *Cassia tora* L. on Benzo[a]pyrene-Mediated DNA Damage toward Hep G2 Cells. *J. Agric. Food Chem.* 49: 2579-2586.
- Yadav, K. and Kalidhar, B. 1994. Alquinone: an anthraquinone from *Cassia alata*. *Planta Med.* 60: 601.
- Yamamoto, M.; Masui, T.; Sugiyama, K.; Yokota, M.; Nakagomi, K. and Nakazawa, H. 1991. Anti-inflammatory active constituents of *Aloe arborescens* Miller. *Agric. Biol. Chem.* 55: 1627-1629.
- Yang, W.; Yu, H.; Xu, W.; Yu, Q. and Cui, Y. 1994. The effects of emodin on electrical and contractive activities of the smooth muscle cells in guinea pig taenia coli. *Shengwu Wuli Xuebao.* 10: 82-86.
- Yeh, F.; Wu, C. and Lee, H. 2003. Signaling pathway for aloe-emodin-induced apoptosis in human H460 lung nonsmall carcinoma cell. *Int. J. Cancer.* 106: 26-33.
- Yen, G.; Duh, P. and Chuang, D. 2000. Antioxidant activity of anthraquinones and anthrone. *Food Chem.* 70: 437-441.

- Yin, C. and Xu, C. 1998. Effect of aloe-emodin on proliferation of vascular smooth muscle cells after arterial injury. *Chung-kuo Chung Hsi I Chieh Ho Tsa Chih*. 18: 420-422.
- Yuan, Z. and Gao, R. 1997. Antioxidant actions of anthraquinolines contained in *Rheum*. *Pharm. Pharmacol. Lett.* 7: 9-12.
- Zhang, L.; Chang, C.; Bacus, S. and Hung, M. 1995. Suppressed transformation and induced differentiation of HER-2/neu-overexpressing breast cancer cells by emodin. *Cancer Res.* 55: 3890-3896.
- Zhang, L. and Hung, M. 1996. Sensitization of HER-2/neu-overexpressing non-small cell lung cancer cells to chemotherapeutic drugs by tyrosine kinase inhibitor emodin. *Oncogene*. 12: 571-576.
- Zhang, L.; Lau, Y.; Xia, W.; Hortobagyi, G.; Hung, M. 1999. Tyrosine kinase inhibitor emodin suppresses growth of HER-2/neu-overexpressing breast cancer cells in athymic mice and sensitizes these cells to the inhibitory effect of paclitaxel. *Clin. Cancer Res.* 5: 343-353.
- Zhan, Y.; Li, D.; Wei, H.; Wang, Z.; Huang, X.; Xu, Q. and Lu, H. 2000. Emodin on hepatic fibrosis in rats. *Chinese Medical Journal*. 113: 599-601.
- Zhao, W.Q.; Polya, G.M.; Wang, B.H.; Gibbs, M.E.; Sedman, G.L. and Ng, K.T. 1995. Inhibitors of cAMP-dependent protein kinase impair long-term memory formation in day-old chicks. *Neurobiology of Learning and Memory*. 64: 106-118.
- Zhu, F.; Liu, X. and Liang, N. 2003. Effect of emodin and apigenin on invasion of human ovarian carcinoma HO-8910PM cells *in vitro*. *Ai Zheng*. 22: 358-362.