

REFERENCES

- Adams, D., Samuel, D., Goulon-Goeau, C., Nakazato, M., Costa, P. M., Feray, C., Plante, V., Ducot, B., Ichai, P., Lacroix, C., Metral, S., Bismuth, H., and Said, G. (2000). The course and prognostic factors of familial amyloid polyneuropathy after liver transplantation. *Brain*. 123 (Pt 7): 1495-504.
- Aleshire, S. L., Bradley, C. A., Richardson, L. D., and Parl, F. F. (1983). Localization of human prealbumin in choroid plexus epithelium. *J Histochem Cytochem*. 31(5): 608-12.
- Almeida, M. R., Alves, I. L., Terazaki, H., Ando, Y., and Saraiva, M. J. (2000). Comparative studies of two transthyretin variants with protective effects on familial amyloidotic polyneuropathy: TTR R104H and TTR T119M. *Biochem Biophys Res Commun*. 270(3): 1024-8.
- Almeida, M. R., Macedo, B., Cardoso, I., Alves, I., Valencia, G., Arsequell, G., Planas, A., and Saraiva, M. J. (2004). Selective binding to transthyretin and tetramer stabilization in serum from patients with familial amyloidotic polyneuropathy by an iodinated diflunisal derivative. *Biochem J*. 381(Pt 2): 351-6.
- Alves, I. L., Altland, K., Almeida, M. R., Winter, P., and Saraiva, M. J. (1997a). Screening and biochemical characterization of transthyretin variants in the Portuguese population. *Hum Mutat*. 9(3): 226-33.
- Alves, M., Conceicao, I., and Luis, M. L. (1997b). Neurophysiological evaluation of sexual dysfunction in familial amyloidotic polyneuropathy--Portuguese type. *Acta Neurol Scand*. 96(3): 163-6.
- Andersson, R. (1970). Hereditary amyloidosis with polyneuropathy. *Acta Med Scand*. 1-2(1): 85-94.
- Andersson, R. (1976). Familial amyloidosis with polyneuropathy. A clinical study based on patients living in northern Sweden. *Acta Med Scand Suppl*. 590: 1-64.
- Ando, Y., and Suhr, O. B. (1998). Autonomic dysfunction in familial amyloidotic polyneuropathy (FAP). *Amyloid*. 5(4): 288-300.

- Andrade, C. (1952). A peculiar form of peripheral neuropathy; familiar atypical generalized amyloidosis with special involvement of the peripheral nerves. *Brain*. 75(3): 408-27.
- Araki, S. (1984). Type I familial amyloidotic polyneuropathy (Japanese type). *Brain Dev.* 6(2): 128-33.
- Araki, S., Mawatari, S., Ohta, M., Nakajima, A., and Kuroiwa, Y. (1968). Polyneuritic amyloidosis in a Japanese family. *Arch Neurol.* 18(6): 593-602.
- Baker, C. M., Manwell, C., Labisky, R. F., and Harper, J. A. (1966). Molecular genetics of avian proteins--V. Egg, blood and tissue proteins of the ring-necked pheasant, *Phasianus colchicus* L. *Comp Biochem Physiol.* 17(2): 467-99.
- Bellovino, D., Morimoto, T., Pisaniello, A., and Gaetani, S. (1998). In vitro and in vivo studies on transthyretin oligomerization. *Exp Cell Res.* 243(1): 101-12.
- Bellovino, D., Morimoto, T., Tosetti, F., and Gaetani, S. (1996). Retinol binding protein and transthyretin are secreted as a complex formed in the endoplasmic reticulum in HepG2 human hepatocarcinoma cells. *Exp Cell Res.* 222(1): 77-83.
- Bennhold, H. (1922). Specific staining of amyloid by Congo red [German]. *Münchener Medizinische Wochenschrift*, 69, 1537 1538.
- Bhat, M. K., Sreekrishna, K., and Cama, H. R. (1977). Heterogeneity of chicken prealbumin & specificity in binding to chicken retinol-binding protein & L-thyroxine. *Indian J Biochem Biophys.* 14(2): 125-8.
- Birch, H. E., and Schreiber, G. (1986). Transcriptional regulation of plasma protein synthesis during inflammation. *J Biol Chem.* 261(18): 8077-80.
- Birnboim, H. C., and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7(6): 1513-23.
- Blake, C., and Serpell, L. (1996). Synchrotron X-ray studies suggest that the core of the transthyretin amyloid fibril is a continuous beta-sheet helix. *Structure*. 4(8): 989-98.
- Blake, C. C. (1981). Prealbumin and the thyroid hormone nuclear receptor. *Proc R Soc Lond B Biol Sci.* 211(1185): 413-31.

- Blake, C. C., Geisow, M. J., Oatley, S. J., Rérat, B., and Rérat, C. (1978). Structure of prealbumin: secondary, tertiary and quaternary interactions determined by Fourier refinement at 1.8 Å. *J Mol Biol.* 121(3): 339-56.
- Blake, C. C., Geisow, M. J., Swan, I. D., Rérat, C., and Rérat, B. (1974). Structure of human plasma prealbumin at 2-5 Å resolution. A preliminary report on the polypeptide chain conformation, quaternary structure and thyroxine binding. *J Mol Biol.* 88(1): 1-12.
- Blake, C. C., and Oatley, S. J. (1977). Protein-DNA and protein-hormone interactions in prealbumin: a model of the thyroid hormone nuclear receptor? *Nature.* 268(5616): 115-20.
- Booth, D. R., Gillmore, J. D., Lachmann, H. J., Booth, S. E., Bybee, A., Soyturk, M., Akar, S., Pepys, M. B., Tunca, M., and Hawkins, P. N. (2000a). The genetic basis of autosomal dominant familial Mediterranean fever. *Qjm.* 93(4): 217-21.
- Booth, D. R., Gillmore, J. D., Persey, M. R., Booth, S. E., Cafferty, K. D., Tennent, G. A., Madhoo, S., Cochrane, S. W., Whitehead, T. C., Pasvol, G., and Hawkins, P. N. (1998). Transthyretin Ile73Val is associated with familial amyloidotic polyneuropathy in a Bangladeshi family. Mutations in brief no. 158. Online. *Hum Mutat.* 12(2): 135.
- Booth, D. R., Lachmann, H. J., Gillmore, J. D., Booth, S. E., and Hawkins, P. N. (2001). Prevalence and significance of the familial Mediterranean fever gene mutation encoding pyrin Q148. *Qjm.* 94(10): 527-31.
- Booth, D. R., Stangou, A., Williams, R. S., Gillmore, J. D., Tennent, G. A., and Hawkins, P. N. (2000b). Transthyretin Ile84Thr is associated with familial amyloid polyneuropathy. *Hum Mutat.* 16(5): 447.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 72: 248-54.
- Branch, W.T., Jr., Robbins, J. and Edelhoch, H. (1971). Thyroxine-binding prealbumin: conformation in aqueous solution. *J Biol Chem.* 246, 6011-6018.

- Branch, W. T., Robbins, J., and Edelhoch, H. (1972). Thyroxine-binding prealbumin. Conformation in urea and guanidine. *Arch Biochem Biophys.* 152(1): 144-51.
- Breathnach, R., Benoist, C., O'Hare, K., Gannon, F., and Chambon, P. (1978). Ovalbumin gene: evidence for a leader sequence in mRNA and DNA sequences at the exon-intron boundaries. *Proc Natl Acad Sci U S A.* 75(10): 4853-7.
- Burton, P., Iden, S., Mitchell, K., and White, A. (1978). Thymic hormone-like restoration by human prealbumin of azathioprine sensitivity of spleen cells from thymectomized mice. *Proc Natl Acad Sci U S A.* 75(2): 823-7.
- Burton, P. M., Horner, B. L., Jones, G. H., Lin, T., Nestor, J. J., Jr., Newman, S. R., Parks, T. L., Smith, A. J., and White, A. (1987). Immuno-enhancing activity of the amino-terminal domain of human prealbumin: isolation, characterization and synthesis. *Int J Immunopharmacol.* 9(3): 297-305.
- Burton, P. M., Hung, P., Lin, T., Lovelace, C., and White, A. (1985). The effects of homogeneous human prealbumin on *in vitro* and *in vivo* immune responses in the mouse. *Int J Immunopharmacol.* 7(4): 473-81.
- Buxbaum, J., and Gallo, G. (1999). Nonamyloidotic monoclonal immunoglobulin deposition disease. Light-chain, heavy-chain, and light- and heavy-chain deposition diseases. *Hematol Oncol Clin North Am.* 13(6): 1235-48.
- Canijo, M., and Andrade, C. (1969). Familial amyloidotic polyneuropathy. Electromyographic study. *J Genet Hum.* 17(3): 281-8.
- Cardoso, I., Merlini, G., and Saraiva, M. J. (2003). 4'-ido-4'-deoxydoxorubicin and tetracyclines disrupt transthyretin amyloid fibrils *in vitro* producing noncytotoxic species: screening for TTR fibril disrupters. *Faseb J.* 17(8): 803-9.
- Coelho, T., Chorão, R., Sousa, A., Alves, I.L. Torres, M.F. Saraiva, M.J.M. (1996). Compound heterozygotes of transthyretin Met30 and transthyretin Met119 are protected from the devastating effects of familial amyloid polyneuropathy. *Neuromuscul Disord (suppl).* 6: S20
- Cohen, A. S. (1966). Preliminary chemical analysis of partially purified amyloid fibrils. *Am J Pharm Sci Support Public Health.* 15(1 Pt 1): 66-83.

- Cohen, A. S., and Calkins, E. (1959). Electron microscopic observations on a fibrous component in amyloid of diverse origins. *Nature*. 183(4669): 1202-3.
- Cohen, S. N., Chang, A. C., and Hsu, L. (1972). Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc Natl Acad Sci U S A*. 69(8): 2110-4.
- Coimbra, A., and Andrade, C. (1971a). Familial amyloid polyneuropathy: an electron microscope study of the peripheral nerve in five cases. I. Interstitial changes. *Brain*. 94(2): 199-206.
- Coimbra, A., and Andrade, C. (1971b). Familial amyloid polyneuropathy: an electron microscope study of the peripheral nerve in five cases. II. Nerve fibre changes. *Brain*. 94(2): 207-12.
- Connors, L. H., Lim, A., Prokaeva, T., Roskens, V. A., and Costello, C. E. (2003). Tabulation of human transthyretin (TTR) variants. *Amyloid*. 10(3): 160-84.
- Connors, L. H., Yamashita, T., Yazaki, M., Skinner, M., and Benson, M. D. (2004). A rare transthyretin mutation (Asp18Glu) associated with cardiomyopathy. *Amyloid*. 11(1): 61-6.
- Conway, K. A., Harper, J. D., and Lansbury, P. T. (1998). Accelerated in vitro fibril formation by a mutant alpha-synuclein linked to early-onset Parkinson disease. *Nat Med*. 4(11): 1318-20.
- Cornwell, G. G., 3rd, Sletten, K., Olofsson, B. O., Johansson, B., and Westermark, P. (1987). Prealbumin: its association with amyloid. *J Clin Pathol*. 40(2): 226-31.
- Costa, P. P., Figueira, A. S., and Bravo, F. R. (1978). Amyloid fibril protein related to prealbumin in familial amyloidotic polyneuropathy. *Proc Natl Acad Sci U S A*. 75(9): 4499-503.
- Cregg, J. M., Barringer, K. J., Hessler, A. Y., and Madden, K. R. (1985). *Pichia pastoris* as a host system for transformations. *Mol Cell Biol*. 5(12): 3376-85.
- Cregg, J.M. and Cereghino, J.L. (2000). Heterologous protein expression in the methylotrophic yeast *Pichia pastoris*. *FEMS Microbiology Reviews* 24, 45-66.
- Cregg, J. M., Cereghino, J. L., Shi, J., and Higgins, D. R. (2000). Recombinant protein expression in *Pichia pastoris*. *Mol Biotechnol*. 16(1): 23-52.

- Clegg, J. M., Madden, K. R., Barringer, K. J., Thill, G. P., and Stillman, C. A. (1989). Functional characterization of the two alcohol oxidase genes from the yeast *Pichia pastoris*. *Mol Cell Biol.* 9(3): 1316-23.
- Clegg, J. M., Vedvick, T. S., and Raschke, W. C. (1993). Recent advances in the expression of foreign genes in *Pichia pastoris*. *Biotechnology (N Y)*. 11(8): 905-10.
- Cserr, H. F. (1971). Physiology of the choroid plexus. *Physiol Rev.* 51(2): 273-311.
- Damas, A. M., and Saraiva, M. J. (2000). Review: TTR amyloidosis-structural features leading to protein aggregation and their implications on therapeutic strategies. *J Struct Biol.* 130(2-3): 290-9.
- de Carvalho, M., Moreira, P., Evangelista, T., Ducla-Soares, J.L., Bento, M., Fernandes, R., Saraiva, M.J. (2000). New transthyretin mutation V28M in a Portuguese kindred with amyloid polyneuropathy. *Muscle Nerve* 23, 1016–1021.
- Dickson, P. W., Aldred, A. R., Marley, P. D., Bannister, D., and Schreiber, G. (1986). Rat choroid plexus specializes in the synthesis and the secretion of transthyretin (prealbumin). Regulation of transthyretin synthesis in choroid plexus is independent from that in liver. *J Biol Chem.* 261(8): 3475-8.
- Dickson, P. W., Aldred, A. R., Menting, J. G., Marley, P. D., Sawyer, W. H., and Schreiber, G. (1987). Thyroxine transport in choroid plexus. *J Biol Chem.* 262(29): 13907-15.
- Dickson, P. W., Howlett, G. J., and Schreiber, G. (1982). Metabolism of prealbumin in rats and changes induced by acute inflammation. *Eur J Biochem.* 129(2): 289-93.
- Drueke, T. B. (2000). Beta2-microglobulin and amyloidosis. *Nephrol Dial Transplant.* 15 Suppl 1: 17-24.
- Elliott, P. M., Mahon, N. G., Matsumura, Y., Hawkins, P. N., Gillmore, J. D., and McKenna, W. J. (2000). Tissue Doppler features of cardiac amyloidosis. *Clin Cardiol.* 23(9): 701.
- Ellis, S. B., Brust, P. F., Koutz, P. J., Waters, A. F., Harpold, M. M., and Gingeras, T. R. (1985). Isolation of alcohol oxidase and two other methanol regulatable genes from the yeast *Pichia pastoris*. *Mol Cell Biol.* 5(5): 1111-21.

- Eneqvist, T., Andersson, K., Olofsson, A., Lundgren, E., and Sauer-Eriksson, A. E. (2000). The beta-slip: a novel concept in transthyretin amyloidosis. *Mol Cell.* 6(5): 1207-18.
- Eneqvist, T., and Sauer-Eriksson, A. E. (2001). Structural distribution of mutations associated with familial amyloidotic polyneuropathy in human transthyretin. *Amyloid.* 8(3): 149-68.
- Felding, P., and Fex, G. (1982). Cellular origin of prealbumin in the rat. *Biochim Biophys Acta.* 716(3): 446-9.
- Felding, P., Fex, G., Westermark, P., Olofsson, B. O., Pitkanen, P., and Benson, L. (1985). Prealbumin in Swedish patients with senile systemic amyloidosis and familial amyloidotic polyneuropathy. *Scand J Immunol.* 21(2): 133-40.
- Ferguson, R. N., Edelhoch, H., Saroff, H. A., Robbins, J., and Cahnmann, H. J. (1975). Negative cooperativity in the binding of thyroxine to human serum prealbumin. Preparation of tritium-labeled 8-anilino-1-naphthalenesulfonic acid. *Biochemistry.* 14(2): 282-9.
- Foss, T. R., Wiseman, R. L., and Kelly, J. W. (2005). The pathway by which the tetrameric protein transthyretin dissociates. *Biochemistry.* 44(47): 15525-33.
- Friedreich, N., Kelkulé, A. (1859). Zur Amyloidfrage. *Virchows Arch Pathol Anat Physiol.* 16: 50-56.
- Fung, W. P., Thomas, T., Dickson, P. W., Aldred, A. R., Milland, J., Dziadek, M., Power, B., Hudson, P., and Schreiber, G. (1988). Structure and expression of the rat transthyretin (prealbumin) gene. *J Biol Chem.* 263(1): 480-8.
- Furuya, H., Nakazato, M., Saraiva, M. J., Costa, S. P., Sasaki, H., Matsuo, H., Goto, I., and Sakaki, Y. (1989). Tetramer formation of a variant type human transthyretin (prealbumin) produced by *Escherichia coli* expression system. *Biochem Biophys Res Commun.* 163(2): 851-9.
- Geisse, S., Gram, H., Kleuser, B., and Kocher, H. P. (1996). Eukaryotic expression systems: a comparison. *Protein Expr Purif.* 8(3): 271-82.
- Gillmore, J. D., Lovat, L. B., Persey, M. R., Pepys, M. B., and Hawkins, P. N. (2001). Amyloid load and clinical outcome in AA amyloidosis in relation to circulating concentration of serum amyloid A protein. *Lancet.* 358(9275): 24-9.

- Glenner, G. G. (1981). The bases of the staining of amyloid fibers: their physico-chemical nature and the mechanism of their dye-substrate interaction. *Prog Histochem Cytochem.* 13(3): 1-37.
- Glenner, G. G. (1983). Alzheimer's disease: multiple cerebral amyloidosis. In R. Katzman ed. Banbury Report 15, Biological aspect of Alzheimer's disease. Cold Spring Harbor Laboratory 137-144.
- Glover, J. (1973). Retinol-binding proteins. *Vitam Horm.* 31: 1-42.
- Goodman, D. S., and Raz, A. (1972). Extraction and recombination studies of the interaction of retinol with human plasma retinol-binding protein. *J Lipid Res.* 13(3): 338-47.
- Grinna, L. S., and Tschopp, J. F. (1989). Size distribution and general structural features of N-linked oligosaccharides from the methylotrophic yeast, *Pichia pastoris*. *Yeast.* 5(2): 107-15.
- Guimarães, A., Monteiro, L., Coutinho, P. (1980). Pathology of the autonomic nervous system in andrade type of familial amyloidotic polyneuropathy. In: Glenner, G.G., Costa, P.P., de Freitas, A.F. (Eds.), Amyloid and Amyloidosis. *Excerpta Medica, Amsterdam*, pp. 88-98.
- Gustavsson, A., Jahr, H., Tobiassen, R., Jacobson, D. R., Sletten, K., and Westermark, P. (1995). Amyloid fibril composition and transthyretin gene structure in senile systemic amyloidosis. *Lab Invest.* 73(5): 703-8.
- Haass, C., and Steiner, H. (2001). Protofibrils, the unifying toxic molecule of neurodegenerative disorders? *Nat Neurosci.* 4(9): 859-60.
- Hagen, G. A., and Elliott, W. J. (1973). Transport of thyroid hormones in serum and cerebrospinal fluid. *J Clin Endocrinol Metab.* 37(3): 415-22.
- Hamilton, J. A., Steinrauf, L. K., Braden, B. C., Liepnieks, J., Benson, M. D., Holmgren, G., Sandgren, O., and Steen, L. (1993). The x-ray crystal structure refinements of normal human transthyretin and the amyloidogenic Val-30-->Met variant to 1.7-A resolution. *J Biol Chem.* 268(4): 2416-24.
- Harms, P. J., Tu, G. F., Richardson, S. J., Aldred, A. R., Jaworowski, A., and Schreiber, G. (1991). Transthyretin (prealbumin) gene expression in choroid plexus is strongly conserved during evolution of vertebrates. *Comp Biochem Physiol B.* 99(1): 239-49.

- Hawkins, J. D., Catalano, R. F., Jr., Gillmore, M. R., and Wells, E. A. (1989). Skills training for drug abusers: generalization, maintenance, and effects on drug use. *J Consult Clin Psychol.* 57(4): 559-63.
- Herbert, J., Cavallaro, T., and Martone, R. (1991). The distribution of retinol-binding protein and its mRNA in the rat eye. *Invest Ophthalmol Vis Sci.* 32(2): 302-9.
- Herbert, J., Wilcox, J. N., Pham, K. T., Fremeau, R. T., Jr., Zeviani, M., Dwork, A., Soprano, D. R., Makover, A., Goodman, D. S., Zimmerman, E. A., and et al. (1986). Transthyretin: a choroid plexus-specific transport protein in human brain. The 1986 S. Weir Mitchell award. *Neurology.* 36(7): 900-11.
- Higgins, D. R., and Clegg, J. M. (1998). Introduction to *Pichia pastoris*. *Methods Mol Biol.* 103: 1-15.
- Holmgren, G., Costa, P. M., Andersson, C., Asplund, K., Steen, L., Beckman, L., Nylander, P. O., Teixeira, A., Saraiva, M. J., and Costa, P. P. (1994). Geographical distribution of TTR met30 carriers in northern Sweden: discrepancy between carrier frequency and prevalence rate. *J Med Genet.* 31(5): 351-4.
- Holmgren, G., Haettner, E., Nordenson, I., Sandgren, O., Steen, L., and Lundgren, E. (1988). Homozygosity for the transthyretin-met30-gene in two Swedish sibs with familial amyloidotic polyneuropathy. *Clin Genet.* 34(5): 333-8.
- Horwitz, J. P., Chua, J., Curby, R. J., Tomson, A. J., Darooge, M. A., Fisher, B. E., Mauricio, J., and Klundt, I. (1964). Substrates for Cytochemical Demonstration of Enzyme Activity. I. Some Substituted 3-Indolyl-Beta-D-Glycopyranosides. *J Med Chem.* 7: 574-5.
- Hugo, L.M., (2002). Three-dimensional structure of the transthyretin-retinol-binding protein complex. *Clin Chem Lab Med.* 40(12):1229–1236
- Ikeda, S., Nakazato, M., Ando, Y., and Sobue, G. (2002). Familial transthyretin-type amyloid polyneuropathy in Japan: clinical and genetic heterogeneity. *Neurology.* 58(7): 1001-7.
- Ingbar, S. H. (1958). Pre-albumin: a thyroxinebinding protein of human plasma. *Endocrinology.* 63(2): 256-9.

- Iozzo, R. V., Cohen, I. R., Grassel, S., and Murdoch, A. D. (1994). The biology of perlecan: the multifaceted heparan sulphate proteoglycan of basement membranes and pericellular matrices. *Biochem J.* 302 (Pt 3): 625-39.
- Jacobson, D. R., McFarlin, D. E., Kane, I., and Buxbaum, J. N. (1992). Transthyretin Pro55, a variant associated with early-onset, aggressive, diffuse amyloidosis with cardiac and neurologic involvement. *Hum Genet.* 89(3): 353-6.
- Jacobson, D. R., Pastore, R. D., Yaghoubian, R., Kane, I., Gallo, G., Buck, F. S., and Buxbaum, J. N. (1997). Variant-sequence transthyretin (isoleucine 122) in late-onset cardiac amyloidosis in black Americans. *N Engl J Med.* 336(7): 466-73.
- Jiang, X., Smith, C. S., Petrassi, H. M., Hammarstrom, P., White, J. T., Sacchettini, J. C., and Kelly, J. W. (2001). An engineered transthyretin monomer that is nonamyloidogenic, unless it is partially denatured. *Biochemistry.* 40(38): 11442-52.
- Kabat, E. A., Moore, D. H., and Landow, H. (1942b). An Electrophoretic Study of the Protein Components in Cerebrospinal Fluid and Their Relationship to the Serum Proteins. *J Clin Invest.* 21(5): 571-7.
- Kabat, E.A., Landow, H. and Moore, D.H. (1942a). Electrophoresis patterns of concentrated cerebrospinal fluid. *Proc Natl Acad Sci USA* 49, 260-263.
- Kanai, M., Raz, A., and Goodman, D. S. (1968). Retinol-binding protein: the transport protein for vitamin A in human plasma. *J Clin Invest.* 47(9): 2025-44.
- Kanda, Y., Goodman, D. S., Canfield, R. E., and Morgan, F. J. (1974). The amino acid sequence of human plasma prealbumin. *J Biol Chem.* 249(21): 6796-805.
- Kelly, J. W. (1996). Alternative conformations of amyloidogenic proteins govern their behavior. *Curr Opin Struct Biol.* 6(1): 11-7.
- Kelly, J. W. (1998). The alternative conformations of amyloidogenic proteins and their multi-step assembly pathways. *Curr Opin Struct Biol.* 8(1): 101-6.
- Kelly, J. W., Colon, W., Lai, Z., Lashuel, H. A., McCulloch, J., McCutchen, S. L., Miroy, G. J., and Peterson, S. A. (1997). Transthyretin quaternary and

- tertiary structural changes facilitate misassembly into amyloid. *Adv Protein Chem.* 50: 161-81.
- Kim, H. S., Byun, S. H., and Lee, B. M. (2005). Effects of chemical carcinogens and physicochemical factors on the UV spectrophotometric determination of DNA. *J Toxicol Environ Health A.* 68(23-24): 2081-95.
- Kishikawa, M., Nakanishi, T., Miyazaki, A., and Shimizu, A. (1999). Enhanced amyloidogenicity of sulfonated transthyretin in vitro, a hypothetical etiology of senile amyloidosis. *Amyloid.* 6(3): 183-6.
- Kisilevsky, R., Lyon, A. W., and Young, I. D. (1992). A critical analysis of postulated pathogenetic mechanisms in amyloidogenesis. *Crit Rev Clin Lab Sci.* 29(1): 59-82.
- Klabunde, T., Petrassi, H. M., Oza, V. B., Raman, P., Kelly, J. W., and Sacchettini, J. C. (2000). Rational design of potent human transthyretin amyloid disease inhibitors. *Nat Struct Biol.* 7(4): 312-21.
- Kopelman, M., Mokady, S., and Cogan, U. (1976). Comparative studies of human and chicken retinol-binding proteins and prealbumins. *Biochim Biophys Acta.* 439(2): 442-8.
- Koutz, P., Davis, G. R., Stillman, C., Barringer, K., Cregg, J., and Thill, G. (1989). Structural comparison of the *Pichia pastoris* alcohol oxidase genes. *Yeast.* 5(3): 167-77.
- Laemmli, U.K. and Favre, M. 1973. Maturation of the head of bacteriophage T4. *J Mol Biol* 80, 575-599.
- Lai, Z., Colon, W., and Kelly, J. W. (1996). The acid-mediated denaturation pathway of transthyretin yields a conformational intermediate that can self-assemble into amyloid. *Biochemistry.* 35(20): 6470-82.
- Lai, Z., McCulloch, J., Lashuel, H. A., and Kelly, J. W. (1997). Guanidine hydrochloride-induced denaturation and refolding of transthyretin exhibits a marked hysteresis: equilibria with high kinetic barriers. *Biochemistry.* 36(33): 10230-9.
- Lashuel, H. A., Lai, Z., and Kelly, J. W. (1998). Characterization of the transthyretin acid denaturation pathways by analytical ultracentrifugation: implications

- for wild-type, V30M, and L55P amyloid fibril formation. *Biochemistry*. 37(51): 17851-64.
- Lewis, W. D., Skinner, M., Simms, R. W., Jones, L. A., Cohen, A. S., and Jenkins, R. L. (1994). Orthotopic liver transplantation for familial amyloidotic polyneuropathy. *Clin Transplant*. 8(2 Pt 1): 107-10.
- Linke, R. P. (2000). Highly sensitive diagnosis of amyloid and various amyloid syndromes using Congo red fluorescence. *Virchows Arch*. 436(5): 439-48.
- Liz, M. A., Faro, C. J., Saraiva, M. J., and Sousa, M. M. (2004). Transthyretin, a new cryptic protease. *J Biol Chem*. 279(20): 21431-8.
- Magnus, J. H., Stenstad, T., Husby, G., and Kolset, S. O. (1992). Isolation and partial characterization of heparan sulphate proteoglycans from human hepatic amyloid. *Biochem J*. 288 (Pt 1): 225-31.
- Martone, R. L., Herbert, J., Dwork, A., and Schon, E. A. (1988). Transthyretin is synthesized in the mammalian eye. *Biochem Biophys Res Commun*. 151(2): 905-12.
- McCutchen, S. L., and Kelly, J. W. (1993). Intermolecular disulfide linkages are not required for transthyretin amyloid fibril formation *in vitro*. *Biochem Biophys Res Commun*. 197(2): 415-21.
- McCutchen, S. L., Lai, Z., Miroy, G. J., Kelly, J. W., and Colon, W. (1995). Comparison of lethal and nonlethal transthyretin variants and their relationship to amyloid disease. *Biochemistry*. 34(41): 13527-36.
- Milland, J., Tsykin, A., Thomas, T., Aldred, A. R., Cole, T., and Schreiber, G. (1990). Gene expression in regenerating and acute-phase rat liver. *Am J Physiol*. 259(3 Pt 1): G340-7.
- Miller, J. B. (1963). Pre-Albumin Variations in Primates. *Nature*. 200: 810.
- Miroy, G. J., Lai, Z., Lashuel, H. A., Peterson, S. A., Strang, C., and Kelly, J. W. (1996). Inhibiting transthyretin amyloid fibril formation via protein stabilization. *Proc Natl Acad Sci U S A*. 93(26): 15051-6.
- Misrahi, A. M., Plante, V., Lalu, T., Serre, L., Adams, D., Lacroix, D. C., and Said, G. (1998). New transthyretin variants SER 91 and SER 116 associated with familial amyloidotic polyneuropathy. Mutations in brief no. 151. Online. *Hum Mutat*. 12(1): 71.

- Mita S., Maeda S., Shimada K. and Araki S. (1984). Cloning and sequence analysis of cDNA for human prealbumin. *Biochem Biophys Res Commun* 124: 558–564.
- Montesino, R., Garcia, R., Quintero, O., and Cremata, J. A. (1998). Variation in N-linked oligosaccharide structures on heterologous proteins secreted by the methylotrophic yeast *Pichia pastoris*. *Protein Expr Purif.* 14(2): 197-207.
- Morrissey, J. H. (1981). Silver stain for proteins in polyacrylamide gels: a modified procedure with enhanced uniform sensitivity. *Anal Biochem.* 117(2): 307-10.
- Motojima, K., and Goto, S. (1990). Dual promoters and tissue-specific expression of rat transthyretin gene. *Biochem Biophys Res Commun.* 173(1): 323-30.
- Murrell, J. R., Schoner, R. G., Liepnieks, J. J., Rosen, H. N., Moses, A. C., and Benson, M. D. (1992). Production and functional analysis of normal and variant recombinant human transthyretin proteins. *J Biol Chem.* 267(23): 16595-600.
- Muto, Y., and Goodman, D. S. (1972). Vitamin A transport in rat plasma. Isolation and characterization or retinol-binding protein. *J Biol Chem.* 247(8): 2533-41.
- Nettleton, E. J., Sunde, M., Lai, Z., Kelly, J. W., Dobson, C. M., and Robinson, C. V. (1998). Protein subunit interactions and structural integrity of amyloidogenic transthyretins: evidence from electrospray mass spectrometry. *J Mol Biol.* 281(3): 553-64.
- Neumann, P., Cody, V., and Wojtczak, A. (2001). Structural basis of negative cooperativity in transthyretin. *Acta Biochim Pol.* 48(4): 867-75.
- Nichols, W. C., and Benson, M. D. (1990). Hereditary amyloidosis: detection of variant prealbumin genes by restriction enzyme analysis of amplified genomic DNA sequences. *Clin Genet.* 37(1): 44-53.
- Nilsson, S. F., and Peterson, P. A. (1971). Evidence for multiple thyroxine-binding sites in human prealbumin. *J Biol Chem.* 246(19): 6098-105.
- Nilsson, S. F., Rask, L., and Peterson, P. A. (1975). Studies on thyroid hormone-binding proteins. II. Binding of thyroid hormones, retinol-binding protein, and fluorescent probes to prealbumin and effects of thyroxine on prealbumin subunit self association. *J Biol Chem.* 250(21): 8554-63.

- Okamoto, T., and Okabe, S. (2000). Ultraviolet absorbance at 260 and 280 nm in RNA measurement is dependent on measurement solution. *Int J Mol Med.* 5(6): 657-9.
- Ong, D. E., Davis, J. T., O'Day, W. T., and Bok, D. (1994). Synthesis and secretion of retinol-binding protein and transthyretin by cultured retinal pigment epithelium. *Biochemistry.* 33(7): 1835-42.
- Oppenheimer, J. H. (1968). Role of plasma proteins in the binding, distribution and metabolism of the thyroid hormones. *N Engl J Med.* 278(21): 1153-62.
- Pages, R. A., Robbins, J., and Edelhoch, H. (1973). Binding of thyroxine and thyroxine analogs to human serum prealbumin. *Biochemistry.* 12(14): 2773-9.
- Palha, J. A. (2002). Transthyretin as a thyroid hormone carrier: function revisited. *Clin Chem Lab Med.* 40(12): 1292-300.
- Perz, J. B., Rahemtulla, A., Giles, C., Szydlo, R. M., Davis, J., Gopaul, D., Gillmore, J., Mathias, C. J., Hawkins, P. N., and Apperley, J. F. (2006). Long-term outcome of high-dose melphalan and autologous stem cell transplantation for AL amyloidosis. *Bone Marrow Transplant.* 37(10): 937-43.
- Peterson, P. A. (1971). Studies on the interaction between prealbumin, retinol-binding protein, and vitamin A. *J Biol Chem.* 246(1): 44-9.
- Peterson, S. A., Klabunde, T., Lashuel, H. A., Purkey, H., Sacchettini, J. C., and Kelly, J. W. (1998). Inhibiting transthyretin conformational changes that lead to amyloid fibril formation. *Proc Natl Acad Sci U S A.* 95(22): 12956-60.
- Planté-Bordeneuve, V., Lalu, T., Misrahi, M., Reilly, M. M., Adams, D., Lacroix, C., and Said, G. (1998). Genotypic-phenotypic variations in a series of 65 patients with familial amyloid polyneuropathy. *Neurology.* 51(3): 708-14.
- Prapunpoj, P., Richardson, S. J., and Schreiber, G. (2002). Crocodile transthyretin: structure, function, and evolution. *Am J Physiol Regul Integr Comp Physiol.* 283(4): R885-96.
- Prapunpoj, P., Yamauchi, K., Nishiyama, N., Richardson, S. J., and Schreiber, G. (2000). Evolution of structure, ontogeny of gene expression, and function of

- Xenopus laevis transthyretin. *Am J Physiol Regul Integr Comp Physiol.* 279(6): R2026-41.
- Quintas, A., Saraiva, M. J., and Brito, R. M. (1997). The amyloidogenic potential of transthyretin variants correlates with their tendency to aggregate in solution. *FEBS Lett.* 418(3): 297-300.
- Quintas, A., Saraiva, M. J., and Brito, R. M. (1999). The tetrameric protein transthyretin dissociates to a non-native monomer in solution. A novel model for amyloidogenesis. *J Biol Chem.* 274(46): 32943-9.
- Quintas, A., Vaz, D. C., Cardoso, I., Saraiva, M. J., and Brito, R. M. (2001). Tetramer dissociation and monomer partial unfolding precedes protofibril formation in amyloidogenic transthyretin variants. *J Biol Chem.* 276(29): 27207-13.
- Raghu, P., Ravinder, P., and Sivakumar, B. (2003). A new method for purification of human plasma retinol-binding protein and transthyretin. *Biotechnol Appl Biochem.* 38(Pt 1): 19-24.
- Raghu, P., and Sivakumar, B. (2004). Interactions amongst plasma retinol-binding protein, transthyretin and their ligands: implications in vitamin A homeostasis and transthyretin amyloidosis. *Biochim Biophys Acta.* 1703(1): 1-9.
- Rask, L., Peterson, P. A., and Nilsson, S. F. (1971). The subunit structure of human thyroxine-binding prealbumin. *J Biol Chem.* 246(19): 6087-97.
- Raz, A., and Goodman, D. S. (1969). The interaction of thyroxine with human plasma prealbumin and with the prealbumin-retinol-binding protein complex. *J Biol Chem.* 244(12): 3230-7.
- Redondo C., Damas A. and Saraiva M. (2000). Designing transthyretin mutants affecting tetrameric structure: implications in amyloidogenicity. *Biochem. J.* 348, 167-172.
- Richardson, S. J., Bradley, A. J., Duan, W., Wettenhall, R. E., Harms, P. J., Babon, J. J., Southwell, B. R., Nicol, S., Donnellan, S. C., and Schreiber, G. (1994). Evolution of marsupial and other vertebrate thyroxine-binding plasma proteins. *Am J Physiol.* 266(4 Pt 2): R1359-70.

- Richardson, S. J., Dziegielewska, K. M., Andersen, N. A., Frost, S., and Schreiber, G. (1998). The acute phase response of plasma proteins in the polyprotodont marsupial *Monodelphis domestica*. *Comp Biochem Physiol B Biochem Mol Biol.* 119(1): 183-8.
- Robbins, J., and Rall, J. E. (1957). The interaction of thyroid hormones and protein in biological fluids. *Recent Prog Horm Res.* 13: 161-202; discussion 202-8.
- Rochet, J. C., and Lansbury, P. T., Jr. (2000). Amyloid fibrillogenesis: themes and variations. *Curr Opin Struct Biol.* 10(1): 60-8.
- Röcken, C., Saeger, W., and Linke, R. P. (1994). Gastrointestinal amyloid deposits in old age. Report on 110 consecutive autopsical patients and 98 retrospective bioptic specimens. *Pathol Res Pract.* 190(7): 641-9.
- Rosen, H. N., Murrell, J. R., Liepnieks, J. J., Benson, M. D., Cody, V., and Moses, A. C. (1994). Threonine for alanine substitution at position 109 of transthyretin differentially alters human transthyretin's affinity for iodothyronines. *Endocrinology.* 134(1): 27-34.
- Sambrook, J., Fritsch, E.F. & Maniatis, T.; Hrsg. (1989). *Molecular Cloning - A Laboratory Manual, 2nd Edition.* Cold Spring Harbour Laboratory Press, New York.
- Sanger, F., Nicklen, S., and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A.* 74(12): 5463-7.
- Saraiva, M. J. (2001). Transthyretin mutations in hyperthyroxinemia and amyloid diseases. *Hum Mutat.* 17(6): 493-503.
- Saraiva, M. J. (2002). Hereditary transthyretin amyloidosis: molecular basis and therapeutical strategies. *Expert Rev Mol Med.* 2002: 1-11.
- Saraiva, M. J., Birken, S., Costa, P. P., and Goodman, D. S. (1984). Amyloid fibril protein in familial amyloidotic polyneuropathy, Portuguese type. Definition of molecular abnormality in transthyretin (prealbumin). *J Clin Invest.* 74(1): 104-19.
- Sasaki, H., Yoshioka, N., Takagi, Y., and Sakaki, Y. (1985). Structure of the chromosomal gene for human serum prealbumin. *Gene.* 37(1-3): 191-7.

- Schoner, B. E., Belagaje, R. M., and Schoner, R. G. (1986). Translation of a synthetic two-cistron mRNA in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 83(22): 8506-10.
- Schormann, N., Murrell, J. R., and Benson, M. D. (1998). Tertiary structures of amyloidogenic and non-amyloidogenic transthyretin variants: new model for amyloid fibril formation. *Amyloid.* 5(3): 175-87.
- Schreiber, G., Aldred, A. R., Jaworowski, A., Nilsson, C., Achen, M. G., and Segal, M. B. (1990). Thyroxine transport from blood to brain via transthyretin synthesis in choroid plexus. *Am J Physiol.* 258(2 Pt 2): R338-45.
- Schreiber, G., Prapunpoj, P., Chang, L., Richardson, S.J., Aldred, R.A. and Munro, S.L. (1999). Evolution of thyroid hormone distribution. *Clin Exp Pharmacol Physiol.* 25, 728-732.
- Schreiber, G., and Richardson, S. J. (1997). The evolution of gene expression, structure and function of transthyretin. *Comp Biochem Physiol B Biochem Mol Biol.* 116(2): 137-60.
- Schwarzman, A. L., Tsiper, M., Wente, H., Wang, A., Vitek, M. P., Vasiliev, V., and Goldgaber, D. (2004). Amyloidogenic and anti-amyloidogenic properties of recombinant transthyretin variants. *Amyloid.* 11(1): 1-9.
- Scorer, C. A., Clare, J. J., McCombie, W. R., Romanos, M. A., and Sreekrishna, K. (1994). Rapid selection using G418 of high copy number transformants of *Pichia pastoris* for high-level foreign gene expression. *Biotechnology (N Y).* 12(2): 181-4.
- Sebastião, M. P., Saraiva, M. J., and Damas, A. M. (1998). The crystal structure of amyloidogenic Leu55 --> Pro transthyretin variant reveals a possible pathway for transthyretin polymerization into amyloid fibrils. *J Biol Chem.* 273(38): 24715-22.
- Selkoe, D. J. (1996). Amyloid beta-protein and the genetics of Alzheimer's disease. *J Biol Chem.* 271(31): 18295-8.
- Seibert, F.B. and Nelson, J.W. (1942). Electrophoretic study of the blood protein response in tuberculosis. *J Biol Chem* 143, 29-38.
- Serag, A. A., Altenbach, C., Gingery, M., Hubbell, W. L., and Yeates, T. O. (2001). Identification of a subunit interface in transthyretin amyloid fibrils: evidence

- for self-assembly from oligomeric building blocks. *Biochemistry*. 40(31): 9089-96.
- Serpell LC, Goldstein G, Dacklin I, Lundgren E & Blake CCF (1996). The ‘edge strand’ hypothesis: prediction and test of a mutational ‘hot-spot’ on the transthyretin molecule associated with FAP amyloidogenesis. *Amyloid* 3, 75–85.
- Serpell, L. C., Sunde, M., and Blake, C. C. (1997). The molecular basis of amyloidosis. *Cell Mol Life Sci.* 53(11-12): 871-87.
- Shastry, B. S. (2003). Neurodegenerative disorders of protein aggregation. *Neurochem Int.* 43(1): 1-7.
- Smeland, S., Kolset, S. O., Lyon, M., Norum, K. R., and Blomhoff, R. (1997). Binding of perlecan to transthyretin *in vitro*. *Biochem J.* 326 (Pt 3): 829-36.
- Smith, F. R., and Goodman, D. S. (1971). The effects of diseases of the liver, thyroid, and kidneys on the transport of vitamin A in human plasma. *J Clin Invest.* 50(11): 2426-36.
- Sousa A, Andersson R, Drugge U, Holmgren G, Sandgren O: (1993). Familial amyloidotic polyneuropathy in Sweden: geographical distribution, age of onset, and prevalence. *Hum Heredity*. 43: 288-294.
- Sousa A, Coelho T, Lobato L, Sequeiros J: (1995). Genetic epidemiology of familial amyloidotic polyneuropathy (FAP)-type I (FAP-MET30). in Kyle R (ed): Amyloid and amyloidosis. New York and London: Parthenon; pp 209-211.
- Sousa, M. M., and Saraiva, M. J. (2003). Neurodegeneration in familial amyloid polyneuropathy: from pathology to molecular signaling. *Prog Neurobiol.* 71(5): 385-400.
- Southwell, B. R., Duan, W., Alcorn, D., Brack, C., Richardson, S. J., Kohrle, J., and Schreiber, G. (1993). Thyroxine transport to the brain: role of protein synthesis by the choroid plexus. *Endocrinology*. 133(5): 2116-26.
- Southwell, B. R., Duan, W., Tu, G. F., and Schreiber, G. (1991). Ontogenesis of transthyretin gene expression in chicken choroid plexus and liver. *Comp Biochem Physiol B.* 100(2): 329-38.
- Sparkes, R. S., Sasaki, H., Mohandas, T., Yoshioka, K., Klisak, I., Sakaki, Y., Heinzmann, C., and Simon, M. I. (1987). Assignment of the prealbumin

- (PALB) gene (familial amyloidotic polyneuropathy) to human chromosome region 18q11.2-q12.1. *Hum Genet.* 75(2): 151-4.
- Stratton, J., Chiruvolu, V., and Meagher, M. (1998). High cell-density fermentation. *Methods Mol Biol.* 103: 107-20.
- Suhr, O. B., Svendsen, I. H., Ohlsson, P. I., Lendoire, J., Trigo, P., Tashima, K., Ranlov, P. J., and Ando, Y. (1999). Impact of age and amyloidosis on thiol conjugation of transthyretin in hereditary transthyretin amyloidosis. *Amyloid.* 6(3): 187-91.
- Tashima, K., Ando, Y., Tanaka, Y., Uchino, M., and Ando, M. (1995). Change in the age of onset in patients with familial amyloidotic polyneuropathy type I. *Intern Med.* 34(8): 748-50.
- Taylor, B. A., Bailey, D. W., Cherry, M., Riblet, R., and Weigert, M. (1975). Genes for immunoglobulin heavy chain and serum prealbumin protein are linked in mouse. *Nature.* 256(5519): 644-6.
- Terazaki, H., Ando, Y., Suhr, O., Ohlsson, P. I., Obayashi, K., Yamashita, T., Yoshimatsu, S., Suga, M., Uchino, M., and Ando, M. (1998). Post-translational modification of transthyretin in plasma. *Biochem Biophys Res Commun.* 249(1): 26-30.
- Terpe, K. (2006). Overview of bacterial expression systems for heterologous protein production: from molecular and biochemical fundamentals to commercial systems. *Appl Microbiol Biotechnol.* 72(2): 211-22.
- Thomas, T., Power, B., Hudson, P., Schreiber, G., and Dziadek, M. (1988). The expression of transthyretin mRNA in the developing rat brain. *Dev Biol.* 128(2): 415-27.
- Toyooka, K., Fujimura, H., Ueno, S., Yoshikawa, H., Kaido, M., Nishimura, T., Yorifuji, S., and Yanagihara, T. (1995). Familial amyloid polyneuropathy associated with transthyretin Gly42 mutation: a quantitative light and electron microscopic study of the peripheral nervous system. *Acta Neuropathol (Berl).* 90(5): 516-25.
- Trimble, R. B., Atkinson, P. H., Tschopp, J. F., Townsend, R. R., and Maley, F. (1991). Structure of oligosaccharides on Saccharomyces SUC2 invertase

- secreted by the methylotrophic yeast *Pichia pastoris*. *J Biol Chem.* 266(34): 22807-17.
- Tschopp, J.F., Sverlow, G., Kosson, R., Craig, W. and Grinna, L. (1987). High-level secretion of glycosylated invertase in the methylotrophic yeast, *Pichia pastoris*. *Bio/Technology* 5, 1305-1308.
- Venter JC, Adams M.D., ----- (2001). The sequence of the human genome. *Science* 291, 1304-1351.
- Verma, R., Boleti, E., and George, A. J. (1998). Antibody engineering: comparison of bacterial, yeast, insect and mammalian expression systems. *J Immunol Methods*. 216(1-2): 165-81.
- Virchow, R. (1854). Zur Cellulose-Frage. *Virchow Arch Path Anat Physiol*, 6, 135-137.
- Wakasugi, S., Maeda, S., Shimada, K., Nakashima, H., and Migita, S. (1985). Structural comparisons between mouse and human prealbumin. *J Biochem (Tokyo)*. 98(6): 1707-14.
- Wallace, M. R., Naylor, S. L., Kluge-Beckerman, B., Long, G. L., McDonald, L., Shows, T. B., and Benson, M. D. (1985). Localization of the human prealbumin gene to chromosome 18. *Biochem Biophys Res Commun.* 129(3): 753-8.
- Westermark, P., Benson, M. D., Buxbaum, J. N., Cohen, A. S., Frangione, B., Ikeda, S., Masters, C. L., Merlini, G., Saraiva, M. J., and Sipe, J. D. (2002). Amyloid fibril protein nomenclature -- 2002. *Amyloid*. 9(3): 197-200.
- Westermark, P., Bergstrom, J., Solomon, A., Murphy, C., and Sletten, K. (2003). Transthyretin-derived senile systemic amyloidosis: clinicopathologic and structural considerations. *Amyloid*. 10 Suppl 1: 48-54.
- Westermark, P., Johansson, B., and Natvig, J. B. (1979). Senile cardiac amyloidosis: evidence of two different amyloid substances in the ageing heart. *Scand J Immunol.* 10(4): 303-8.
- Westermark, P., Pitkanen, P., Benson, L., Vahlquist, A., Olofsson, B. O., and Cornwell, G. G., 3rd. (1985). Serum prealbumin and retinol-binding protein in the prealbumin-related senile and familial forms of systemic amyloidosis. *Lab Invest.* 52(3): 314-8.

- Westermark, P., Sletten, K., Johansson, B., and Cornwell, G. G., 3rd. (1990). Fibril in senile systemic amyloidosis is derived from normal transthyretin. *Proc Natl Acad Sci U S A.* 87(7): 2843-5.
- Westermark, P., Wernstedt, C., Wilander, E., and Sletten, K. (1986). A novel peptide in the calcitonin gene related peptide family as an amyloid fibril protein in the endocrine pancreas. *Biochem Biophys Res Commun.* 140(3): 827-31.
- Wilfinger, W. W., Mackey, K., and Chomczynski, P. (1997). Effect of pH and ionic strength on the spectrophotometric assessment of nucleic acid purity. *Biotechniques.* 22(3): 474-6, 478-81.
- Wiseman, R. L., Green, N. S., and Kelly, J. W. (2005). Kinetic stabilization of an oligomeric protein under physiological conditions demonstrated by a lack of subunit exchange: implications for transthyretin amyloidosis. *Biochemistry.* 44(25): 9265-74.
- Woeber, K. A., and Ingbar, S. H. (1968). The contribution of thyroxine-binding prealbumin to the binding of thyroxine in human serum, as assessed by immunoadsorption. *J Clin Invest.* 47(7): 1710-21.
- Wojtczak, A. (1997). Crystal structure of rat transthyretin at 2.5 Å resolution: first report on a unique tetrameric structure. *Acta Biochim Pol.* 44(3): 505-17.
- Wood, M.J. and Komives, E.A. (1999). Production of large quantities of isotopically labeled protein in *Pichia pastoris* by fermentation. *J Biomol NMR* 13, 149-159.
- Yamamoto, K., Hsu, S. P., Yoshida, K., Ikeda, S., Nakazato, M., Shiomi, K., Cheng, S. Y., Furihata, K., Ueno, I., and Yanagisawa, N. (1994). Familial amyloid polyneuropathy in Taiwan: identification of transthyretin variant (Leu55-->Pro). *Muscle Nerve.* 17(6): 637-41.
- Yoshinaka, T. Nakazato, M. Ikeda, S. and Ohnishi, A. (1992). Homozygosity for the transthyretin M-30 gene in three Japanese siblings with type I familial amyloidotic neuropathy. *Neurology.* 42(2): 2045-2047