

CONTENT

	Page
บทคัดย่อ.....	(3)
ABSTRACT.....	(5)
ACKNOWLEDGEMENT.....	(7)
CONTENT.....	(9)
LIST OF TABLES.....	(11)
LIST OF ILLUSTRATIONS.....	(13)
CHAPTER	
1. INTRODUCTION.....	1
2. REVIEW OF LITERATURES.....	4
2.1. COMPOSITIONS AND CHARACTERISTICS OF FATTY AND DARK-FLESH FISH	4
2.2. SURIMI PROCESSING FOR THE FATTY AND DARK-FLESH FISH	5
2.3. THE NOVEL SURIMI PROCESSING	9
2.4. HEMOGLOBIN	10
OBJECTIVES	22
EXPECTED OUTCOMES	22
SCOPE OF THE RESEARCH	23
3. EFFECT OF PH, ADP AND MUSCLE SOLUBLE COMPONENTS ON COD HEMOGLOBIN CHARACTERISTICS AND EXTRACTABILITY.....	24
3.1. INTRODUCTION	24
3.2. MATERIALS	25
3.3. METHODS	26
3.4. RESULTS AND DISCUSSION	32
3.5. CONCLUSION	43
3.6. REFERENCES	43

CONTENT (Continued)

	Page
4. EFFECT OF AN ALKALINE TREATMENT AND FISH MUSCLE COMPONENTS ON EXTRACTABILITY/SOLUBILITY OF HEMOGLOBIN.....	47
4.1. INTRODUCTION	47
4.2. MATERIALS	48
4.3. METHODS	49
4.4. RESULTS AND DISCUSSION	53
4.5. CONCLUSION	60
4.6. REFERENCES	60
5. REMOVAL OF HEME PROTEINS AND COLOR OF HERRING WASHED MINCE/PROTEIN ISOLATE BY WASHING OR THE ALKALINE SOLUBILIZATION PROCESS	63
5.1. INTRODUCTION	63
5.2. MATERIALS	64
5.3. METHODS	64
5.4. RESULTS AND DISCUSSION	67
5.5. CONCLUSION	76
5.6. REFERENCES	77
6. CONCLUSION.....	80
7. FUTURE WORKS.....	82
REFERENCES.....	85
APPENDIX.....	95
VITAE.....	100

LIST OF TABLES

	Page
Table 3-1 Per cent extractability of hemoglobin of washed and unwashed cod mince at pH 6.0 and 7.2.....	37
Table 3-2 Per cent extractable hemoglobin and soluble protein of the unwashed cod mince originally at pH 6.0 after extraction at pH 6.0 or 7.0 with or without added salt.....	38
Table 3-3 Effect of press juice and its components on the percentage of extractable hemoglobin of washed cod mince at pH 6.0.....	40
Table 3-4 Effect of press juice or its high and low molecular weight components on per cent soluble hemoglobin after incubation with SR at pH 6.0 for 30 min.....	41
Table 3-5 Per cent extractable hemoglobin of washed cod mince with added ADP.....	42
Table 4-1 Relative soluble hemoglobin in buffer system and relative extractable hemoglobin in cod homogenate after various pH treatments.....	54
Table 4-2 The relative percent soluble hemoglobin after various treatments.....	56
Table 4-3 Effect of the final pH after the alkaline treatment on percent soluble hemoglobin of cod and herring with added SR either from cod or herring.....	58
Table 4-4 Effect of pH on percent soluble myosin and percent soluble hemoglobin in presence and absence of myosin after alkaline treatment.....	59
Table 5-1 Extractable heme protein content of herring whole muscle at pH 7.0 and 8.0 with and without NaCl.....	69
Table 5-2 Soluble protein of herring whole muscle removed by different washing protocols.....	71
Table 5-3 Color values of washed herring whole muscle and protein isolate.....	74
Table 5-4 Soluble protein of herring whole muscle removed by different washing conditions.....	75
Table 1 Effect of the sarcoplasmic reticulum of cod or herring on the percentage of insoluble hemoglobin after incubation at pH 6.0 or 7.0 for 30 min.....	98
Table 2 Effect of hemoglobin addition at pH 10.8 on the per cent soluble hemoglobin remaining after the centrifugation at various precipitate pH values.....	98

LIST OF TABLES (Continued)

	Page
Table 3	Effect of existence of cod muscle soluble fraction (press juice) on per cent insoluble hemoglobin treated with the alkaline solubilization process.99
Table 4	Effect of freezing on extractable heme protein from light and whole muscle of herring99

LIST OF ILLUSTRATIONS

	Page
Figure 2-1 Protoporphyrin IX.....	11
Figure 2-2 The typical absorbance spectra of hemoglobin and its derivatives.....	13
Figure 2-3 Relative oxygenation of trout hemoglobin.....	14
Figure 3-1 Typical visible light absorbance spectrum of cod hemoglobin (10 μ M) in 50 mM phosphate buffer at pH 6.0 or 7.2.	33
Figure 3-2 Relative oxygenation of cod hemoglobin at various pH values.....	34
Figure 3-3 Effect of ADP and pH 7.0 on relative oxygenation of cod hemoglobin.....	35
Figure 3-4 Effect of extraction pH on percentage of extractable hemoglobin and protein of the unwashed mince originally at pH 6.0.	38
Figure 3-5 Effect of pH and sodium chloride on per cent soluble hemoglobin after incubation with and without cod SR.	39
Figure 4-1 Percent insoluble hemoglobin after incubation with SR at various pH values.....	55
Figure 5-1 Extractable heme protein content of herring light and whole muscle during iced storage.....	67
Figure 5-2 Amount of heme proteins in herring whole muscle removed by different methods.	71
Figure 5-3 Extractable heme protein of herring whole muscle removed by pre-washing and the alkaline solubilization process.....	73
Figure 1 Typical visible light absorbance spectrum of cod and herring hemoglobins in 50 mM phosphate at pH 8.0.	96
Figure 2 Relative oxygenation of cod and herring hemoglobin at various pH values...97	