

REFERENCES

- AOAC. 1999. Official Methods of Analysis. 16th Ed. Association of Official Analytical Chemists, Washington, DC.
- Abe, Y. 1994. Quality of kamaboko gel prepared from walleye pollock surimi with an additive containing transglutaminase. *Nippon Suisan Gakkaishi*. 60: 381-387.
- Akahane, Y. and Shimizu, Y. 1990. Effect of setting incubation on the water-holding capacity of salt-ground fish meat and its heated gel. *Nippon Suisan Gakkaishi*. 56: 139-143.
- Akahane, T., Chihara, S., Niki, T.P., Sano, T., Tsuchiya, T., Noguchi, S.F., Ookami, H. and Matsumoto, J.J. 1985. Different scanning calorimetric studies on thermal behaviours of myofibrillar proteins. *Bull. Jpn. Soc. Sci. Fish.* 51: 1841-1846.
- Akahane, T., Chihara, S., Yoshida, Y., Tsuchiya, T., Nogchi, S., Ookami, H. and Matsumoto, J.J. 1984. Roles of constituent proteins in gel properties of cooked meat gels. *Bull. Jap. Soc. Sci. Fish.* 50: 1029-1033.
- Albert, T. 1989. Stabilization energies of protein conformation. In *Prediction of protein structure and the principles of protein conformation* (Fasman, G.D., ed.). p. 161-192. Plenum. New York.
- Almas, K.A. 1981. Chemistry and Microbiology of Fish Processing. Dept. of Biochemistry. Norwegian Inst. of Technol. Univ. of Trondheim. Norway.
- Alvarez, C. and Tejada, M. 1997. Influence of texture of suwari gels on kamaboko gels made from sardine (*Sardina pilchardus*) surimi. *J. Sci. Food Agric.* 79: 472-480.
- Alvarez, C., Couso, I. and Tejada, M. 1995. Sardine surimi gels as affected by salt concentration, blending heat treatment and moisture. *J. Food Sci.* 60: 622-626.
- Alvarez, C., Couso, I. and Tejada, M. 1999. Microstructure of suwari and kamaboko sardine surimi gels. *J. Sci. Food Agric.* 79: 839-844.
- Amano, K., Yamada, K. and Bito, M. 1963. Detection and identification of formaldehyde in gadoid fish. *Bull. Jpn. Soc. Sci. Fish.* 29: 695-701.
- An, H., Peters, M.Y. and Seymour, T.A. 1996. Roles of endogenous enzymes in surimi gelation. *Trends. Food Sci. Technol.* 7: 321-327.

- An, H., Weerasinghe, V., Seymour, T.A. and Morrissey, M.T. 1994. Cathepsin degradation of Pacific whiting surimi proteins. *J. Food Sci.* 59: 1013-1017.
- Ando, H., Adachi, M., Unread, K., Matsuura, A., Nonaka M., Uchio, R., Tanaka, H. and Motoki, M. 1989. Purification and characteristics of novel transglutaminase derived from microorganisms. *Agric. Biol. Chem.* 53: 2613-2617.
- Ang, J.F. and Hultin, H.O. 1989. Denaturation of cod myosin during freezing after modification with formaldehyde. *J. Food Sci.* 54: 814-818.
- Arai, K., Kawamura, K. and Gayashi, C. 1973. The relative thermostability of actomyosin ATPase from the dorsal muscle of various fish species. *Bull. Jpn. Soc. Sci. Fish.* 39: 1077-1082.
- Araki, H. and Seki, N. 1993. Comparison of reactivity of transglutaminase to various fish actomyosins. *Nippon Suisan Gakkaishi.* 95: 711-716.
- Armenta-LÓpez, R., Guerrero, L.I. and Huerta, S. 2002. Astaxanthin extraction from shrimp waste by lactic fermentation and enzymatic hydrolysis of the carotenoprotein complex. *J. Food Sci.* 67. 1002-1006.
- Asagami, T., Ogiwara, M., Wakameda, A. and Noguchi, S. 1995. Effect of microbial transglutaminase on the quality of frozen surimi made from various kinds of fish species. *Fish. Sci.* 61: 267-272.
- Ashie, I.N.A. and Lanier, T.C. 1999. High pressure effects on gelation of surimi and turkey breast muscle enhanced by microbial transglutaminase. *J. Food Sci.* 64: 704-708.
- Ashie, I.N.A. and Lanier, T.C. 2000. Transglutaminase in seafood processing. In *Seafood enzymes utilization and influence on postharvest seafood quality* (Haard, N.F. and Simpson, B.K., eds.). p. 147-166. Marcel Dekker, Inc. New York.
- Ashie, I.N.A. and Simpson, B.K. 1997. Purification and characterization of cathepsin L from hepatopancreas of carp *Cyprinus carpio*. *Comp. Biochem. Physiol.* 115B: 531-537.
- Baker-Royer, E.N. and Dodson, E.J. 1992. Crystallographic refinement of the structure of actindin at 1.7 Å resolution by fast fourier least-squares methods. *Acta. Cryst. Sect. A*36: 559-572.
- Badii, F. and Howell, N.K. 2002. Changes in the texture of cod and haddock fillets during frozen storage. *Food Hydrocolloids.* 16: 313-319.

- Barely, T.H., Schapira, D.V. and Williams, G.T. 1983. The moisture and proximate composition of commercial fresh and frozen shrimp species as affected by phosphate treatments. *J. Food Sci.* 51: 878-881.
- Beas, V.E., Wagner, J.R., Crupkin, M. and Anon, M.C. 1990. Thermal denaturation of hake (*Merluccius hubbsi*) myofibrillar proteins. A differential scanning calorimetric and eletrophoretic study. *J. Food Sci.* 55: 683-696.
- Bechtel, J.P. 1986. Muscle development and contractile proteins. In *Muscle as Food* (Bechtel, J.P., ed.). p. 1-35. Academic Press. London.
- Benjakul, S. and Morrissey, M.T. 1997. Protein hydrolysates from Pacific whiting solid wastes. *J. Agric. Food Chem.* 45: 3423-3430.
- Benjakul, S., Seymour, T.A., Morrissey, M.T. and An, H. 1997. Physicochemical changes in Pacific whiting muscle proteins during ice storage. *J. Food Sci.* 62: 729-733.
- Benjakul, S. and Bauer, F. 2000. Physicochemical and enzymatic changes of cod muscle proteins subjected to different freeze-thaw cycles. *J. Sci. Food Agric.* 80: 1143-1150.
- Benjakul, S., Visessanguan, W. and Srivilai, C. 2001a. Gel properties of bigeye snapper (*Priacanthus tayenus*) surimi as affected by setting and porcine plasma protein. *J. Food Qual.* 24: 453-471.
- Benjakul, S., Visessanguan, W., Ishizaki, S. and Tanaka, M. 2001b. Differences in gelation characteristic of natural actomyosin from two species of bigeye snapper, *Priacanthus tayenus* and *P. macracanthus*. *J. Food Sci.* 66: 1311-1318.
- Benjakul, S., Visessanguan, W., Riebroy, S., Ishizaki, S. and Tanaka, M. 2002. Gel-forming properties of surimi produced from bigeye snapper, *Priacanthus tayenus* and *P. macracanthus*, stored in iced. *J. Sci. Food Agric.* 82: 1142-1451.
- Benjakul, S. and Visessangaun, W. 2003. Transglutaminase-mediated setting in bigeye snapper surimi. *Food Res. Int.* 36: 253-266.
- Benjakul, S., Visessanguan, W. and Tanaka, M. 2003a. Patial purification and characterization of trimethylamine-N-oxide demethylase from lizardfish kidney. *Comp. Biochem. Physiol.* 135B: 359-371.

- Benjakul, S., Visessanguan, W., Thongkaew, C. and Tanaka, M. 2003b. Comparative study on physicochemical changes of muscle proteins from some tropical fish during frozen storage. *Food Res. Int.* 36: 787-795.
- Benjakul, S., Visessanguan, W. and Tueksuban, J. 2003c. Changes in physico-chemical properties and gel-forming ability of lizardfish (*Saurida tumbil*) during post-mortem storage in ice. *J. Food Sci.* 80: 535-544.
- Benjakul, S., Chantarasuwan, C. and Visessangaun, W. 2003d. Effect of medium temperature setting on gelling characteristics of surimi from some tropical fish. *Food Chem.* 82: 567-574.
- Benjakul, S., Visessanguan, W. and Tanaka, M. 2004. Induced formation of dimethylamine and formaldehyde by lizardfish kidney trimethylamine-N-oxide demethylase. *Food Chem.* 84: 297-305.
- Bennour, M., Marrakchi, A.E. Bouchart, N., Hamama, A. and Ousdda, M. E. 1991. Chemical and microbiological assessments of mackerel (*Scomber scombrus*) stroed in ice. *J. Food Prot.* 54: 784-792.
- Bryant, C.M. and McClements, D.J. 1998. Molecular bases of protein functionality with special consideration of cold-set gels derived from heat-denatured whey. *Trends. Food Sci. Technol.* 9: 143-151.
- Buege, J.A. and Aust, S.D. 1978. Microsomal lipid peroxidation. *Methods Enzymol.* 52: 302-304.
- Busconi, L., Folco, E.J., Martone, C.B. and Sanchez, J.J. 1989. Postmortem changes in cytoskeletal elements of fish muscle. *J. Food Biochem.* 13: 443-451.
- Buttkus, H. 1974. On the nature of the chemical and physical bonds which contribute to some structural properties of protein foods: A hypothesis. *J. Food Sci.* 39: 484-489.
- Cao, M.J., Osatomi, K., Suzuki, M., Hara, K., Tachibana, K. and Ishihara, T. 2000. Purification and characterization of two anionic trypsins from the hepatopancreas of carp. *Fish. Sci.* 66: 1172-1179.
- Careche, M. and Li-Chan, E. 1997. Structural changes and raman spectroscopic studies of cod myosin upon modification with formaldehyde or frozen storage. *J. Food Sci.* 62: 717-723.

- Careche, M., Cofrades, S., Carballo, J. and Colmenero, F.J. 1998. Emulsifying and gelation properties during freezing and frozen storage of hake, pork, and chicken actomyosins as affected by addition of formaldehyde. *J. Agric. Food Chem.* 46: 813-819.
- Careche, M. and Tejada, M. 1990. Effect of neutral and oxidized lipids on functionality in megrim and sardine during frozen storage. *Food Chem.* 37: 275-287.
- Castell, L., Hedenreich, W. and Enrique, A. 1980. Frozen storage affects the compressive creep behavior of the porcine intervertebral disc. *J. Fish Sci.* 22: 2867-2876.
- Chalmers, M., Careche, M. and Mackie, I.M. 1992. Properties of actomyosin isolated from cod (*Gadus nitorhua*) after various periods of storage in ice. *J. Sci. Food Agric.* 58: 375-383.
- Chan, J.K. and Gill, T.A. 1994. Thermal aggregation of mixed fish myosins. *J. Agric. Food Chem.* 42: 2649-2655.
- Chan, J.K., Gill, T.A. and Paulson, A.T. 1992. Cross-link ability of myosin heavy chains from cod, herring and silver hake during thermal setting. *J. Food Sci.* 57: 906.
- Chan, J.K., Gill, T.A., Thompson, J.W. and Singer, D.S. 1995. Herring surimi during low temperature setting, physicochemical and textural properties. *J. Food Sci.* 60: 1248-1253.
- Cobb, B.F., Vanderzant, C. Thompson, C.A. and Custer, C.S. 1973. Chemical characteristics, bacterial counts and potential shelf-life of shrimp from various locations on the northwestern Gulf of Mexico. *J. Milk Food Technol.* 36: 463-468.
- Colmenero, F.J. and Borderias, A.J. 1983. A study of the effects of frozen storage of certain functional properties of meat and fish protein. *J. Food Technol.* 18: 731-737.
- Connell, J.J. 1960. Changes in the adenosinetriphosphatase activity and sulphhydryl groups of cod flesh during frozen storage. *J. Sci. Food Agric.* 11: 245-249.
- Connell, J. J. 1975. Control of fish quality. Surrey: Fishing News.
- Conway, E.J. and Byrne, A. 1936. An absorption apparatus for the micro-determination of certain volatile substances I. The micro determination of ammonia. *J. Biochem.* 27: 419-429.
- Damodaran, S. 1996. Functional properties. In *Food Proteins Properties and Characterization* (Nakai, S. and Modler, H. W., eds.). p. 167-233. VCH Publishers. New York.
- Davies, J.R., Bardsley, R.O. and Ledward, D.A. 1988. Myosin thermal stability in fish muscle. *J. Sci. Food Agric.* 45: 61-68.

- Decker, E.A., Xiong, Y.L., Calvert, J.T., Crum, A.D. and Blanchard, S.P. 1993. Chemical physical and functional properties of oxidized turkey white muscle myofibrillar proteins. *J. Agric. Food Chem.* 41: 186189.
- De Jong, G. and Koppelman, S. 2002. Transglutaminase catalyzed reactions: impact on food applications *J. Food Sci.* 67: 2798-2806.
- Dingle, J.R. and Hines, J.A. 1975. Protein instability in minced flesh from fillets and frames of several commercial Atlantic fishes during storage at -5 °C. *J. Fish. Res.bd. Can.* 32: 775-783.
- Ebashi, S. and Endo, M. 1968. Calcium and muscle contraction. *Prog. Biophys. Mol. Biol.* 18: 123-183.
- Ebashi, S., Kodama, A. and Ebashi, M. 1986. Troponin I. Preparation and physiological function. *J. Biochem.* 64 : 465-477.
- Eckhoff, M.K., Aidos, I., Hernre, G.I. and Lie, O. 1998. Collagen content in farmed Atlantic salmon (*Salmo salar*, L.) and subsequent changes in solubility during storage on ice. *Food Chem.* 62: 197-200.
- Ellman, G.L. 1959. Tissue sulfhydryl groups. *Archives of Biochemistry Biophysics.* 82: 70-77.
- Finch, C.A. 1983. Chemical modification and some cross-linking reactions of water soluble polymers. In *Chemistry and technology of water-soluble polymers* (Finch, C.A., ed.). p. 81-112. Plenum Press. New York.
- Finley, J.W. 1988. Effects of processing on proteins: an overview. In *Protein quality and the effects of processing* (Phillips, R.D. and Finley, J.W., eds.). p. 1-7. Marcel Dekker, Inc. New York.
- Fiske, C.H. and Subbarow, Y. 1925. The colorimetric determination of phosphorus. *J. Biol. Chem.* 66: 375-400.
- Foegeding, E.A., Lanier, T.C. and Hultin, H.O. 1996. Characteristics of edible muscle tissues. In *Food Chemistry* (Fennema, O.R., ed.). p. 879-942. Marcel Dekker, Inc. New York.
- Folk, J.E. 1970. Tranglutaminase. *Methods Enzymol.* 17: 889-894.
- Folk, J.E. 1980. Transglutaminase. *Ann. Rev. Biochem.* 49: 517-531.
- Folk, J.E. And Chung, S.I. 1985. Transglutaminase. *Methods Enzymol.* 113: 358-375.

- Folk, J.E. and Finlayson, J.S. 1977. The ϵ -(γ -glutamyl)lysine cross-link and the catalytic role of transglutaminase. *Adv. Protein Chem.* 31: 1-133.
- Gerber, U., Jucknischke, U., Putzien, S. and Fuchsbauer, H. L. 1994. A rapid and simple method for the purification of transglutaminase from *Streptoverticillium moharaense*. *J. Biochem.* 299: 825-829.
- Gill, T.A. and Conway, J.T. 1989. Thermal aggregation of cod (*Gadus morhua*) muscle proteins using 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide as a zero length cross-linker. *Agric. Biol. Chem.* 53: 2553-2555.
- Gill, T.A. and Paulson, A.T. 1982. Location, characterization and partial purification of TMAO-ase. *Comp. Biochem. Physiol.* 371B: 49-56.
- Gill, T.A., Chan, J.K., Phonchareon, K.F. and Paulson, A.T. 1992. Effect of salt concentration and temperature heat-induced aggregation and gelation of fish myosin. *Food Res. Int.* 25: 333-341.
- Gokodlu, N., Ozden, O. and Erkan, N. 1998. Physical, chemical and sensory analyses of freshly harvested sardines (*Sardina pilchardus*) stored at 4°C. *Can. J. Aquat. Food Prod. Technol.* 7: 5-15.
- Gorman, J.J. and Folk, J.E. 1981. Sturtural features of glutaminesubstrares for transglutaminase: specificity of human factor XIIIa and the guinea pig liver enzyme toward synthetic peptides. *J. Biochem.* 256: 2712-2715.
- Grabowska, J. and Sikorski, Z.E. 1976. The gel-forming capacity of fish myofibrillar proteins. *Lebensm. Wiss. U. Technol.* 9: 33-37.
- Gram, L. and Huss, H.H. 1996. Microbial spoilage of fish and fish products. *Inter. J. Food Microbiol.* 33: 121-137.
- Greenberg, C.S., Birckichler, P.J. and Rice, R.H. 1991. Transglutaminase: multifunctional cross-linking enzymes that stabilize tissues. *FASEB.* 5: 3071-3077.
- Guan, X., Yao, H., Chen, Z., Shan, L. and Zhang, M. 2006. Some functional properties of oat bran protein concentrate modified by trypsin. *Food Chem.* (in press)
- Haard, N.F., Simpson, B.K. and Sikorski, Z.E. 1994. Biotechnological applications of seafood proteins and other nitrogenous compounds Ch.13. *In Seafood Proteins*

- (Sikorski, Z.E., Pan, B.S. and Shihidi, F., eds.). p. 194-216. Chapman & Hall. New York.
- Hamada, I., Tsaji, K., Nakayama, T. and Niwa, E. 1977. Oxidative denaturation of actomyosin. Bull. Jpn. Soc. Sci. Fish. 43: 1105-1109.
- Hamann, D.D. 1992. Viscoelastic properties of surimi seafood products. In Viscoelastic Properties of Foods (Rao, M.A. and Steffe, J.F., eds.). p. 157-171. Marcel Dekker, Inc. New York.
- Harada, K. 1975. Studies on enzymes on enzymes catalyzing the formation of formaldehyde and dimethylamine in tissue of fishes and shellfishes. J. Shimonoseki University Fisheries. 18: 296-302.
- Harris, P. and Tall, J. 1994. Rancidity in fish. In Rancidity in Foods (Allen, J. C. and Hamiton, R.J., eds.). p. 256-272. Chapman & Hall. London.
- Hashimoto, A., Kobayashi, A. and Arai, A. 1982. Thermostability of fish myofibrillar Ca-ATPase and adaptation to environmental temperature. Bull. Jap. Soc. Sci. Fish. 48: 671-684.
- Hashimoto, K., Watabe, S., Kono, M. and Shiro, K. 1979. Muscle protein composition of sardine and mackerel. Bull. Jpn. Soc. Sci. Fish. 45: 1435-1441.
- Hastings, R.J., Rodger, G.W., Park, R., Matthews A.D. and Anderson, E.M. 1985. Differential scanning calorimetry of fish muscle: the effect of processing and species variation. J. Food Sci. 50: 503-510.
- Hatae, K., Tobimatsu, A., Takeyama, M. and Matsumoto, J.J. 1986. Contribution of the connective tissues on the texture difference of various fish species. Bull. Jpn. Soc. Sci. Fish. 52: 2001-2007.
- Hayakawa, S. and Nakai, S. 1985. Contribution of hydrophobicity, netcharge, and sulphydryl groups to thermal properties of ovalbumin. Can. Inst. Food Sci. Technol. J. 18: 290-295.
- Hayat, M.A. 1981. Fixation for electron microscopy. In Electron Microscopy in Biology (Harris, J.R., ed.). p.1-14. Oxford University Press. New York.
- Hebard, C.E., Flick, G.J. and Martin, R.E. 1982. Occurrence and significance of trimethylamine oxide and its derivatives in fish and shellfish. In Chemistry and Biochemistry of Marine

- Food Products (Martin, R.E., Flick, G.J. and Ward, D.R., eds.). p. 149-304. AVI Publishing Company. Connecticut, USA.
- Hermansson, A.M. 1979. Aggregation and denaturation involved in gel formation. In Functionality and Protein Structure (Pour-El, A., ed.). p. 81. American Chemical Society: Washington.
- Herrera, J.J.R, Pastoriza, L. and Samperdro, G. 2000. Inhibition of formaldehyde production in frozen-stored minced blue whiting (*Micromesistius poutassou*) muscle by cryostabilizers: an approach from the glassy state theory. *J. Agric. Food Chem.* 48: 5256-5262.
- Hill, A.R., Irvine, D.M. and Bullock, D.H. 1982. Precipitation and recovery of whey proteins. A review. *Can. Inst. J. Food Sci. Technol.* 15: 155-160.
- Ho, M.L., Leu, S.Z., Hsieh, J.F., and Jiang, S.T. 2000. Technical approach to simplify the purification method and characterization of microbial transglutaminase produced from *Streptoverticillium ladakanum*. *J. Food Sci.* 65: 76-80.
- Huang, Y. P., Seguro, K., Motoki, M. and Tawada, K. 1992. Cross-linking of contractile proteins from skeletal muscle by treatment with microbial transglutaminase. *J. Biochem.* 112: 229-234.
- Huidobro, A. and Tejada, M. 1994. Emulsifying capacity of fish mince from several species during frozen storage. *J. Sci. Food Agric.* 61: 333-338.
- Huidobro, A., Mohamed, G. F. and Tejada, M. 1998. Aggregation of myofibrillar proteins in hake, sardine and mixed minces during frozen storage. *J. Agric. Food Chem.* 46: 2601-2608.
- Huss, H.H. 1995. Quality and quality changes in fresh fish. FAO Fishery Technical Paper. 348: 1-195.
- Ikura, K., Kometani, T., Yoshikawa, M., Sasaki, R. and Chiba, H. 1980. Crosslinking of casein components by transglutaminase. *Agric. Biol. Chem.* 44: 1567-1573.
- Iso, N., Mizuno, H., Ogawa, H., Mochizuki, Y. and Masuda, N. 1991. Differential scanning calorimetry on fish meat paste. *Nippon Suisan Gakkaishi.* 57: 337-340.
- Ishikawa, S. 1978. Fish jelly product (kamaboko) and frozen minced meat (frozen surimi) made of sardine-II. Influence of the temperature during the manufacturing process on the jelly strength of kamaboko. *Bull. Tokai Reg. Fish Res. Lab.* 94: 37-44.
- Itoh, Y., Yoshinaka, R. and Ikeda, S. 1979. Behaviour of the sulphydryl reagents on the gel formation of carp actomyosin by heating. *Bull. Jap. Soc. Sci. Fish.* 45: 1019-1024.

- Iwata, K., Kobashi, K. and Hase, J. 1974. Studies on muscle alkaline protease III. Distribution of alkaline protease in muscle of fresh water fish, marine fish and in internal organs of carp. Bull. Jap. Soc. Sci. Fish. 40: 201-209.
- Jiang, S.T. 2000. Enzymes and their effects on seafood texture. In *Seafood Enzymes: utilization and influence on postharvest seafood quality* (Haard, N.F. and Simpson, B.K., eds.). p. 411-450. Marcel Dekker, Inc. New York.
- Jiang, S.T., Hsieh, J.F., Ho, M.L. and Chung, Y.C. 2000a. Microbial transglutaminase affects gel properties of golden threadfin-bream and Pollack surimi. J. Food Sci. 65: 694-699.
- Jiang, S.T., Hsieh, J.F., Ho, M.L. and Chung, Y.C. 2000b. Combination effects of microbial transglutaminase, reducing agent, and protease inhibitor on the quality of hairtail surimi. J. Food. Sci. 65: 241-245.
- Jiang, S.T., Leu, S.Z. and Tsai, G.J. 1998a. Cross-linking of mackerel surimi actomyosin by microbial transglutaminase and ultraviolet irradiation. J Agric. Food. Chem. 46: 5278-5282.
- Jiang, S., Hwang, D. and Chen, C. 1998b. Effect of storage temperatures on the formation of disulfides and denaturation of milkfish actomyosin (*Chanos chanos*). J. Food Sci. 53: 1333-1335.
- Jiang, S.T., Lan, C.C. and Tsao, C.Y. 1986. New approach to improve the quality of minced fish products from freeze-thawed cod and mackerel. J. Food Sci. 51: 310-312, 351.
- Jiang, S.T., Wang, F.J. and Chen, C.S. 1989. Properties of actin and stability of the actomyosin reconstituted from milkfish (*Chanos chanos*) actin and myosin. J. Agric. Food Chem. 37: 1232-1235.
- Jo, C. and Anh, D.U. 2000. Volatiles and oxidatives changes in irradiated pork sausage with different fatty acid composition and tocopherol content. J. Food Sci. 65: 270-275.
- Kamal, M., Watabe, S. and Hashimoto, K. 1991. Postmortem changes in myofibrillar ATPase of sardine ordinary and dark muscles. Nippon Suisan Gakkaishi. 57 : 1177-1184.
- Kamath, G.G., Lanier, T.C., Foegeding, E.A. and Hamann, D.D. 1992. Nondisulfide covalent cross-linking of myosin heavy chain in “setting” of Alaska pollock and Atlantic croaker surimi. J. Food Biochem. 16: 151-172.

- Kanaji, T., Ozaki, H., Takao, T., Kawajiri, H., Ide, H., Motoki, M. and Shimonishi, Y. 1993. Primary Structure of Microbial Transglutaminase from *Streptoverticillium* sp. Strain s-8112. *J. Biol. Chem.* 268: 11565-11572.
- Karakoltsidis, P.A., Zotos, A. and Constantinides, S.M. 1995. Composition of the commercially important Mediterranean finfish, crustaceans, and molluscs. *J. Food Comp. Anal.* 8: 258-273.
- Kato, A. and Nakai, S. 1980. Hydrophobicity determined by a fluorescence probe method and its correlation with surface properties of proteins. *Biochem. Biophys. Acta.* 624: 13-20.
- Kato, T., Tomimatsu, K. and Toba, S. 1991. Manufacture of storage-stable retort mapuo-doufu. Jap. Kokai Tokyo Koho JP. 03168059.
- Kielley, W.W. and Bradley, L.B. 1956. The relationship between sulphydryl groups and the activation of myosin adenosinetriphosphatase. *J. Biol. Chem.* 218: 653-659.
- Kimura, M., Seki, N. and Kimura, I. 2000. Occurrence and some properties of trimethylamine-N-oxide demethylase in myofibrillar fraction from walleye Pollack muscle. *Fish Sci.* 66: 755-729.
- Kimura, I., Sugimoto, M., Toyoda, K., Seki, N., Arai, K. I. and Fugita, T. 1991. A study on the cross-linking reaction of myosin in kamaboko "suwari" gels. *Nippon Suisan Gakkaishi.* 57: 1389-1396.
- Kolakowska, A. 2002. Lipid oxidation in food systems. In *Chemical and Functional Properties of Food Lipids* (Sikorski, Z.E. and Kolakowska, A., eds.). p. 133-160. CRC Press. USA.
- Krueger, D.J. and Fennema, O.R. 1989. Effect of chemical additives on toughening of fillets of frozen storage Alaska pollack (*Theragra chacogramma*). *J. Food Sci.* 54: 1101-1106.
- Kubow, S. 1992. Routes of formation and toxic consequences of lipid oxidation in foods. *Free Rad. Bio. Med.* 12: 63-81.
- Kumazawa, Y., Seguro, K., Takamura, M. and Motoki, M. 1993. Formation of ϵ -(γ -glutamyl) lysine cross-link in cured horse mackerel meat induced by drying. *J. Food Sci.* 58: 1062-1067.

- Kumazawa, Y., Numazawa, T., Seguro, K. and Motoki, M. 1995. Suppression of surimi gel setting by transglutaminase inhibitors. *J. Food Sci.* 60: 715-717.
- Kurokawa, T. 1979. Kamaboko-foaming ability of frozen and ice stored lizardfish. *Bull. Jap. Soc. Sci. Fish.* 45: 1551-1555.
- Kurth, L. and Rogers, P. 1984. Transglutaminase catalyzed cross-linking of myosin to soya protein, casein and gluten. *J. Food Sci.* 49: 573-589.
- Ladrat, C., Verrez-Bagnis, V., Noel, J. and Fleurence, J. 2003. *In vitro* proteolysis of myofibrilar and sarcoplasmic proteins of white muscle of sea bass (*Dicentrarchus labrax L.*): effects of cathepsins B, D and L. *Food Chem.* 81: 517-525.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of head of bacteriophage T4. *Nature.* 227: 680-685.
- Lanier, T.C. 1992. Measurement of surimi composition and functional properties. In *Surimi Technology* (Lanier, T.C. and Lee, C.M., eds.). p. 123-166. Marcel Dekker, Inc. New York.
- Lanier, T.C. 2000. Surimi gelation chemistry. In *Surimi and Surimi Seafood* (Park, J.W., ed.). p. 237-265. Marcel Dekker, Inc. New York.
- Lanier, T.C., Lin, T.S., Hamann, D.D. and Thomas, F.B. 1981. Effects of alkaline protease in mince fish on texture of heat-processed gels. *J. Food Sci.* 47: 1643-1649.
- Lawrence, R., Comsolacion, F. and Jelen, P. 1986. Formation of structured protein foods by freeze texturization. *Food Technol.* 40(4): 77-82.
- Lee, C.M. 1984. Surimi process technology. *Food Technol.* 38(11): 69-80.
- Lee, H.G., Lanair, T.C., Hamann, D.D. and Knopp, J.A. 1997. Transglutaminase effects on low temperature gelation of fish protein sols. *J. Food Sci.* 62: 20-24.
- Lee, N.H., Seki, N., Kato, N., Nakagawa, S., Terui, S. and Arai, K. 1990. Gel forming ability and cross-linking ability of myosin heavy chain in salted meat paste from threadfin bream. *Nippon Suisan Gakkaishi.* 56: 329-336.
- Leelapongwattana, K., Benjakul, S., Visessanguan, W. and Howell, N.K. 2005. Physicochemical and biochemical changes during frozen storage of minced flesh of lizardfish (*Saurida micropectorialis*). *Food Chem.* 90: 141-150.

- Li-Chen, E., Nakai, S. and Wood, D.F. 1985. Relationship between functional (fat-binding, emulsifying) and physicochemical properties of muscle proteins: Effect of heating, freezing, pH and species. *J. Food Sci.* 50: 1034-1040.
- Lin, T.M. and Park, J.W. 1996. Protein solubility in Pacific whiting by proteolysis during storage. *J. Food Sci.* 61: 536-539.
- Lorand, L., Urayama, T., Kiewiet, J.W.C. and Nossel, H.L. 1969. Diagnostic and genetic studies on fibrin-stabilizing factor with a new assay based on amine incorporation. *J. Clin. Invest.* 48: 1054-1064.
- Lorand, L. 1983. Post-translational pathways for generating ϵ -(γ -glutamyl) lysine crosslinks. *Ann. NY Acad. Sci.* 421: 10-27.
- Lowry, Q.H., Rosebrough, N.J., Farr, L.A. and Randall, R.J. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 256-275.
- Lundstrom, R.C., Correia, F.F. and Wilhelm, K.A. 1982a. Enzymatic dimethylamine and formaldehyde production in minced American plaice and blackback flounder mixed with a red TMAOase active fraction. *J. Food Sci.* 47: 1305-1310.
- Lundstrom, R.C., Correia, F.F. and Wilhelm, K.A. 1982b. Dimethylamine production in fresh red hake (*Urophycis chuss*): the effect of packaging material, oxygen permeability and cellular damage. *J. Food Biochem.* 6: 229-241.
- Ma, C.Y. 1985. Functional properties of oat concentrate treated with linoleate or trypsin. *Can. Inst. Food Sci. Technol.* 18: 79-84.
- MacDonald, G.A., Lelievre, J. and Wilson, N.D.C. 1990. The strength of gels prepared from washed and unwashed minces of hoki (*Macruronus novaezelandiae*) stored in ice. *J. Food Sci.* 55: 976-980.
- MacDonald, G.A., Lelievre, J. and Wilson, N.D.C. 1992. Effect of frozen storage on the gel-forming properties of hoki (*Macruronus novaezelandiae*). *J. Food Sci.* 57: 69-71.
- Mackie, I.M. 1994. Fish protein. In *New and Developing Sources of Food Proteins* (Hudson, B.F.J., ed.). p. 95-143. Chapman & Hall. New York.
- Mahmoud, M.I., Malone, W.T. and Cordle, C.T. 1992. Enzymatic hydrolysis of casein: effect of degree of hydrolysis on antigenicity and physical properties. *J. Food Sci.* 57: 51-65.

- Marrakchi, E., Bennour, A. M., Bouchriti, N., Hamama, A. and Tagafait, H. 1990. Sensory, chemical and microbiological assessments of Moroccan sardines (*Sardina pilchardus*) stored in ice. J. Food Prot. 53: 600-605.
- Martinez, I., Friis, T.J. and Careche, M. 2001. Post mortem muscle protein degradation during ice-storage of Arctic (*Pandalus borealis*) and tropical (*Penaeus japonicus* and *Penaeus monodon*) shrimps: a comparative eletrophoretic and immunological study. J. Sci. Food Agric. 81: 1199-1208.
- Matsumoto, J.J. 1979. Denaturation of fish muscle proteins during frozen storage. In Protein at low Temperatures (Fennema, O., ed.). p. 206. ACS. Adv. In Chem. Series. Washington, D.C. ACS.
- Matsumoto, J.J. 1980. Chemical deterioration of muscle proteins during frozen storage. In Chemical Deterioration of Protein (Whitaker, J.K. and Fujimaki, J.R., eds.). p. 95. ACS Symp. Series 123, ACS. Washington D.C.
- Mazorra-Manzano, M.A., Pacheco-Aguilar, R. Diaz-Rojas, E.I. and Lugo-Sanchez, M.E. 2000. Post-mortem changes in black skipjack muscle during storage on ice. J. Food Sci. 65: 774-779.
- McCormick, R.J. 1994. Structure and properties of tissue. In Muscle foods meat poultry and seafood technology (Kinsman, D.M., Kotula, A.W. and Breidenstein, B.C., eds.). p. 25-62. Chapman & Hall. New York.
- Morales, O.G., Ramirez, J.A., Vivanco, D.I. and Vazquez, D.I. 2001. Surimi of fish species from the gulf of Mexico: evaluation of the setting phenomenon. Food Chem. 75: 43-48.
- Morawetz, H. 1972. Conformation transitions in macromolecules. Adv. Protein. Chem. 26: 243.
- Morrisey, M.T., Wu, J.W., Lin, D.D. and An, H. 1993. Protease inhibitor effects on torsion measurements and autolysis of Pacific whiting of surimi. J. Food Sci. 58: 1050-1054.
- Motoki, M. and Seguro, K. 1998. Transglutaminase and its use for food processing. Trends. Food Sci. Technol. 9: 204-210.
- Muguruma, M., Sakamoto, K., Numata, M., Yamada, H. and Nakamura, T. 1990. Studies on application of transglutaminase to meat and meat products, II. Effect of microbial transglutaminase on gelation of myosin B, myosin and actin. Nippon Suisan Gakkaishi. 56: 446-453.

- Multilangi, W.A.M., Panyam, D. and Kilara, A. 1996. Functional properties of hydrolysates from proteolysis of heat-denatured whey protein isolate. *J. Food Sci.* 61: 270-274.
- Nakahara, C., Nozawa, H. and Seki, N. 1999. A comparison of cross-linking of fish myofibrillar proteins by endogenous and microbial transglutaminase. *Fish. Sci.* 65: 138-144.
- Nambudiri, D.D. and Gopakumar, K. 1992. ATPase and lactate dehydrogenase activities in frozen stored fish muscle as indices of cold storage deterioration. *J. Food Sci.* 57 : 72-76.
- Nawar, W.W. 1996. Lipids. In *Food Chemistry* (Fennema, O.R., ed.). p. 225-314. Marcel Dekker, Inc. New York.
- Ng, C.S. 1978. Measurement of free and expressible drips. In *Laboratory Manual on Analytical Methods and Procedure for Fish and Fish Products*. (Hasegawa, H., ed.) p. 1-2. Southeast Asian Fisheries Development Center. Singapore.
- Nip, W.K. and Moy, J.H. 1988. Microstructural changes of ice-chilled and cooked freshwater prawn, *Macrobrachium rosenbergii*. *J. Food Sci.* 53: 319-322.
- Nishimoto, S., Hashimoto, A., Seki, N., Kimura, I., Toyoda, K. and Arai, K. 1987. Influencing factors on changes in myosin heavy chain jelly strength of salted meat paste from Alaska pollack during setting. *Nippon Suisan Gakkaishi*, 56: 1341.
- Niwa, E. 1992. Chemistry of surimi gelation. In *Surimi Technology* (Lanier, T.C. and Lee, C.M. eds.). p. 389-428. Marcel Dekker, Inc. New York.
- Niwa, E. 1975. Role of hydrophobic bonding in gelation of fish flesh paste. *Bull. Jap. Soc. Sci. Fish.* 41: 907-908.
- Niwa, E., Nakayama, T. and Hamada, I. 1982. Participation of SS bonding in the appearance of setting. *Bull. Jap. Soc. Sci. Fish.* 48: 727-724.
- Niwa, E., Sato, J., Suzuki, R., Nakayama, T. and Hamada, I. 1981. Fluorometric study of setting properties of fish flesh sol. *Bull. Jap. Soc. Sci. Fish.* 47: 817-819.
- Nonaka, M., Matsura, Y., Nakano, K. and Motoki, M. 1997. Improvement of the pH-solubility profile of sodium caseinate by using Ca^{2+} -independent microbial transglutaminase with gelatin. *Food Hydrocolloids.* 11: 347-349.

- Nonaka, M., Tanaka, H., Okiyama, A., Motoki, M., Anso, H., Umeda, K. and Matsukuwa, A. 1989. Polymerization of several proteins by Ca^{2+} independent transglutaminase derived from microorganisms. *Agric. Biol. Chem.* 36: 1078-1081.
- Numakura, T., Kimura, I., Toyoda, K. and Fujita, T. 1990. Temperature dependent changes in gel strength and myosin heavy chain of salt ground meat from walleye pollack during setting. *Nippon Suisan Gakkaishi.* 56: 329-336.
- Ochiai, Y. and Chow, C.J. 2000. Myosin ATPase. In *Seafood enzymes* (Haard, N.F. and Simpson, B.K., eds.). p. 69-90. Marcel Dekker, Inc. New York.
- Ogawa, M., Ehara, T., Tamiya, T. and Tsuchiya, T. 1993. Thermal stability of fish myosin. *Comp. Biochem. Physiol.* 106B: 517-521.
- Ogawa, M., Tamiya, T. and Tsuchiya, T. 1994. Structural changes of carp myosin during heating. *Fish. Sci.* 60: 723-727.
- Ohba, Y., Morimitsu, Y. and Watarai, A. 1979. Reaction of formaldehyde with Calf-thymus nucleohistone. *Eur. J. Biochem.* 100: 285-293.
- Okada, M. 1992. History of surimi technology in Japan. In *Surimi Technology* (Lanier, T.C. and Lee, C.M., eds.). p. 3-22. Marcel Dekker, Inc. New York.
- Okada, S., Nur-E-Borhan, S.A. and Yamaguchi, K. 1998. Carotenoid composition in the exoskeleton of commercial black tiger prawns. *Fish. Sci.* 60: 213-215.
- Ouali, A. and Valin, C. 1981. Effect of muscle lysosomal enzyme and calcium activated neutral proteinase on myofibrillar ATPase activity with aging change. *Meat Sci.* 5: 233-245.
- Owusu-Ansah, Y.J. and Hultin, H.O. 1987. Effect of *in situ* formaldehyde production on solubility and cross-linking of proteins of minced red hake muscle during frozen storage. *J. Food Biochem.* 11: 17-34.
- Pacheco-Aguilar, R., Lugo-Sanchez, M.E. and Robles-Burgueno, M.R. 2000. Postmortem biochemical characteristic of Monterey sardine muscle stored at 0°C . *J. Food. Sci.* 65: 40-47.
- Panyam, D. and Kilara, A. 1996. Enhancing the functionality of food proteins by enzymatic modification. *Trends. Food Sci. Technol.* 7: 120-125.
- Park, J.W. 1994. Functional protein additives in surimi gel. *J. Food Sci.* 59: 525-527.

- Park, J.W. and Lanier, T.C. 1989. Scanning calorimetric behavior of tilapia myosin and actin due to processing of muscle and protein denaturation. *J. Food Sci.* 54: 49-51.
- Pearsons, A.M. and Young, R.B. 1989. Muscle and meat biochemistry. Academic Press, Inc. San Diego. CA.
- Perez-Villarreal, B. and Pozo, R. 1990. Chemical composition and iced spoilage of albacore (*Thunnus alalunga*). *J. Food Sci.* 55: 535-544.
- Pigott, G.M. and Tucker, B.W. 1990. Seafood: Effect of Technology on Nutrition. p.362. Marcel Dekker, Inc. New York.
- Pomeranz, Y. 1991. Proteins: General. In *Functional Properties of Food Components* (Pomeranz, Y., ed.). p. 147-189. Academic Press. Sandigo.
- Poulter, R.G. Ledward, D.A., Godber, S., Hall, G. and Rowlands, B. 1985. Heat stability of fish muscle proteins. *J. Food Technol.* 20: 203-217.
- Racicot, L.D., Lundstrom, R.C., Wihelm, K.A., Ravesi, E.M. and Licciardello, J.J. 1984. Effect of oxidizing and reducing agents on trimethylamine N-oxide demethylase activity in red hake muscle. *J. Agric. Food Chem.* 32: 459-464.
- Ragnarrsson, K. and Regenstien, J.M. 1989. Changes in electrophoretic patterns of gadoid and non-gadoid muscle during frozen storage. *J. Food Sci.* 54: 819-823.
- Rahaman, M., Yasmin, L., Kamal, M.D., Mazid, M.A. and Islam, N. M.D. 2001. Effect of delayed icing on the quality changes in brackish water shrimp *Penaeus monodon* during ice storage. *J. Biol. Sci.* 11: 1390-1394.
- Ramirez-Suarez, J.C. and Xiong, Y.L. 2002. Transglutaminase crosslinking of whey/myofibrillar proteins and the effect on protein gelation. *J. Food Sci.* 67: 2885-2891.
- Ramirez-Suarez, J.C. and Xiong, Y.L. 2003. Effect of transglutaminase-induced cross-linking on gelation of myofibrillar/soy protein mixtures. *Meat Sci.* 65: 899-907.
- Ramirez-Suarez, J.C., Xiong, Y.L. and Wang, B. 2001. Transglutaminase cross-linking of bovine cardiac myofibrillar proteins and its effect on protein gelation. *J. Muscle Food.* 12: 85-96.
- Ramirez, J.A., Rodriguez-Sosa, R., Morales, O.G. and Vazquez, M. 2000. Surimi gels from striped mullet (*Mugil cephalus*) employing microbial transglutaminase. *Food Chem.* 70: 443-449.

- Regenstein, J.E., Jauregui, C.A. and Baker, R.C. 1983. The effect of pH, polyphosphates and different salts on water retention properties of ground trout muscle. *J. Food Biochem.* 8: 123-131.
- Rehbein, H. and Schreiber, W. 1984. TMAO-ase activity in tissue of fish species from the northeast Atlantic. *Comp. Biochem. Physiol.* 79B.: 447-452.
- Riaz, F. and Qadri, R. B. 1985. Quality changes in lobster (*Panulirus poliphagus*) muscle during storage in ice. *J. Agric. Food. Chem.* 33: 117-122.
- Riebroy, S., Benjakul, S., Visessanguan, W and Tanaka, M. 2006. Effect of iced storage of bigeye snapper (*Priacanthus tayenus*) on the chemical composition, properties and acceptability of Som-fug, a fermented Thai fish mince. *Food Chem.* (in press).
- Robinson, H.W. and Hodgen, C.G. 1940. The biuret reaction in the determination of serum protein. I. A study of condition necessary for the production of the stable color which bears a quantitative relationship to the protein concentration. *J. Biol. Chem.* 135: 707-725.
- Roura, S.I. and Crupkin, M. 1995. Biochemical and functional properties of myofibrils from pre- and post-spawned hake (*Merluccius hubbsi Marini*) stored in ice. *J. Food Sci.* 60: 269-272.
- Roura, S.I., Monteccia, C., Goldemberg, A.L., Trucco, R.E., and Crupkin, M. 1990. Biochemical and physicochemical properties of actomyosin from pre and post-spawned hake (*Merluccius hubbsi Marini*) stored on ice. *J. Food Sci.* 55 : 688-692.
- Roussel, H. and Cheftel, J.C. 1990. Mechanism of gelation of sardine protein: influence of thermal processing and various additives on the texture and protein solubility of kamaboko gel. *Int. J. Food Sci. Technol.* 25: 260-280.
- Rowland, B., Finne, G. and Tillman, R. 1982. A morphological study of muscle proteolysis in the tails of *Macrobrachium rosenbergii*. *Proc. 7th Annual Trop. Subtrop. Fish. Technol. Conf. Amer. TAMU-SG-82-10.* p. 105.
- Ryder, J.M., Buisson, D.H., Scott, D.N. and Fletcher, G.C. 1984. Storage of New Zealand jack mackerel (*Trachurus novazelandiae*) in ice: chemical, microbiological and sensory assessment. *J. Food Sci.* 49: 1453-1456.

- Saeed, S., Fawthrop, S. A. and Howell, N. K. 1999. Electron spin resonance (ESR) study on free-radical transfer in fish lipid-protein interactions. *J. Sci. Food Agric.* 79: 1809-1816.
- Saeki, H., Iseya, Z., Sugiura, S. and Seki, N. 1995. Gel forming characteristics of frozen surimi from chum salmon in the presence of proterase inhibitors. *J. Food Sci.* 60: 917-928.
- Sakamoto, H., Kumazawa, Y. and Motoki, M. 1994. Strength of protein gels prepared with microbial transglutaminase as related to reaction conditions. *J. Food Sci.* 59: 866-871.
- Sakamoto, H., Kumazawa, Y., Toiguchi, S., Seguro, K., Soeda, T. and Motoki, M. 1995. Gel strength enhancement by addition of microbial transglutaminase during onshore surimi manufacture. *J. Food Sci.* 60: 300-304.
- Samejima, K., Ishioroshi, M. and Yasui, T. 1981. Relative roles of the head and tail portions of the molecule in heat induced gelation of myosin. *J. Food Sci.* 46: 1412-1416.
- Sano, T., Noguchi, S.F., Matsumoto, J.J. and Tsuchiya, T. 1990. Thermal gelation characteristics of myosin subfragments. *J. Food Sci.* 55: 55-58.
- Sano, T., Ohno, T., Fuchino, H.O., Matsumoto, J. and Tsuchiya, T. 1994. Carp natural actomyosin: thermal denaturation mechanism. *J. Food Sci.* 59: 1002-1008.
- Sasajima, M. 1973. Studies on the psychrotolerant bacteria in fish and shellfish. IV. Relation between the number of trimethylamine oxide reducing psychrotrophic bacteria and their activity. *Bull. Jap. Soc. Sci. Fish.* 39: 511-518.
- Sato, K., Ohashi, C., Ohtsuki, K. and Kawabata, M. 1991. Type V collagen in trout (*Salmo gairdneri*) muscle and its solubility change during chilled storage of muscle. *J. Agric. Food Chem.* 39: 1222-1225.
- Seguro, K., Kumazawa, Y., Ohtsuka, T., Toiguchi, S. and Motoki, M. 1995. Microbial transglutaminase ϵ -(γ -glutamyl)lysine crosslink effects on elastics properties of kamaboko gels. *J. Food Sci.* 60: 305-311.
- Seguro, K., Nio, N. and Motoki, M. 1996. Some characteristics of a Microbial protein Cross-linking Enzyme: Transglutaminase in Macromolecular. *Interactions in Food Technology (ACS Symposium Series 650)*. p. 271-280. American Chemical Society.
- Seki, N., Ikeda, M. and Narita, N. 1979. Changes in ATPase activities of carp myofibril during ice-storage. *Bull. Jap. Soc. Sci. Fish.* 45 791-799.

- Seki, N. and Narita, N. 1980. Changes in ATPase activities and other properties of carp myofibrillar proteins during iced storage. *Bull. Jap. Soc. Sci. Fish.* 46: 207-213.
- Seki, N., Uno, H., Lee, N.H., Kimura, I., Toyoda, K., Fujita, T. and Arai, K. 1990. Transglutaminase activity in Alaska pollack muscle and surimi, and its reaction with myosin B. *Nippon Suisan Gakkaishi.* 56: 125-132.
- Seki, N., Nakahara, C., Takeda, H., Maruyama, N. and Nozawa, H. 1998. Dimerization site of carp myosin heavy chains by the endogenous transglutaminase. *Fish. Sci.* 64: 314-319.
- Sekine, T., Barnett, L.M. and Kielley, W.W. 1962. The active site of myosin adenosine triphosphatase. *J. Biol. Chem.* 237: 2769-2772.
- Shenouda, S.Y.K. 1980. Theories of protein denaturation during frozen storage of fish flesh. *Adv. Food Res.* 26: 275-311.
- Shimizu, Y. 1985. Biochemical and functional properties of material fish. In Proceedings of the International Symposium on Engineered Seafood Including Surimi (Seattle, W.A., Martin, R.E., and Collete, R.L., eds.). Nov 19-21. p. 148-167. National Fish. Institute. Washington DC.
- Shimizu, Y., Machida, R. and Takenami, S. 1981. Species variations in the gel-forming characteristics of fish meat paste. *Nippon Suisan Gakkaishi.* 47: 95-104.
- Sikorski, Z.E. 1994. The contents of proteins and other nitrogenous compounds in marine animals. In *Seafood Protein* (Sikorski, Z.E., Pan B.S. and Shahidi, F., eds.). p. 6-12. Chapman & Hall. New York.
- Sikorski, Z.E. and Borderias, J.A. 1990. Collagen in the muscles and skin of marine animals. In *Seafood: Resources, Nutritional Composition and Preservation* (Sikorski, Z.E., ed.). p. 58-70. CRC Press, Inc. Florida.
- Sikorski, Z.E., Kolakoska, A. and Pan, B.S. 1990a. The nutritive composition of the major groups of marine food organisms. In *Seafood: Resources, Nutritional Composition and Preservation* (Sikorski, Z.E., ed.). p. 29-54. CRC Press, Inc. Florida.
- Sikorski, Z., Kolakowska, A. and Burt, J. R. 1990b. Postharvest biochemical and microbial changes. In *Seafood: Resources, nutritional composition and preservation* (Sikorski, Z.E., ed.) p. 55-57. CRC Press, Inc. Florida.

- Sikorski, Z.E., Olley, J. and Kostuch, S. 1976. Protein changes in frozen fish. CRC. Crit. Rev. Food Sci. Nutr. 8: 97-129.
- Simpson, K.L. 1982. Carotenoid pigments in seafood. In Chemistry and biochemistry of marine food products R.E. Martin (Flick, G.J. and Ward, D.R., eds.). p. 115-148. AVI Publishing. Connecticut.
- Smith, D.M. 1991. Factors influencing heat induced gelation of muscle proteins. In Interactions of Food Proteins (Paris, N. and Bradford, R., eds.). p. 243-256. American Chemical Society. Washington DC.
- Smith, M.S. 1988. Meat proteins: functional properties in comminuted meat products. Food Technol. 21: 116-121.
- Smyth, A.B., Smith, D.M. and Ó-Neill, E. 1998. Disulfide bonds influence the heat-induced gel properties of chicken breast muscle myosin. J. Food. Sci. 63: 584-589.
- Sompongse, W., Itoh, Y. and Obatake, A. 1996. Role of SH_a in the polymerization of myosin heavy chain during iced storage of carp actomyosin. Fish. Sci. 62 : 110-113.
- Sotelo, C.G., Pinerio, C. and Perez-Martin, R.I. 1995. Denaturation of fish proteins during frozen storage: Role of formaldehyde. Z-Zebensm. Unters Forsch. 200: 14-23.
- Srinivasan, S. and Hultin, H.O. 1997. Chemical, physical and functional properties of cod proteins modified by a nonenzymic freeradical-generating system. J. Agric. Food Chem. 45: 310-320.
- Stabursvik, E. and Martens, H. 1980. Thermal denaturation of proteins in post rigor muscle tissue as studies by differential scanning colorimetry. J. Sci. Food Agric. 31: 1934-1042.
- Stamman, K., Gerdes, D. and Caporaso, F. 1990. Modified atmosphere packaging of seafood. Food Sci. Nutr. 29: 301-331.
- Steel, R.D.D. and Torrie, J.H. 1980. Principle and procedure of statistics: A biomaterial approach (2nd ed.) p. 862. MacGraw-Hill. New York.
- Suzuki, T. 1981. Fish and Krill Protein. In Protein Technology. Appl. Sci. Publ, Ltd. London. UK.
- Tachibana, K., Misima, T., and Tsuchimoto, M. 1993. Changes of ultrastructure and cytochemical Mg²⁺-ATPase activity in ordinary muscle of cultured and wild red sea bream during storage in ice. Nippon Suisan Gakkaishi. 54: 721-727.

- Taguchi, T., Ishizaka, H., Tanaka, M., Nagashima, Y. and Amano, K. 1987. Protein-protein interaction of fish myosin fragments. *J. Food Sci.* 52: 1130-1136.
- Tanabe, M. and Saeki, H. 2001. Effect of Maillard reaction with glucose and ribose on solubility at low ionic strength and filament-forming ability of fish myosin. *J. Agric. Food Chem.* 49: 3403-3407.
- Tejada, M., Mohamed, G.F. and Huidobro, A. 2002. Addition of sardine to hake minces and subsequent effect on dimethylamine and formaldehyde formation. *J. Sci. Food Agric.* 82: 351-359.
- Thannhauser, T.W., Konishi, Y. and Scheraga, H.A. 1987. Analysis for disulfide bonds in peptides and protein. *Methods Enzymol.* 143: 115-119.
- Tokiwa, T. and Matsumiya, J. 1969. Fragmentation of fish myofibril. Effect of storage condition and muscle cathepsin. *Bull. Jap. Soc. Sci. Fish.* 35: 1099-1109.
- Tome, D., Kozlowski, A. and Mahon, F. 1985. Carbon-13 NMR study on the combination of formaldehyde with bovine serum albumin. *J. Agric. Food Chem.* 33: 449-457.
- Tsai, G.J., Lin, S.M. and Jaing, S.T. 1996. Transglutaminase from *Streptoverticillum ladakanum* and application to mince fish product. *J. Food Sci.* 61: 1234-1238.
- Tsai, S.F., Wang, J.H. and Jiang, S.T. 1989. Stability of muscle proteins from some subtropical fish. *J. Agric. Food Chem.* 37: 749-752.
- Tsai, G.J., Tsai, F.C. and Jiang, S.T. 1995. Cross-linking of mackerel muscle proteins by microbial transglutaminase. *J. Fish. Soc. Taiwan.* 22: 125-135.
- Tsukamasa, Y. and Shimizu, Y. 1990. Setting property of sardine and Pacific mackerel meat. *Nippon Suisan Gakkaishi.* 56: 1105-1112.
- Tsukamasa, Y., Sato, K., Shimizu, Y., Imai, C., Sugiyama, M., Minegishi, Y. and Kaqabata, M. 1993. ϵ -(γ -glutamyl) lysine crosslink formation in sardine myofibril sol during setting at 25 °C. *J. Food Sci.* 59: 785-787.
- Venugopal, V., Alur, M.D. and Lewis, N.F. 1983. Extracellular protease from *Pseudomonas marinoglutinosa*: some properties and its action on fish actomyosin. *J. Food Sci.* 48: 671-702.
- Vojdani, F. 1996. Solubility. In *Methods of Testing Protein Functionality* (Hall, G. M., ed.). p. 11-60. Blackie Academic and Professional. New York.

- Wan, J., Kimura, I., Satake, M. and Seki, N. 1994. Effect of calcium ion concentration on the gelling properties and transglutaminase activity of walleye pollack surimi paste. *Fish. Sci.* 60: 107-114.
- Wan, J., Kimura, I. and Seki, N. 1995. Inhibitory factors of transglutaminase in salted salmon meat paste. *Fish Sci.* 61: 968-972.
- Watabe, S., Ushino, H., Iwamoto, M., Yamanako, H. and Hashimoto, K. 1989. Temperature-dependency of rigor-mortis of fish muscle Myofibrillar Mg^{2+} -ATPase activity and Ca^{2+} uptake by sarcoplasmic reticulum. *J. Food Sci.* 54 : 1107-1115.
- Wicker, L., Lanier, T.C., Hamann, D.D. and Akahane, T. 1986. Thermal transitions in myosin-ANS fluorescence and gel rigidity. *J. Food Sci.* 51: 1540-1543.
- Wright, D.J. and Wilding, P. 1984. Differential scanning calorimetric study of muscle and its proteins: myosin and subfragments. *J. Sci. Food Agric.* 35: 357-372.
- Wong, S.S. 1991. Zero-length cross-linking reagents. In *Chemistry of Protein Conjugation and Cross-linking* (Wong, S.S., ed.). p. 195-220. CRC Press. Boca Raton:
- Wu, H.C., Shiau, C.Y., Chen, H.M. and Chiou, T.K. 2003. Antioxidant activities of carnosine, anserine and some free amino acids and their synergistic effects in combination. *J. Food. Drug Anal.* 11: 148-153.
- Xiong, Y.L. 1997. Structure-function relationships of muscle protein. In *Food Protein and Their Applications* (Damodaran, S. and Paraf, A., eds.). p. 341-392. Marcel Dekker, Inc. New York.
- Xiong, Y.L. and Brekke, C.J. 1989. Changes in protein solubility and gelation properties of chicken myofibrils during storage. *J. Food Sci.* 54: 1141-1146.
- Xiong, Y.L. and Decker, E.A. 1995. Alterations of muscle protein functionality by oxidative and antioxidative processed. *J. Muscle Foods.* 6: 139-160.
- Xiong, Y.L., Lou, X., Wang, C., Moody, W.G. and Harmon, R.J. 2000. Protein extraction from chicken myofibrils irrigated with various polyphosphate and NaCl solutions. *J. Food Sci.* 65: 69-100.
- Yamagushi, M. and Sekine, T. 1966. Sulhydryl groups involved in the active site of myosin. A adenosine triphosphate. I. Specific blocking of the SH group responsible for the inhibitory phase in "B : phasic response" of the catalytic activity. *J. Biochem.* 59: 24-33.

- Yamashita, M. and Konagaya, S. 1991. Hydrolytic action of salmon cathepsin B and L to muscle structural proteins in respect of muscle softening. Nippon Suisan Gakkaishi. 57: 1917-1922.
- Yamashita, K., Arai, K. and Nishita, K. 1978. Thermo-stabilites of synthetic actomyosins in various combinations of myosin and actin from scallop and rabbit muscles. Bull. Jap. Soc. Sci. Fish. 44: 485-489.
- Yamazawa, M. 1990. Effect of high temperature heating on physical properties of kamaboko gel. Nippon Suisan Gakkaishi. 56: 497-503.
- Yanar, Y. and Celik, M. 2005. Seasonal variations of fatty acid composition in wild marine shrimps (*Penaeus semisulcatus* De Hann, 1844 *Metapenaeus monoceros* Fabricius, 1789) from the Eastern Mediterranean Sea. Food Sci. Technol. Intern. 11: 391-395.
- Yasunaga, K., Abe, Y., Yamazawa, M. and Arai, K. 1996. Heat-induced change in myosin heavy chains in salt-ground meat with a food additive containing transglutaminase. Nippon Suisan Gakkaishi. 62: 659-668.
- Zeigler, G.R. and Acton, J.C. 1984. Machanism of gel forming by proteins of muscle tissue. Food Technol. 38: 77-82.