

CONTENTS

	Page
Contents	viii
List of tables	xv
List of figures	xvii
Chapter	
1. Introduction	1
Literature review	2
1. Chemical composition of fish and shell fish	2
1.1 Myofibrillar protein	3
- Myosin and paramyosin	4
- Actin	5
- Tropomyosin	6
- Troponin	6
1.2 Sarcoplasmic protein	7
1.3 Stroma protein	8
2. Post mortem changes of fish	8
2.1 Changes of proteins	8
2.1.1 Degradation of proteins	8
2.1.2 Denaturation of proteins	9
2.1.3 Cross-linking of proteins	11
3. Functional properties of fish protein	12
3.1 Water holding capacity	12
3.2 Solubility	13
3.3 Gel-forming ability	13
3.3.1 Protein denaturation	14
3.3.2 Aggregation	15
4. Suwari (setting)	18
5. Transglutaminase (TGase)	20

CONTENTS (Continued)

	Page
6. Microbial transglutaminase	22
Objectives	27
2. Materials and Methods	28
1. Materials	28
1.1 Fish samples	28
1.2 Chemicals	28
2. Instruments	29
3. Methods	29
3.1 Determination of chemical composition and property of white shrimp, bigeye snapper and lizardfish mince	29
3.1.1 Proximate analysis	29
3.1.2 Determination of nitrogenous constituent	29
3.1.3 Thermal denaturation	30
3.2 Study on the effect of setting condition and MTGase on gel forming ability of some marine fish and shrimp meats	30
3.2.1 Determination of MTGase activity	30
3.2.2 Effect of setting condition and MTGase on gel forming ability	30
3.2.2.1 Determination of breaking force and deformation	32
3.2.2.2 Determination of whiteness	32
3.2.2.3 Determination of protein patterns	32
3.2.2.4 Determination of expressible moisture content	33
3.2.2.5 Determination of microstructure	33
3.3 Effect of post-mortem storage on crossing and gel enhancing ability MTGase on mince from some marine fish and shrimp	33
3.3.1 Preparation of ice-stored some marine fish and shrimp	33
3.3.2 Analyses	34

CONTENTS (Continued)

	Page
3.3.2.1 Determination of TCA-soluble peptide content	34
3.3.2.2 Determination of Thiobarbituric acid reactive substances (TBARS)	34
3.3.2.3 Determination of pH	34
3.3.2.4 Determination of total volatile bases (TVB) and trimethylamine (TMA) contents	34
3.3.2.5 Determination of formaldehyde content	35
3.3.2.6 Determination of protein patterns	35
3.3.2.7 ATPase activity	35
3.3.3 Effect of MTGase on gel-forming ability of some marine fish and shrimp stored in ice for different times	36
3.4 Effect of protein substrate denaturation on cross-linking activity of MTGase	36
3.4.1 Preparation of natural actomyosin (NAM)	36
3.4.2 Thermal stability of NAM	36
3.4.3 Effect of MTGase on cross-linking and physico-chemical properties of natural actomyosin with different degrees of denaturation	37
3.4.3.1 Determination of protein solubility	37
3.4.3.2 Determination of Surface hydrophobicity	37
3.4.3.3 Determination of Total sulfhydryl group content	37
3.4.3.4 Determination of Disulfide bond content	38
3.4.3.5 Determination of protein patterns	38
3.4.3.6 Determination of free amino group content	38
3.5 Effect of MTGase on cross-linking and physicochemical properties of natural actomyosin added with different levels of formaldehyde	38

CONTENTS (Continued)

	Page
3.6 Effect of MTGase on cross-linking and physicochemical properties of natural actomyosin with different degrees of hydrolysis	39
3.6.1 Preparation of NAM with different degrees of hydrolysis	39
3.6.1.1 Determination of free amino group content	39
3.6.1.2 Determination of protein patterns	39
3.6.1.3 Determination of Surface hydrophobicity	39
3.6.2 Effect of MTGase on cross-linking and physico-chemical properties of natural actomyosin with different degrees of hydrolysis	39
4. Statistic analytical	40
3. Results and Discussion	41
3.1 Chemical composition and property of white shrimp, bigeye snapper and lizardfish mince	41
3.1.1 Proximate composition	41
3.1.2 Proteins and non-protein nitrogenous compounds	41
3.1.3 Thermal denaturation	44
3.2 Effect of setting condition and MTGase on gel forming ability of some marine fish and shrimp meats	45
3.2.1 Breaking force and deformation	45
3.2.2 Color and whiteness	48
3.2.3 Expressible moisture content	52
3.2.4 Protein patterns	53
3.2.5 Microstructure	56
3.3 Effect of post-mortem storage on protein cross-linking and gel enhancing ability of MTGase on mince from some marine fish and shrimp	58
3.3.1 Chemical changes during iced storage	58
3.3.1.1 Changes in pH	58

CONTENTS (Continued)

	Page
3.3.1.2 Changes in TVB-N and TMA-N contents	59
3.3.1.3 Changes in TBARS	61
3.3.1.4 Changes in formaldehyde content	63
3.3.1.5 Changes in TCA-soluble peptide content	64
3.3.1.6 Changes in protein patterns	65
3.3.1.7 Changes in ATPase activity	66
3.3.2 Effect of MTGase on gel-forming ability of some marine fish and shrimp stored in ice for different times	70
3.3.2.1 Breaking force and deformation	70
3.3.2.2 Color and whiteness	74
3.3.2.3 Expressible moisture content	78
3.3.2.4 Protein patterns	81
3.3.2.5 Microstructure	83
3.4 Effect of protein substrate denaturation on cross-linking activity of MTGase	85
3.4.1 Thermal stability	85
3.4.2 Effect of MTGase on cross-linking and physico-chemical properties of NAM with different degrees of denaturation	86
3.4.2.1 Changes in free amino group content	86
3.4.2.2 Changes in solubility	89
3.4.2.3 Changes in protein patterns	92
3.4.2.4 Changes in sulfhydryl group content	95
3.4.2.5 Changes in disulfide bond content	98
3.4.2.4 Changes in surface hydrophobicity	100
3.5 Effect of MTGase on cross-linking and physicochemical properties of NAM added with different levels of formaldehyde	102
3.5.1 Changes in free amino group content	102

CONTENTS (Continued)

	Page
3.5.2 Changes in solubility	104
3.5.3 Changes in protein patterns	106
3.5.4 Changes in sulfhydryl group content	108
3.5.5 Changes in disulfide bond content	110
3.5.6 Changes in surface hydrophobicity	112
3.6 Effect of MTGase on cross-linking and physicochemical properties of NAM with different degrees of hydrolysis	114
3.6.1 Changes in physicochemical properties of NAM with different degrees of hydrolysis	114
3.6.1.1 Changes in surface hydrophobicity	114
3.6.1.2 Changes in free amino group content	115
3.6.1.3 Changes in protein patterns	116
3.6.2 Effect of MTGase on cross-linking and physico-chemical properties of NAM with different degrees of hydrolysis	117
3.6.2.1 Changes in free amino group content	117
3.6.2.2 Changes in solubility	119
3.6.2.3 Changes in protein patterns	121
3.6.2.4 Changes in sulfhydryl group content	122
3.6.2.5 Changes in disulfide bond content	125
3.6.2.6 Changes in surface hydrophobicity	127
4. Conclusion	129
References	130
Appendix	154
Vitae	160

LIST OF TABLES

Table	Page
1. Composition of fish and shellfish meat	2
2. Contractile proteins in food myosystems	3
3. Conformational changes which may occurring during the thermal denaturation of natural actomyosin	14
4. Suwari forming capacity of different fish species	19
5. Source and characteristics of some TGases	21
6. Proximate composition of mince from some marine fish and shrimp muscles	41
7. Nitrogenous constituents in the mince from some marine fish and shrimp muscles	42
8. T_{max} and enthalpy (ΔH) of mince from some marine fish and shrimp muscles	44
9. L^* , a^* , b^* -values and whiteness of gels from white shrimp mince added with various levels of MTGase and different setting conditions	49
10. L^* , a^* , b^* -values and whiteness of gels from bigeye snapper mince added with various levels of MTGase and different setting conditions	50
11. L^* , a^* , b^* -values and whiteness of gels from lizardfish mince added with various levels of MTGase and different setting conditions	51
12. Expressible moisture content of mince gels from some marine fish and shrimp added with various levels of MTGase and different setting conditions	53
13. L^* , a^* , b^* -values and whiteness of mince gels from white shrimp stored in ice for different times without and with MTGase addition (0.6 units/g) and set under different conditions	75
14. L^* , a^* , b^* -values and whiteness of mince gels from bigeye snapper stored in ice for different times without and with MTGase addition (0.6 units/g) and set under different conditions	76
15. L^* , a^* , b^* -values and whiteness of mince gels from lizardfish stored in ice for different times without and with MTGase addition (0.6 units/g) and set under different conditions	77

LIST OF TABLES (Continued)

Table	Page
16. Thermal inactivation rate constant ($K_D \times 10^{-5} S^{-1}$) of NAM from some marine fish and shrimp meats	86

LIST OF FIGURES

Figure	Page
1. Model of myosin molecule	5
2. (a) A thin filament of muscle formed by the filament of tropomyosin molecules wound in each of the two grooves of the actin helix. (b) Proposed model for configuration of actin, tropomyosin and troponin (Tn) subunits	7
3. Formation of a gel network structure	16
4. Reaction catalyzed by TGase	20
5. Primary structure of microbial transglutaminase (MTGase).	23
6. Scheme for gel preparation	31
7. Electrophoretic pattern of various protein fractions of mince from some marine fish and shrimp	43
8. Breaking force and deformation of mince gels from some marine fish and shrimp added with various levels of MTGase and set under different conditions	47
9. SDS-PAGE pattern of mince gels from some marine fish and shrimp added with various levels of MTGase and set under different conditions	55
10. Microstructure of gels from some marine fish and shrimp added without and with MTGase (0.6 units/g) and set under different conditions	57
11. Changes in pH of some marine fish and shrimp during the iced storage	59
12. Changes in TVB-N (A) and TMA-N (B) contents of some marine fish and shrimp during the iced storage	61
13. Changes in TBARS of some marine fish and shellfish during the ice storage	62
14. Changes in formaldehyde content of some marine fish and shellfish during the iced storage	64
15. Changes in TCA-soluble peptide content of some marine fish and shellfish during the iced storage	65
16. SDS-PAGE pattern of muscle proteins from some marine fish and shrimp during the iced storage	66

LIST OF FIGURES (Continued)

Figure	Page
17. ATPase activities of natural actomyosin of some marine fish and shrimp during the iced storage	68
18. Ca ²⁺ -sensitivity of natural actomyosin extracted from some marine fish and shrimp during the iced storage	69
19. Breaking force and deformation of mince gels from some marine fish and shrimp stored in ice for different times without and with MTGase addition (0.6 units/g) and set under different conditions	73
20. Expressible moisture content of mince gels from some marine fish and shrimp stored in ice for different times without and with MTGase addition (0.6 units/g) and set under different conditions	80
21. SDS-PAGE patterns of mince gels from some marine fish and shrimp stored in ice for different times without and with MTGase addition (0.6 units/g) and set under different conditions	81
22. Microstructure of mince gels from some marine fish and shrimp stored in ice for 10 days without and with MTGase addition (0.6 units/g) and set under different conditions	84
23. Changes in free amino group content of some marine fish and shrimp NAM with pre-heating at 50°C for different times as influenced by MTGase addition at different levels	88
24. Changes in solubility of some marine fish and shrimp NAM with pre-heating at 50°C for different times as influenced by MTGase addition at different levels	91
25. Changes in protein patterns of some marine fish and shrimp NAM with pre-heating at 50°C for different times as influenced by MTGase addition	94
26. Changes in sulfhydryl group content of some marine fish and shrimp NAM with pre-heating at 50°C for different times as influenced by MTGase addition at different levels	97

LIST OF FIGURES (Continued)

Figure	Page
27. Changes in disulfide bond content of some marine fish and shrimp NAM with pre-heating at 50°C for different times as influenced by MTGase addition at different levels	99
28. Changes in surface hydrophobicity of some marine fish and shrimp NAM with pre-heating at 50°C for different times as influenced by MTGase addition at different levels	101
29. Changes in free amino group content of some marine fish and shrimp NAM added with different levels of formaldehyde as influenced by MTGase addition at different levels	103
30. Changes in solubility of some marine fish and shrimp NAM added with different levels of formaldehyde as influenced by MTGase addition at different levels	105
31. Protein patterns of some marine fish and shrimp NAM added with different levels of formaldehyde as influenced by MTGase addition	107
32. Changes in sulfhydryl group content of some marine fish and shrimp NAM added with different levels of formaldehyde as influenced by MTGase addition at different levels	109
33. Changes in disulfide bond content of some marine fish and shrimp NAM added with different levels of formaldehyde as influenced by MTGase addition at different levels	111
34. Changes in surface hydrophobicity of some marine fish and shrimp NAM added with different levels of formaldehyde as influenced by MTGase addition at different levels	113
35. Surface hydrophobicity of NAM from some marine fish and shrimp with different degrees of hydrolysis	114
36. Free amino group content of NAM from some marine fish and shrimp with different degrees of hydrolysis	115

LIST OF FIGURES (Continued)

Figure	Page
37. SDS-PAGE of NAM from some marine fish and shrimp with different degrees of hydrolysis	116
38. Changes in free amino group content of NAM from some marine fish and shrimp as influenced by different degrees of hydrolysis	118
39. Changes in solubility of NAM from some marine fish and shrimp as influenced by different degrees of hydrolysis	120
40. Protein patterns of NAM hydrolysate from some marine fish and shrimp with different degrees of hydrolysis added without and with MTGase at different levels and incubated at 40°C for 30 min	122
41. Changes in sulfhydryl group content of NAM from some marine fish and shrimp as influenced by different degrees of hydrolysis	124
42. Changes in disulfide bond content of NAM from some marine fish and shrimp as influenced by different degrees of hydrolysis	126
43. Changes in surface hydrophobicity of NAM from some marine fish and shrimp as influenced by different degrees of hydrolysis	128