

## REFERENCES

- Adler-Nissen, J. 1986. Enzymic hydrolysis of food proteins. Elsevier Applied Science Publishers. Barking. UK.
- Adler-Nissen, J and Olsen, H.S. 1979. The influence of peptide chain length on taste and functional properties of enzymatically modified soy protein. *In* Functionality and protein structure. (Pour-El, A., ed.). American Chemical Society Symposium Series 92. Washington, DC.
- Ahn, D.U., Ajuyah, A., Wolfe, F.H. and Sim, J.S. 1993. Oxygen availability affects prooxidant catalyzed lipid oxidation of cooked turkey patties. *J. Food Sci.* 58: 278-282.
- Ahn, D.U., Olson, D.G., Jo, C., Chen, X., Wu C. and Lee, J.J. 1998. Effect of muscle type, packaging, and irradiation on lipid oxidation, volatile production and color in raw pork patties. *Meat Sci.* 49: 37-39.
- Ajandouz, E. H., Tchiakpe, L. S., Ore, F. D., Benajibas, A., and Puigserver, A. 2001. Effect of pH on caramelisation and Maillard reaction kinetics in fructose-lysine model systems. *J of Food Sci.* 66: 926-932.
- Amarowicz, R. and Shahidi, F. 1997. Antioxidant activity of peptide fractions of capelin protein hydrolysates. *Food Chem.* 58: 355-359.
- Angeles Navarrete del Toro, M. and Garcia-Carreno, F.L. 2002. Evaluation of the progress of protein hydrolysis. *In* Current Protocols in Food Analytical Chemistry, (B.2.2.7). Wiley, New York.
- AOAC. 2000. Official Method of Analysis, 17<sup>th</sup> ed. Association of Official Chemists. Gaithersberg, Maryland.
- Arnao, M. B., Cano, A. and Acosta, M. 2001. The hydrophilic and lipophilic contribution to total antioxidant activity. *Food Chem.* 73: 239-244.
- Artharn, A., Benjakul, S., Prodpran, T. and Tanaka, M. 2007. Properties of a protein-based film from round scad (*Decapterus maruadsi*) as affected by muscle types and washing. *Food Chem.* 103: 867-874.
- Aspmo, S.I., Horn, S.J. and Eijsink, V.G.H. 2005. Enzymatic hydrolysis of Atlantic cod (*Gadus morhua* L.) viscera. *Process Biochem.* 40: 1957-1966.

- Baek, H.H. and Cadwallader, K.R. 1995. Enzymatic hydrolysis of crayfish processing by-products. *J. Food Sci.* 60: 929-935.
- Bandyopadhyay, M., Chakraborty, R. and Raychaudhuri, U. 2007. A process for preparing a natural antioxidant enriched dairy product (Sandesh). *LWT-Food Sci. Technol.* 40: 842-851.
- Benjakul, S., Lertittikul, W. and Bauer, F. 2005. Antioxidant activity of Maillard reaction products from a porcine plasma protein-sugar model system. *Food Chem.* 93: 189-196.
- Benjakul, S. and Morrissey, M. 1997. Protein hydrolysates from Pacific whiting solid wastes. *J. Agric. Food Chem.* 45: 3423-3430.
- Benjakul, S., Seymour, T. S., Morrissey, M. T. and An, H. 1997. Physicochemical changes in Pacific whiting muscle proteins during iced storage. *J. Food Sci.* 62: 729-733.
- Benjakul, S., Visessanguan, W., Phongkanpai, V. and Tanaka, M. 2005. Antioxidative activity of caramelisation products and their preventive effect on lipid oxidation in fish mince. *Food Chem.* 90: 231-239.
- Benzie, I. F. F. and Strain, J.J. 1996. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power" the FRAP assay. *Anal Biochem.* 239: 70-76.
- Beuge, J. A. and Aust, S.D. 1978. Microsomal lipid peroxidation. *Methods Enzymol.* 52: 302-310.
- Bishov, S.J. and Henick, A.S. 1975. Antioxidant effect of protein hydrolyzates in freeze-dried model systems. Synergistic action with a series of phenolic antioxidants. *J. Food Sci.* 40: 345-348.
- Blendford, D.E. 1994. Protein hydrolysate: functionalities and uses in nutritional products. *Int. Food Ingr.* 3: 45-49.
- Bligh, E.G. and Dyer, W.J. 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 37: 911-931.
- Buinov, A.A., Ginzburg, A.S. and Syroedov, V.I. 1977. Hygroscopic properties of fish protein hydrolysates, dried as a foam. *Izvestiya Vysshikh Uchebnykh Zavedenii, Pishchevaya Tekhnologiya.* 3: 110-113.

- Carabasa-Giribet, M. and Ibarz-Ribas, A. 2000. Kinetics of colour development in aqueous glucose systems at high temperatures. *J. Food Eng.* 44: 181-189.
- Castell, C.H., Maclean, J. and Moore, B. 1965. Rancidity in lean fish muscle. IV. Effect of sodium chloride and other salts. *J. Fish Res. Board. Can.* 22: 929-944.
- Chan, W.K.M., Decker, E.A., Lee, J.B. and Butterfield, D.A. 1994. EPR spin-trapping studies of the hydroxyl radical scavenging activity of carnosine and related dipeptides. *J. Agric. Food Chem.* 42: 1407-1410.
- Chaveesuk, R., Smith, J.P. and Simpson, B.K. 1993. Production of fish sauce and acceleration of sauce fermentation using proteolytic enzymes. *J. Aqua. Food Prod. Technol.* 2(3): 59-77.
- Cheftel, J.C., Cuq, J.L. and Lorient, D. 1985. Amino acids; peptides, and proteins. *In Food Chemistry* (Fennema O.R. ed.), pp. 245–370. Marcel Dekker. New York.
- Chen, R.H. and Hwa, H.D. 1996. Effect of molecular weight of chitosan with the same degree of deacetylation on thermal, mechanical and permeability properties of the prepared membrane. *Carbohydr. Polym.* 29: 353–358.
- Chen, H. and Meyers, S. P. 1982. Extraction of astaxanthin pigment from crawfish waste using a soy oil process. *J. Food Sci.* 47: 892–900.
- Chen, H.M., Muramoto, K. and Yamauchi, F. 1995. Structural analysis of antioxidative peptides from soybean  $\beta$ -conglycinin. *J. Agric. Food Chem.* 43: 574-578.
- Chen, H.M., Muramoto, K., Yamauchi, F., Fujimoto, K. and Nokihara, K. 1998. Antioxidative properties of histidine-containing peptides designed from peptide fragments found in the digests of a soybean protein. *J. Agric. Food Chem.* 46: 49-53.
- Chen, H.M., Muramoto, K., Yamauchi, F. and Nokihara, K. 1996. Antioxidant activity of designed peptides based on the antioxidative peptide isolated from digests of a soybean protein. *J. Agric. Food Chem.* 44: 2619-2623.
- Chen, C.H., Pearson, A.M. and Gray, J.L. 1992. Effect of synthetic antioxidants (BHA, BHT and PG) on mutagenicity of IQ-like compounds. *Food Chem.* 43: 177-183.
- Clifford-Hall, I.I.I. 2001. Sources of natural antioxidants: oilseeds, nuts, cereals, legumes, animal products and microbial sources. *In Antioxidants in Food: practical applications.* (Pokorny, J., Yanishlieva, N. and Gordon, M. eds.), pp. 159-209. Woodhead Publishing. Abington, England.

- Cortas, N.K. and Wakid, N.W. 1990. Determination of inorganic nitrate in serum and urine by a kinetic cadmium-reduction method. *Clin. Chem.* 36: 1440–1443.
- Coward-kelly, G., Agbogbo, F.K. and Holtzapfle. 2006. Lime treatment of shrimp head waste for the generation of highly digestible animal feed. *Carbohydr. Polym.* 97: 1515-1520.
- Damodaran, S. 1996. Enzymatic hydrolysis. *In Food Chemistry*. 3<sup>rd</sup> ed. (Fennema, O.R. ed.), pp. 422-423. Marcel Dekker. New York.
- Daukšas, E., Šližyte, R., Rustard, T. and Storrø1, I. 2004. Bitterness in fish protein hydrolysates and methods for removal. *J. Aquat. Food Prod. Technol.* 13(2): 101-114.
- Decker, A.E. 1998. Antioxidant mechanism. *In Food lipid: chemistry, nutrition, and biotechnology*. (Akoh, C.C and Min, D.B., eds.), pp. 397-448. Marcel Decker. New York.
- Decker, E.A. and Crum, A.D. 1993. Antioxidant activity of carnosine in cooked ground pork. *Meat Sci.* 34: 245-253.
- Decker, E.A. and Hultin, H.O. 1990. Factors influencing catalysis of lipid oxidation by the soluble fraction of mackerel muscle. *J. Food Sci.* 55: 947–950.
- Decker, H. and Tucek, F. 2000. Tyrosinase/catecholoxidase activity of hemocyanins: structural basis and molecular mechanism. *Trends Biochem. Sci.* 25: 392-397.
- Dinis, F.M. and Martin, A.M. 1997. Effects of the extent of enzymatic hydrolysis on functional properties of shark protein hydrolysate. *Lebensm.-Wiss. u.-Technol.* 30: 266–272.
- Duh, P.D. 1998. Antioxidant activity of burdock (*Arctium lappa linne*): its scavenging effect on free radical and active oxygen. *J. Am. Oil Chem. Soc.* 75: 455-461.
- Dziezak, J.D. 1986. Antioxidants: The ultimate answer to oxidation. *Food Technol.* 40: 94-102.
- Egan, H., Kirk, R. and Sawyer, R. 1981. Measurement of rancidity. *In Peason's Chemical Analysis of Foods*. 8<sup>th</sup> ed. pp. 534-539, Churchill Livingstone, New York.
- Egorov, S.Y., Kurella, E.G., Boldrynev, A.A. and Kasnovski, A.A. 1992. The quenching of singlet molecular oxygen by carnosine and anserine in aqueous solution. *Bioorg. Khim.* 18: 142-144.
- Fagbenro, O.A. 1996. Preparation, properties and preservation of lactic acid fermented shrimp heads. *Food Res. Int.* 29: 595-599.

- Farombi, E.O., Britton, G. and Emerole, G.O. 2000. Evaluation of the antioxidant activity and partial characterisation of extracts from browned yam flour diet. *Food Res. Int.* 33: 493-499.
- Fennema, O. R. 1996. Water and Ice. *In Food Chemistry* (Fennema, O. R. ed.), pp. 17-94. Marcel Dekker. New York.
- Fernandez-Lopez, J., Sevilla, L., Sayas-Barbera, M.E., Navarro, C., Marin, F. and Perez-Alvarez, J.A. 2003. Evaluation of the antioxidant potential of hyssop (*Hyssopus officinalis* L.) and rosemary (*Rosmarinus officinalis* L.) extract in cooked pork meat. *J. Food Sci.* 68: 660-664.
- Fountoulakis, M. and Lahm, H.W. 1998. Hydrolysis and amino acid composition analysis of proteins. *J. Chromatogr. A.* 109: 109-134.
- Frankel, E. N., Huang, S. W. and Aeschbach, R. 1997. Antioxidant activity of green tea in different lipid systems. *J. Am. Oil Chem. Soc.* 74: 1309-1315.
- Friedman, M. 1999. Chemistry, biochemistry, nutrition, and microbiology of lysinoalanine, lanthionine and histidinoalanine in food and other proteins. *J. Agric. Food Chem.* 47: 1295-1319.
- Gbogouri, G.A., Linder, M., Fanni, J. And Parmentier, M. 2004. Influence of hydrolysis degree on the function properties of salmon byproducts hydrolysates. *Food Chem. Toxic.* 69: 615-622.
- German, J.B. and Kinsella, J.E. 1985. Lipid oxidation in fish tissue: Enzymatic initiation via lipoxygenase. *J. Agric. Food. Chem.* 33: 680-683.
- Giese, J. 1996. Antioxidants : Tools for preventing lipid oxidation. *Food Technol.* 50 (11): 73-81.
- Gil, B., Yoon, S.H. and Choi, E.S. 2002. A fryer design with constant specific surface area. *J. Am. Oil Chem. Soc.* 79: 511-513.
- Gildberg, A., Batista, I. and Strøm, E. 1989. Preparation and characterization of peptones obtained by a two-step enzymatic hydrolysis of whole fish. *Biotechnol. Appl. Biochem.* 11: 413-423.
- Gildberg, A. and Stenberg, E. 2001. A new process for advanced utilization of shrimp waste. *Process Biochem.* 36: 809-812.

- Gonzalez-Tello, P., Camacho, F., Jurado, E., Paez, M.P. and Guadix, E.M. 1994. Enzymatic hydrolysis of whey proteins. II. Molecular-weight range. *Biotechnol. Bioeng.* 44: 529-532.
- Guadix, A., Camacho, F. and Guadix, E.M. 2006. Production of whey protein hydrolysate with reduced allergenicity in a stable membrane reactor. *J. Food Eng.* 72: 398-405.
- Guérard, F., Dufossé, L., De La Broise, D. and Binet, A. 2001. Enzymatic hydrolysis of proteins from yellowfin tuna (*Thunnus albacares*) wastes using Alcalase. *J. Mol. Catal. B: Enzymol.* 11: 1051-1059.
- Hale, M.B. 1972. Making fish protein concentrate by enzymatic hydrolysis. NOAA technical report NMFS SSRF-675. p.1-31. Department of commerce. Seattle, USA.
- Haque, Z.U. 1993. Influence of milk peptides in determining the functionality of milk proteins: a review. *J. Dairy Sci.* 76: 311-320.
- Hatate, H., Nagata, Y., and Kochi, M. 1990. Antioxidative effect of bovine serum albumin hydrolyzates and their synergistic effect with antioxidants. *Yukagaku.* 39: 42-46.
- He, H., Chen, X., Sun, C., Zhang, Y. and Gao, P. 2006. Preparation and functional evaluation of oligopeptide-enriched hydrolysate from shrimp (*Acetes chinensis*) treated with crude protease from *Bacillus* sp. SM98011. *Biores. Technol.* 97: 385-390.
- Hider, R.C., Liu, Z.D. and Khodr, H. H. 2001. Metal chelation of polyphenols. *Meth. Enzymol.* 335: 190-203.
- Hoyle, N.T. and Merritt, J.H. 1994. Quality of fish protein hydrolysate from herring (*Clupea harengus*). *J. Food Sci.* 59: 76-80.
- Hyun, C.K. and Shin, H.K. 2000. Utilization of bovine blood plasma proteins for the production of angiotensin I converting enzyme inhibitory peptides. *Process Biochem.* 36: 65-71.
- Ichihashi, H., Kohno, H., Kannan, K., Tsumura, A. and Yamsaki, S. 2001. Mutielement analysis of purpleback flying squid using high resolution inductively coupled plasma-mass spectrometry (HR-ICP-MS). *Environ. Sci. Technol.* 35: 3103-3108.
- Imm, J.Y. and Lee, C.M. 1999. Production of seafood flavor from red hake (*Urophycis chuss*) by enzymatic hydrolysis. *J. Agric. Food Chem.* 47: 2360-2366.

- Jadhav, S.J., Nimbalkar, S.S., Kulkarni, A.D. and Madhavi, D.L. 1996. Lipid oxidation in biological and food systems. *In Food antioxidants*. (Madhavi, D.L. *et al.*, eds.), p. 5-64. Marcel Dekker. New York.
- Jakobsen, M. and Bertelsen, G. 2000. Colour stability and lipid oxidation of fresh beef. Development of a response surface model for predicting the effects of temperature, storage time and modified atmosphere composition. *Meat Sci.* 54: 49-57.
- Jao, C.L. and Ko, W.C. 2002. 1,1-Diphenyl-2-picrylhydrazyl (DPPH) radical scavenging by protein hydrolysates from tuna cooking juice. *Fish. Sci.* 68: 430-435.
- Jayaprakasha, G.K., Singh, R.P. and Sakariah, K.K. 2001. Antioxidant activity of grape seed (*Vitis vinefera*) extracts on peroxidation models *in vitro*. *Food Chem.* 73: 285-290.
- Je, J.Y., Park, P.J. and Kim, S.K. 2005. Antioxidant activity of a peptide isolated from Alaska pollack (*Theragra chalcogramma*) frame protein hydrolysate. *Food Res. Int.* 38: 45-50.
- Je, J.Y., Park, P.J., Kwon, J.Y. and Kim, S.K. 2004. A novel angiotensin I converting enzyme inhibitory peptide from Alaska Pollack (*Theragra chalcogramma*) frame protein hydrolysate. *J. Agric. Food Chem.* 52: 7842-7845.
- Je, J.Y., Qian, Z.J., Byun, H.G. and Kim, S.K. 2007. Purification and characterization of an antioxidant peptide obtained from tuna backbone protein by enzymatic hydrolysis. *Process Biochem.* 42: 840-846.
- Jeon, Y.J., Byun, H.G. and Kim, S.K. 1999. Improvement of functional properties of cod frame protein hydrolysates using ultrafiltration membranes. *Process Biochem.* 35: 471-478.
- Jimenez-Escrig, A., Rincon, M., Pulido, R. and Saura-Calixto, F. 2001. Guava fruit (*Psidium guajava L.*) as a new source of antioxidant dietary fiber. *J. Agric. Food Chem.* 49: 5489-5493.
- Johnson, A.H. 1974. Hydrolysis. *In Encyclopedia of food technology* (Johnson, A.H. and Peterson, M.S. eds.), pp. 514-517. The Avi publishing company, Inc. Connecticut, USA.
- Johnston-Banks, F.A., 1990. Gelatin. *In Food gels* (Harris, P. ed.), pp. 233-289. Elsevier Applied Science. London.
- Jun, S. Y., Park, P. J., Jung, W. K., and Kim, S. K. 2004. Purification and characterization of an antioxidative peptide from enzymatic hydrolysate of yellowfin sole (*Limanda aspera*) frame protein. *Eur. Food Res. Technol.* 219: 20-26.

- Jung, W.K., Mendis, E., Je, J.Y., Park, P.J., Son, B.W., Kim, H.C., Choi, Y.K. and Kim, S. 2006. Angiotensin I converting enzyme inhibitory peptide from yellowfin sole (*Limanda aspera*) frame protein and its antihypertensive effect in spontaneously hypertensive rats. Food Chem. 94: 26-32.
- Kamil, J.Y.V.A., Jeon, Y.J. and Shahidi, F. 2002. Antioxidative activity of chitosan of different viscosity in cooked comminuted fish of herring (*Clupea harengus*). Food Chem. 79: 69-77.
- Karpinska, M., Borowski, J. and Danowska-Oziewicz, M. 2001. The use of antioxidants in ready-to-serve food. Food Chem. 72:5-9.
- Khokhar, S. and Apenten, R.K. O. 2003. Iron binding characteristics of phenolic compounds: some tentative structure–activity relations. Food Chem. 81: 133–140.
- Kim, S.Y., Je, J.Y. and Kim, S.K. 2006. Purification and characterization of antioxidant peptide from hoki (*Johnius belengerii*) frame protein by gastrointestinal digestion. J. Nutr. Biochem. 18: 31-38.
- Kim, S.K., Kim, Y.T., Byun, H.G., Nam, K.S., Joo, D.S. and Shahidi, F. 2001. Isolation and characterization of antioxidative peptide from gelatin hydrolysate of Alaska pollack skin. J. Agric. Food Chem. 49: 1984-1989.
- Kinsella, J.E. 1976. Functional properties of proteins in food: a survey. Crit. Rev. Food Sci. Nutr. 8: 219–280.
- Kinsella, J.E. 1981. Functional properties of proteins: possible relationships between structure and function in foams. Food Chem. 7: 273-288.
- Klompong, V., Benjakul, S., Kantachote, D. and Shahidi, F. 2007. Antioxidative activity and functional properties of protein hydrolysate of yellow stripe trevally (*Selaroides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. Food Chem. 102: 1317-1323.
- Kohama, Y., Matsumoto, S., Oka, H., Teramoto, T., Okabe, M. and Mimura, T. 1998. Isolation of angiotensin-converting enzyme inhibitor from tuna muscle. Biochem. Biophys. Res. Com. 155: 332-337.



- Kolakoska, A., Olley, J. and Dunstan, G.A. 2002. Fish lipids. *In* Chemical and functional Properties of Food Lipids (Sikorski, Z.E. and Kolakoska, A. eds.), pp. 221-264. CRC Press. Florida.
- Kristinsson, H.G. and Rasco, B.A. 2000a. Fish protein hydrolysates: production, biochemical and functional properties. *Crit. Rev. Food Sci. Nutr.* 40: 43-81.
- Kristinsson, H.G. and Rasco, B.A. 2000b. Biochemical and functional properties of Atlantic salmon (*Salmo salar*) muscle proteins hydrolyzed with various alkaline proteases. *J. Agric. Food Chem.* 48: 657–666.
- Kuhn, H. 1999. Lipoxygenases. *In* Prostaglandins, Leukotrienes and Other Eicosanoids (Marks, F. and Fustenberger, G. eds.), p 109. Wiley-VCH. Weinheim.
- Lahl, W.J. and Braun, S.D. 1994. Enzymatic production of protein hydrolysates for food use. *Food Technol.* 48(10): 68-71.
- Lee, B, J. and Hendricks, D, G. 1997. Antioxidant effects of L-carnosine on liposomes and beef homogenates. *J. Food Sci.* 62: 931-934.
- Leffler, A. 1986. Proteolytic enzymes: sources and applications. *Food Technol.* 40(1): 63-69.
- Leong, L. P. and Shui, G. 2002. An investigation of antioxidant capacity of fruits in singapore markets. *Food Chem.* 76. 69-75.
- Lertittikul, W. 2005. Porcine plasma protein-sugar Maillard reaction products (MRPs): some factors affecting antioxidant activity and their application. Master of Science Thesis. Prince of Songkla University. 159 p.
- Liaset, B., Lied, E. and Espe, M. 2000. Enzymatic hydrolysis of by-products from the fish-filleting industry; chemical characterisation and nutritional evaluation. *J. Sci. Food Agric.* 80: 581-589.
- Liaset, B., Nortvedt, R., Lied, E. and Espe, M. 2002. Studies on the nitrogen recovery in enzymic hydrolysis of Atlantic salmon (*Salmo salar* L.) frames by Protamex (TM) protease. *Process Biochem.* 37: 1263–1269.
- Liceaga-Gesualdo, A.M. and Li-Chan, E.C.Y. 1999. Functional properties of fish protein hydrolysates from herring (*Clupea harengus*). *J. Food Sci.* 64: 1000–1004.
- Linder, M., Fanni, J., Parmentier, M., Sergent, M. and Phan-Than-Luu, R. 1995. Protein recovery from veal bones by enzymatic hydrolysis. *J. Food Sci.* 60: 949-952.

- Lopetcharat, K. and Park, J.W. 2002. Characteristics of fish sauce made from Pacific whiting and surimi by products during fermentation stage. *J. Food Sci.* 67: 511-516.
- Lopez-Cervantes, J., Sanchez-Machado, D.I. and Rosas-Rodriguez, J.A. 2006. Analysis of free amino acids in fermented shrimp waste by high-performance liquid chromatography. *J. Chromatogr. A.* 1105: 106–110.
- Mackie, J.M. 1982. Fish protein hydrolysates. *Process Biochem.* 17: 26-31.
- Madhavi, D.L., Despande S.S. and Salunkhe D.K. 1996. Toxicological aspects of food antioxidants. *In Food antioxidants.* pp.267-311. Marcel Dekker. New York.
- Mahmoud, M.I. 1994. Physicochemical and functional properties of protein hydrolysates in nutritional products. *Food Technol.* 48:89–95.
- Mahmoud, M.I., Malone, W.T. and Cordle, C.T. 1992. Enzymatic hydrolysis of casein: effect of degree of hydrolysis on antigenicity and physical properties. *J. Food Sci.* 57: 1223-1229.
- Mandeville, S., Yaylayan, V. and Simpson, B. K. 1992. Proximate analysis, isolation and identification of amino acids and sugars from raw and cooked commercial shrimp waste. *Food Biotechnol.* 6: 51–64.
- Mansour, E.H. and Khalil, A.H. 2000. Evaluation of antioxidant activity of some plant extracts and their applications to ground beef patties. *Food Chem.* 69: 135-141.
- Matmaroh, K., Benjakul, S. and Tanaka, M. 2006. Effect of reactant concentrations on the Maillard reaction in a fructose–glycine model system and the inhibition of black tiger shrimp polyphenoloxidase. *Food Chem.* 98: 1–8.
- Matthaus, B. 2002. Antioxidant activity of extracts obtained from residues of different oilseeds. *J. Agric. Food Chem.* 50: 3444-3452.
- McCarthy, T.L., Kerry, J.P., Kerry, J.F., Lynch, P.B. and Buckley, D.J. 2001. Evaluation of the antioxidant potential of natural food/plant extracts as compared with synthetic antioxidants and vitamin E in raw and cooked pork patties. *Meat Sci.* 58: 45-52.
- Mendis, E., Rajapakse, N., Byun, H.G. and Kim, S.K. 2005. Investigation of jumbo squid (*Dosidicus gigas*) skin gelatin peptide for their *in vitro* antioxidant effects. *Life Sci.* 77: 2166-2178.

- Mendis, E., Rajapakse, N. and Kim, S.K. 2005. Antioxidant properties of a radical-scavenging peptide purified from enzymatically prepared fish skin gelatin hydrolysate. *J. Agric. Food Chem.* 53: 581-587.
- Meyers, SP. 1986. Utilization of shrimp processing waste. *Infofish Marketing Digest.* 4: 18-19.
- Mohr, V. 1980. Enzymes technology in the meat and fisheries industries. *Process Biochem.* 15: 18-21.
- Monahan, F.J. 2000. Oxidation of lipids in muscle foods: fundamental and applied concerns. *In* Antioxidants in muscle foods. (Decker, E.A., Faustman, C. and Lopez-Bote, C.J. eds), p. 3. Wiley-Interscience. New York.
- Morales, F. J. and Jimenez-Perez, S. 2001. Free radical scavenging capacity of Maillard reaction products as related to colour and fluorescence. *Food Chem.* 72: 119-125.
- Morales, F. J., Romeo, C. and Jimenez-Perez. 1996. Fluorescence associated with Maillard reaction in milk and milk-resembling systems. *Food Chem.* 57: 423-428.
- Moran, J. F., Klucas, R. V., Grayer, R. J., Abian, J. and Becana, M. 1997. Complexes of iron with phenolic compounds from soybean nodules and other legume tissues: prooxidant and antioxidant properties. *Free Rad. Biol. Med.* 22: 861-870.
- Moridani, M. Y., Pourahmad, J., Bui, H., Siraki, A., and O'Brien, P. J. 2003. Dietary flavonoid iron complexes as cytoprotective superoxide radical scavengers. *Free Rad. Biol. Med.* 34: 243-253.
- Morr, C.V. 1985. Composition, physicochemical and functional properties of reference whey protein concentrates. *J. Food Sci.* 50: 1406-1411.
- Mullally, M.M., O'Callaghan, D.M., FitzGerald, R.J., Donnelly, W.J. and Dalton, J.P. 1995. Zymogen activation in pancreatic endoproteolytic preparations and influence on some whey protein characteristics. *J. Food Sci.* 60: 227-233.
- Murueta, J.H.C., Toro, M.A.N. and Carreno, F.G. 2007. Concentrate of fish protein from bycatch species produced by various drying processes. *Food Chem.* 100: 705-711.
- Nakayama, T., Osawa, T., Mendosa, E.N.T., Laurena, A.C. and Kawakishi, S. 1994. Comparative study of antioxidative assays of plant materials. *In* Postharvest biochemistry of plant food-materials in the tropics. (Uritani, I. *et al.*, eds.), p. 241-251. Japan Scientific Societies Press. Japan.

- Namiki, M. 1990. Antioxidant/antimutagens in food. *Crit. Rev. Food Sci. Nutr.* 29: 273-300.
- Nawar, W.W. 1996. Lipid. *In Food Chemistry.* (Fennema, O.R. ed.), pp. 210-243. New York: Marcel Dekker.
- Nielsen, P.M. 1997. Functionality of protein hydrolysates. *In Food protein and their applications.* (Damodaran, S. and Paraf, A. eds.), pp. 443-471. New York: Marcel Dekker.
- Niki, E. 1987. Antioxidants in relation to lipid peroxidation. *Chem. Phys. Lipids.* 44: 227-253.
- Nilsang, S., Lertsiri, S., Suphantharika, M. And Assavanig, A. 2005. Optimization of enzymatic hydrolysis of fish soluble concentrate by commercial proteases. *J. Food Eng.* 70: 571-578.
- Nwanna, L.C., Balogun, A.M., Ajenifuja, Y.F. and Enujigha, V.N. 2004. Replacement of fish meal with chemically preserved shrimp head in the diets of African catfish, *Clarias gariepinus*. *Food Agric. Environ.* 2: 79-83.
- Ogasawara, M., Katsumata, T. And Egi, M. 2006. Taste properties of Maillard-reaction products prepared from 1000 to 5000 Da peptide. *Food Chem.* 99: 600-604.
- Oleary, V., Darley-USmar, V.M., Russell, L.J. and Stone, D. 1992. Pro-oxidant effects of lipoxygenase-derived peroxides on the copper-initiated oxidation of low-density lipoprotein. *J. Biochem Sci.* 282: 631-634.
- Ono, S., Hosokawa, M., Miyashita, K. and Takahashi, K. 2003. Isolation of peptides with angiotensin I-converting enzyme inhibitory effect derived from hydrolysate of upstream chum salmon muscle. *J. Food Sci.* 68: 1611-1614.
- Onoderalore, A.C. and Shahidi, F. 1996. Protein dispersions and hydrolysates from shark (*Isurus oxyrinchus*). *J. Aquat. Food Prod. Technol.* 5(4): 43-59.
- Osawa, T. 1994. Novel natural antioxidants for utilization in food and biological system. *In Postharvest biochemical of plant food-material in the tropics.* (Uritani, I. *et al.*, eds.). p. 241-251. Japan Scientific Societies Press. Japan.
- Pan, B.S. 1990. Recovery of shrimp waste for flavorant. *In Advances in Fisheries Technology and Biotechnology for Increased Profitability.* (Voigt, M. N. and Botta, J. R. eds.), pp. 437-447. Basel, Switzerland: Technomic Pub.
- Pena-Ramos, E.A. and Xiong, Y.L. 2003. Whey and soy protein hydrolysates inhibit lipid oxidation in cooked pork patties. *Meat Sci.* 64: 259-263.

- Peterson, B.R. 1981. The impact of the enzymatic hydrolysis process on recovery and use of proteins. *In* Enzymes and food processing. (Birch, G.G. *et al.*, eds), p. 269-299. Elsevier Applied Science Publishers. London, UK.
- Peterson, M.S. 1978. Fish protein concentrate. *In* Encyclopedia of food science (Peterson, M.S. and Johnson, A.H. eds.), pp. 266-271. The Avi publishing company, Inc. Connecticut
- Pinho, O., Ferreira, I.M.P.L.V.O., Oiveira, M.B.P.P. and Ferreira, M.A. 2000. Quantification of synthetic phenolic antioxidants in liver pates. *Food Chem.* 68: 353-357.
- Pokorny, J. 1999. Antioxidants in food preservation. *In* Handbook of food preservation. (Shafiur, R.M. ed.), p. 309-337. Marcel Dekker. New York.
- Pokorny, J. and Schmidt, S. 2001. Natural antioxidant functionality during food processing. *In* Antioxidant in food: Practical applications. (Pokorny, J., Yanishlieva, N. and Gordon, M. eds.), pp. 331-351. Woodhead Publishing. Abington, England.
- Pokorny, J. and Korczak, J. 2001. Preparation of natural antioxidants. *In* Antioxidant in food: Practical applications. (Pokorny, J., Yanishlieva, N. and Gordon, M. eds.), pp.311-330. Woodhead Publishing. Abington, England.
- Pomeranz, Y. 1991. Water. *In* Functional properties of food components. New York: Academic Press.
- Pour-El, A. 1981. Protein functionality: classification, definition, and methodology. *Protein Funct. Foods.* 147: 1-19.
- Pryor, W.A. and Porter, N.A. 1990. Suggested mechanisms for the production of 4-hydroxy-2-nonenal from the autoxidation of polyunsaturated fatty acids. *Free Rad. Biol. Med.* 8: 541-543.
- Puerta, T. 1999. Inhibition of leukocytes lipoxygenase by phenolics from virgin olive oil. *Biochem. Pharm.* 57: 448-449.
- Quaglia, G.B. and Orban, E. 1990. Influence of enzymatic hydrolysis on structure and emulsifying properties of sardine (*Sardina pilchardus*) protein hydrolysates. *J. Food Sci.* 55: 1571-1619.
- Rajalakshmi, D. and Narasimhan, S. 1996. Food antioxidants: sources and method of evaluation. *In* Food antioxidants. (Madhavi, D.L. *et al.*, eds.). p. 65-158. Marcel Dekker. New York.

- Rajapakse, N., Mendis, E., Jung, W.K., Je, J.Y. and Kim, S.K. 2005. Purification of a radical scavenging peptide from fermented mussel sauce and its antioxidant properties. *Food Res. Int.* 38: 175-182.
- Rebeca, B.D., Penavera, M.T. and Diazcastaneda, M. 1991. Production of fish protein hydrolysates with bacterial proteases yield and nutritional value. *J. Food Sci.* 56: 309–314.
- Rice-Evans, C. A., Miller, N. J. and Paganga, G. 1996. Structure antioxidant activity relationships of flavonoids and phenolic acids. *Free Radi. Biol. Med.* 20: 933–956.
- Riisom, T., Sims, R.J. and Fiorti, J.A. 1980. Effect of amino acids on the autoxidation of safflower oil in emulsions. *J. Am. Oil Chem. Soc.* 57: 351–359.
- Ruttanapornvareesakul, Y., Ikeda, M., Hara, K., Osako, K., Kongpun, O. and Nozaki, Y. 2005. Effect of shrimp head protein hydrolysates on the state of water and denaturation of fish myofibrils during dehydration. *Fish. Sci.* 71: 220-228.
- Ruttanapornvareesakul, Y., Ikeda, M., Hara, K., Osatomi, K., Osako, K., Kongpun, O. and Nozaki, Y. 2006. Concentration-dependent suppressive effect of shrimp head protein hydrolysate on denaturation-induced denaturation of lizardfish myofibrils. *Biores. Technol.* 97: 762-769.
- Sachindra, N.M., Bhaskar, N. and Mahendrakar, N.S. 2006. Recovery of carotenoids from shrimp waste in organic solvents. *Waste Manage.* 26: 1092–1098.
- Sakanaka, S., Tachibana, Y., Ishihara, N. and Juneja, L.R. 2004. Antioxidant activity of egg-yolk protein hydrolysates in a linoleic acid oxidation system. *Food Chem.* 86: 99–103.
- Sang, W. and Jin, Z.T. 2004. Lipid oxidation of fish liver oil affected by light, antioxidants and temperature. *J. Food Process. Preserv.* 28: 1-10.
- Sathivel, S., Bechtel, P., Babbitt, J., Smiley, S., Crapro, C., Reppond, K. and Prinyawiwatkul, W. 2003. Biochemical and functional properties of herring (*Clupea harengus*) byproduct hydrolysates. *J. Food Sci.* 68: 2196-2200.
- Seymour, T.A., Li, S.J. and Morrissey, M.T. 1996. Characterization of a natural antioxidant from shrimp shell waste. *J. Agric. Food Chem.* 44: 682-685.
- Shahidi, F. 1994. Proteins from seafood processing discards. *In* Seafood proteins (Sikorski Z.E., Pan, B.S. and Shahidi, F. eds.), p. 178. Chapman and Hall, Inc. New York.

- Shahidi, F. and Amarowicz, R. 1996. Antioxidant activity of protein hydrolysates from aquatic species. *J. Am. Oil Chem. Soc.* 73, 1197-1199
- Shahidi, F., Arachchi, J.K.V. and Jeon, Y. J. 1999. Food applications of chitin and chitosans. *Trends in Food Sci. Technol.* 10: 37-51.
- Shahidi, F., Han, X.Q. and Synowiecki, J. 1995. Production and characteristics of protein hydrolysates from capelin (*Mallotus villosus*). *Food Chem.* 53: 285-293.
- Shahidi, F., Synowiecki, J. and Naczk. 1992. Utilization of shellfish processing discards. *In* *Seafood Science and Technology*. (Graham, B. E. ed.), pp. 300-304. Fishing News Books. Canada.
- Shahidi, F. and Wanasundara, P.K.J.P.D. 1992. Phenolic antioxidants. *Crit. Rev. Food Sci. Nutr.* 32: 67-103.
- Siddhuraju, P. and Becker, K. 2007. The antioxidant and free radical scavenging activities of processed cowpea (*Vigna unguiculata (L.) walp.*) seed extracts. *Food Chem.* 101: 10-19.
- Sies, H., Stahl, W. and Sundquist, A. 1992. Antioxidant functions of vitamins, vitamin E and C, beta-carotene and other carotenoids. *Ann. NY Acad. Sci.* 368: 7-19.
- Sikorski, Z.E., Kolakowska, A. and Pan, B.S. 1990. The nutritive composition of the major groups of marine food organisms. *In* *Seafood: resources, nutritional composition and preservation* (Z.E. Sikoski, ed.), pp. 29-54. CRC Press. Florida.
- Silva, A.S., Hernandez, J.L. and Losada, P.P. 2004. Modified atmosphere packaging and temperature effect on potato crisps oxidation during storage. *Anal. Chem Acta.* 524: 185-189.
- Šližyte, R., Daukšas, E., Falch, E., Storrø1, I. and Rustad, T. 2005a. Yield and composition of different fractions obtained after enzymatic hydrolysis of cod (*Gadus morhua*) by-products. *Process Biochem.* 40: 1415-1424.
- Šližyte, R., Daukšas, E., Falch, E., Storrø1, I. and Rustad, T. 2005b. Characteristics of protein fractions generated from hydrolysed cod (*Gadus morhua*) by-products. *Process Biochem.* 40: 2021-2033.
- Spinelli, J., Lehman, L. and Wieg, D. 1974. Composition, processing and utilization of red crab (*Pleuroncoides planipes*). *J. Fish. Res. Board Can.* 31: 1025-1029.

- Sriket, P., Benjakul, S., Visessanguan, W. and Kijroongrojana, K. 2006. Comparative studies on chemical composition and thermal properties of black tiger shrimp (*Penaeus monodon*) and white shrimp (*Penaeus vannamei*) meats. Food Chem. 103: 1199-1207.
- Stallings, C. W., Kroa, B. A., Carroll, R. T., Mertzner, A.L. and Funk, M.O. 1990. Crystallization and preliminary x-ray characterization of a soybean seed lipoxygenase. J. Mol. Biol. 211, 685-687.
- Stanhke, L.H. 1995. Dried sausages fermented with *Staphylococcus xylosus* at different temperatures and different ingredient levels-part II volatile components. Meat Sci. 41: 193-209.
- Steel, R.G.D. and Torrie, J.H. 1980. Principle and procedure of statistics. 2<sup>nd</sup> ed. New York: McGraw-Hill.
- Sudaryono, A., Hoxey, M.J., Kailis, S.G. and Evans, L.H. 1995. Investigation of alternative protein sources in practical diets for juvenile shrimp, *Penaeus monodon*. Aquaculture. 134: 313-323.
- Sudaryono, A., Tsvetnenko, E. and Evans, L.H. 1996. Digestibility studies on fisheries by-product based diets for *Penaeus monodon*. Aquaculture. 143: 331-340.
- Suetsuna, K., Ukeda, H. and Ochi, H. 2000. Isolation and characterization of free radical scavenging activities peptides derived from casein. J. Nutr. Biochem. 11: 128-131.
- Suetsuna, K. 2000. Antioxidant peptides from the protease digest of prawn (*Penaeus japonicus*) muscle. Marine Biotechnol. 2: 5-10.
- Sun, B.X. and Fukuhara, M. 1997. Effects of coadministration of butylated hydroxytoluene, butylated hydroxyanisole and flavonoids on the activation of mutagens and drug-metabolizing enzymes in mice. Toxicol. 122: 61-72.
- Suphamongkhon, K. 2002. Office of the Thai trade representative. Thai trade representative Thailand's Policy on international Trade and Investment after the 1997 Economic Crisis for the International Boston Seafood Show, Massachusetts Boston, 12 March.
- Taga, M.S., Miller, E. E. and Pratt, D. E. 1984. Chaiaseeds as source of natural lipid antioxidant. J. Am. Oil Chem. Soc. 61: 928-931.



- Tang, S., Kerry, J.P., Sheedan, D., Buckley, D.J. and Morrissey, P.A. 2001. Antioxidative effect of added tea catechin on susceptibility of cooked red meat, poultry and fish patties to lipid oxidation. *Food. Res. Intern.* 34: 651-657.
- Taylor, M.J. and Richardson, T. 1980. Antioxidant activity of cysteine and protein sulfhydryls in a linoleate emulsion oxidized by hemoglobin. *J. Food Sci.* 45:1223–1227, 1230.
- TCPS. 324. 2004. Thai community Product standard. Thai industrial standard institute. Ministry of industry, Bangkok, Thailand.
- Thai Industrial Standard. 1983. Local fish sauce standard. Department of industry, Bangkok, Thailand.
- Thiansilakul, Y., Benjakul, S. and Shahidi, F. 2007a. Antioxidative activity of protein hydrolysate from round scad muscle using Alcalase and Flavourzyme. *J. Food Biochem.* 31: 266-287.
- Thiansilakul, Y., Benjakul, S. and Shahidi, F. 2007b. Compositions, functional properties and antioxidative activity of protein hydrolysates prepared from round scad (*Decapterus maruadsi*). *Food Chemistry.* 103: 1385-1394.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L. and Byrne, D.H. 2006. Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *J. Food Comp. Anal.* 19: 669-675.
- Thanonkaew, A., Benjakul, S. and Visessanguan, W. 2006. Chemical composition and thermal property of cuttlefish (*Sepia pharaonis*) muscle. *J. Food Comp. Anal.* 19: 127-133.
- Trung, T.S., Thein-Han, W.W., Qui, N.T., Ng, C.H. and Stevens, W.F. 2006. Functional characteristics of shrimp chitosan and its membranes as affected by the degree of deacetylation. *Biores. Technol.* 97: 659-663.
- Turgeon, S.L., Gautier, S.F. and Paquin, P. 1992. Emulsifying property of whey peptide fractions as a function of pH and ionic strength. *J. Food Sci.* 57: 601-634.
- Ukeda, H., Matsuda, H., Osajima, K., Matsufuji, H., Matsui, T. and Osajima, Y. 1992. Peptides from peptic hydrolysate of heated sardine meat that inhibit angiotensin I-converting enzyme. *Nippon Nogeikagaku Kaishii.* 66: 25-29.
- Venugopal, V. and Shahidi, F. 1994. Thermostable water dispersions of myofibrillar proteins from Atlantic mackerel (*Scomber scombrus*). *J. Food Sci.* 59: 265-268.

- Venugopal, V. and Shahidi, F. 1995. Value-added products from underutilized fish species. *Crit. Rev. Food Sci. Nutr.* 35: 431–453.
- Viera, G.H.F., Martin, A.M., Saker-Sampaio, S., Omar, S. and Goncalves, R.C.F. 1995. Studies on the enzymatic hydrolysis of Brazilian lobster (*Panulirus* spp.) processing wastes. *J. Sci. Food Agric.* 69: 61-65.
- Weber, K.C., Honório, K.M., Bruni, A.T. and Albérico, B.F.S. 2006. The use of classification methods for modeling the antioxidant activity of flavonoid compounds. *J. Mol. Model.* 12: 915-920.
- Webster, J.D., Ledward, D.A. and Lawrie, R.A. 1982. Protein hydrolysates from meat industry by-products. *Meat Sci.* 7: 147-157.
- Weska, R.F., Moura, J.M., Batista, L.M., Rizzi, J. and Pinto, L.A.A. 2007. Optimization of deacetylation in the production of chitosan from shrimp wastes: Use of response surface methodology. *J. Food Eng.* 80: 749-753.
- Wettasinghe, M. and Shahidi, F. 1999. Antioxidant and free radical scavenging properties of ethanolic extracts of defatted borage (*Borago officinalis* L.) seeds. *Food Chem.* 67: 399–414.
- Wijewickreme, A.N., Kitts, D.D. and Durance, T.D. 1997. Reaction conditions influence the elementary composition and metal chelating affinity of nondialyzable model Maillard reaction products. *J. Agric. Food Chem.* 45: 4577-4583.
- Wilcox, A.L. and Marnett, L.J. 1993. Polyunsaturated fatty acid alkoxyl radicals exist as carbon-centered epoxyallylic radicals: a key step in hydroperoxide-amplified lipid peroxidation. *Chem. Res. Toxicol.* 6: 413-416.
- Wilde, P.J. and Clark, D.C. 1996. Foam formation and stability. *In* methods for testing protein functionality. (Hall, G.M., ed.). p. 110-148. Blackie Academic and Professional. London, UK.
- Wilding, P., Lilliford, P.J. and Regenstein, J.M. 1984. Functional properties of proteins in foods. *J. Chem. Technol. Biotechnol.* 34B: 182–189.
- Windsor, M.L. 2001. Fish protein concentrate. (online). Search from: <http://www.Fao.org/waidocs/tan/.htm> [11 September 2007]

- Wu, H.C., Chen, H.M. and Shiau, C.Y. 2003. Free amino acid and peptide as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). Food Res. Int. 36: 949-957.
- Yamaguchi, N., Yokoo, Y. and Fujimaki, M. 1975. Studies on antioxidative activities of amino compounds on fats and oils. Part II. Antioxidative activities of dipeptides and their synergistic effects of tocopherol. Nippon Shokuhin Kogyo Gakkaishi 22, 425-430.
- Yanar, Y. and Celik, M. 2005. Seasonal variations of fatty acid composition in wild marine shrimps (*Penaeus semisulcatus* De Haan, 1844 and *Metapenaeus monoceros* Fabricus, 1789) from the eastern Mediterranean Sea. Food Sci. Technol. Int. 11: 391-395.
- Yanishlieva-Maslarova, N.Y. 2001. Inhibiting oxidation. In Antioxidants in food: practical applications. (Pokony, J., Yanishlieva, N. and Gordon, M. eds.), pp. 22-70. Woodhead Publishing. Abington, England.
- Yoshimura, Y., Iijima, T., Watanabe, T. and Nakazawa, H. 1997. Antioxidative effect of Maillard reaction products using glucose-glycine model system. J. Agric. Food Chem. 45: 4106-4109.