

REFERENCES

- Aaslyng, D.M., Bejerholm, C., Ertbjerg, P., Bertram, C.H. and Andersen, J.H. 2003. Cooking loss and juiciness of pork in relation to raw meat quality and cooking procedure. *Food Qual. Pref.* 14: 277-288.
- Adacghi, K., Hirata, T., Nagai, K. and Sakaguchi, M. 2001. Hemocyanin a most likely inducer of black spots in kuruma prawn *Penaeus japonicus* during storage. *J. Food Sci.* 66: 1130-1136.
- Akiyama, D.M., Dominy, W.G. and Lawrence, A.L. 1992. Penaeid shrimp nutrition. In: *Marine shrimp culture: principles and practices* (Fast, A.W. and Lester, L.J. eds.). p. 535–568. Elsevier. Amsterdam.
- Almas, K.A. 1981. Chemistry and microbiology of fish processing. Dept. of Biochemistry, Norwegian Inst. of Technol., Univ. of Trondheim. Norway.
- Ang, J.I.T.F. and Hultin, H.O. 1989. Denaturation of cod myosin during freezing after modification with formaldehyde. *J. Food Sci.* 54: 814-818.
- AOAC. 1999. *Official Method of Analysis*, 16th ed. Association of Official Analytical Chemists. Washington, DC.
- Arai, K. 1966. Acid soluble nucleotide in muscle of marine invertebrates. *Bull. Jpn. Soc. Sci. Fish.* 32: 174-180.
- Arai, S. and Fujimaki, M. 1991. Modification with proteolytic enzymes. In *Food enzymology* (Fox, P.F. ed.). p. 86-87. Chapman & Hall. New York.
- Armenta-Lopez, R., Guerrero, L. and Huerta, S. 2002. Astaxanthin extraction from shrimp waste by lactic fermentation and enzymatic hydrolysis of the carotenoprotein complex. *J. Food Sci.* 67: 1002-1006.
- Ashie, I.N.A. and Simpson, B.K. 1997. Purification and characterization of cathepsin L from hepatopancreas of carp *Cyprinus carpio*. *Comp. Bichem. Physiol.* 115B: 531-537.
- Aubourg, S.P., Pineiro, C., Gallardo, J.M. and Velazquez, J.B. 2004. Biochemical changes and quality loss during chilled storage of farmed turbot (*Psetta maxima*). *Food Chem.* 90: 445-452.

- Baden, S.P., Depledge, M.H. and Hagerman, L. 1994. Glycogen depletion and altered copper and manganese handling in *Nephrops norvegicus* following starvation and exposure to hypoxia. Mar. Ecol. Prog. Ser. 103: 65-72.
- Badii, F. and Howell, N.K. 2001. A comparison of biochemical changes in cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*) fillets during frozen storage. J. Sci. Food Agric. 82: 87-97.
- Badii, F. and Howell, N.K. 2002. Changes in texture of cod and haddock fillets during frozen storage. Food Hydrocol. 16: 313-319.
- Ban, M. Ushio, H. and Yamanaka, H. 1998. Postmortem biochemical changes in muscle of *Buccinum striatissimum* during storage. Fish. Sci. 64: 482-486.
- Beas, V.E., Wagner, J.R., Crupkin, M. and Anon, M.C. 1990. Thermal denaturation of hake (*Merluccius hubbsi*) myofibrillar proteins. A differential scanning calorimetric and electrophoretic study. J. Food Sci. 55: 683-687.
- Benjakul, S. and Bauer, F. 2000. Physicochemical and enzymatic changes of cod muscle proteins subjected to different freeze-thaw cycles. J. Sci. Food Agric. 80: 1143-1150.
- Benjakul, S. and Bauer, F. 2001. Biochemical and physicochemical changes in catfish (*Silurus glanis* Linne) muscle as influenced by different freeze-thaw cycles. Food Chem. 72: 207-217.
- Benjakul, S., Seymour, T.S., Morrissey, M.T. and An, H. 1997. Physicochemical changes in Pacific whiting muscle proteins during iced storage. J. Food Sci. 62: 729-733.
- Benjakul, S., Visessanguan, W., Riebroy, S., Ishizaki, S. and Tanaka, M. 2002. Gel-forming properties of surimi produced from bigeye snapper, *Priacanthus tayenus* and *P. macracanthus*, stored in ice. J. Sci. Food Agric. 82: 1442-1451.
- Benjakul, S., Visessanguan, W., Thongkaew, C. and Tanaka, M. 2003. Comparative study on physicochemical changes of muscle proteins from some tropical fish during frozen storage. Food Res. Intern. 36: 787-795.
- Benner, R.A., Staruszkiewicz, J.R., Rogers, P.L. and Otwell, W.S. 2003. Evaluation of putrescine, cadaverine, and indole as chemical indicators of decomposition in Penaeid shrimp. J. Food Sci. 68: 2178-2185.

- Bergman, I. and Loxley, R. 1963. Two improved and simplified methods of the spectrophotometric determination of hydroxyproline. *Anal. Chem.* 35: 1961-1965.
- Bishop, J.B. and Burton, R.S. 1993. Amino acid synthesis during hyperosmotic stress in *Penaeus aztecus* postlarvae. *Comp. Biochem. Physiol.* 106: 49-56.
- Bjeldanes, L.F., Shutz, D.E. and Morris, M.M. 1978. Etiology of scombroid poisoning-cadaverine potentiation of histamine toxicity in guinea pigs. *Food Cosmet. Toxicol.* 16: 157-162.
- Bligh, E.G. and Dyer, W.J. 1959. Lipid extraction from fish muscle. *Can. J. Biochem. Physiol.* 37: 911-931.
- Boon, D.D. 1975. Discoloration in processed crabmeat: a review. *J. Food Sci.* 40:756-61.
- Boonyaratpalin, M., Thongrod, S., Supamattaya, K., et al. 2001. Effects of β -carotene source, *Dunaliella salina*, and astaxanthin on pigmentation, growth, survival and health of *Penaeus monodon*. *Aquac. Res.* 32: 182-190.
- Botta, J.R., Bonnell, G. and Squires, B.E. 1987. Effect of method of catching and time of season on sensory quality of fresh raw Atlantic cod (*Gadus morhua*). *J. Food Sci.* 52: 928-931.
- Bottino, N.R., Gennity, J., Lilly, M.L., Simmons, E. and Finne, G. 1980. Seasonal and nutritional effects on the fatty acids of three species of shrimp *Penaeus setiferus*, *P. aztecus* and *P. duoratum*. *Aqua.* 19: 139-145.
- Bottino, N.R., Lilly, M.L. and Finne, G. 1979. Fatty acid stability of Gulf of Mexico brown shrimp (*Penaeus aztecus*) held on ice and in frozen storage. *J. Food Sci.* 44: 1778-1779.
- Brauer, J.M.E., Leyva, J.A.S., Alvarado, L.B. and Sandez, O.R. 2003. Effect of dietary protein on muscle collagen, collagenase and shear force of farmed white shrimp (*Litopenaeus vannamei*). *Eur. Food Res. Technol.* 217: 277-280.
- Bredie, W.L.P. and De Boer, E. 1992. Evaluation of the MPN, Anderson- Baird- Parker, Petrifilm *E. coli* and Fluorocult ECD method for enumeration of *Escherichia coli* in foods of animal origin. *J. Food Microbiol.* 161: 197-208.
- Britton G. 1996. Carotenoids. In: Natural food colorants. 2nd ed. (Hendry, G.A.F and Houghton J.D, eds.) p. 197-243. New York: Blackie Academic and Professional.

- Buchanan, R.L. 1991. Microbiological criteria for cooked ready-to eat shrimp and crab meat. *Food Technol.* 45(4): 157-160.
- Busconi, L., Folco, E.J., Martone, C.B. and Sanchez, J.J. 1989. Post mortem changes in cytoskeletal elements of fish muscle. *J. Food Biochem.* 13: 443-451.
- Buttkus, H. 1974. On the nature of the chemical and physical bonds which contribute to some structural properties of protein foods: a hypothesis. *J. Food Sci.* 39: 484-489.
- Cano-López, A., Simpson, B.K. and Haard, N.F. 1987. Extraction of carotenoprotein from shrimp process wastes with the aid of trypsin from Atlantic cod. *J. Food Sci.* 52: 503-506.
- Castell, C.H., Maclean, J. and Moore, B 1965. Rancidity in lean fish muscle. IV. Effect of sodium chloride and other salts. *J. Fish. Res. Board. Can.* 22: 929-944.
- Catacutan, M.R., 1991. Growth and fatty acid composition of *Penaeus monodon* juveniles fed various lipids. *Isr. J. Aquac. Bamidgeh.* 43: 47–56.
- Chaitiamvong, S. and Supongpan, M., 1992. A Guide to Penaeoid Shrimps Found in Thai Waters. Australian Institute of Marine Science, Townsville, Australia.
- Chanmugam, P., Donovan, J., Wheeler, C.J. and Hwang, D.H. 1983. Differences in the lipid composition of fresh water prawn (*Macrobrachium rosenbergii*) and marine shrimp. *J. Food Sci.* 48: 1440-1462.
- Chen, H.M. and Meyers, S.P. 1983. Ensilage treatment of crawfish waste for improvement of astaxanthin pigment extraction. *J. Food Sci.* 48: 1516-1521.
- Cheng, K., Hu, C., Liu, Y., Zheng, S. and Qi, X. 2006. Effect of dietary calcium, phosphorus and calcium/phosphorus ratio on the growth and tissue mineralization of *Litopenaeus vannamei* reared in low-salinity water. *Aquaculture.* 251: 472-483.
- Cheuk, W., Finne, G., and Nickelson II, R. 1979. Stability of adenosine deaminase and adenosine monophosphate deaminase during ice storage of pink and brown shrimp from the Gulf of Mexico. *J. Food Sci.* 44: 1625- 1628.
- Chia, S.S., Baker, R.C. and Hotchkiss, J.H. 1983. Quality comparison of thermoprocessed fishery products in cans and retortable pouches. *J. Food Sci.* 48: 1521–1531.
- Chien, Y.H. and Jeng, S.C. 1992. Pigmentation of kuruma prawn, *Penaeus japonicus* Bate, by various pigment sources and levels and feeding regimes. *Aquaculture.* 102: 333- 346.

- Chinivasagam, H. N., Bremner, H. A. and Reeves, R. 1998. Can spoilage bacteria cause black spot (melanosis) in stored prawns. *Lett. Appl. Microbiol.* 27: 5-8.
- Cobb, B.F., Vanderzant, C. Thompson, C.A. and Custer, C.S. 1973. Chemical characteristics, bacterial counts and potential shelf-life of shrimp from various locations on the northwestern Gulf of Mexico. *J. Milk Food Technol.* 36: 463-468.
- Connell, J.J. 1964. Fish muscle proteins and some effects on them of processing. In *Symposium on Foods: proteins and their reaction*. (Shultz, H.W. and Anglemeier A.F. ed.). p. 255. AVI Publ. Co. Westport, CT.
- Connell, J.J. 1995. Quality deterioration and extrinsic quality defects in raw material. In *Control of Fish Quality*. Fishing News Books Ltd. Survey, England.
- Crawford, D.L. 1981. Meat yield and shell removal functions of shrimp processing. Oregon State University Extension Marine Advisory Program A Land Grant/Sea Grant Cooperative Special Report 597. Oregon State.
- Cuzon, G. and Aquacop. 1998. Nutritional review of *Penaeus stylirostris*. *Rev. Fish. Sci.* 6: 129-141.
- Dall, W. 1965. Studies on the physiology of a shrimp, *Metapenaeus* sp. (Crustacea: Decapoda: Penaeidae). III. Composition and stucture of the integument. *Austr. J. Mar. Fresh. Res.* 16: 13 – 24.
- Davies, J.R., Bardsley, R.O., Ledward, D.A. and Poulter, R.G. 1988. Myosin thermal stability in fish muscle. *J. Sci. Food Agric.* 45: 61-68.
- Davis, D.A. and Gatlin III, D.M. 1996. Dietary mineral requirements of fish and marine crustaceans. *Rev. Fish. Sci.* 4: 75-99.
- Decker, E.A. and Hultin, O.H. 1992. Lipid oxidation in muscle food via redox: iron. In *Lipid oxidation in food* (Angelo, A.S. ed.). p. 33–54.: American Chemical Society. Washington, DC.
- Decker, E.A., Xiong, Y.L., Calvert, J.T., Crum, A.D. and Blanchard, S.P. 1993. Chemical, physical and functional properties of oxidized turkey white muscle myofibrillar proteins. *J. Agric. Food Chem.* 41: 186-189.
- Decker, H. and Tuczak, F. 2000. Tyrosinase/catecholoxidase activity of hemocyanins: structural basis and molecular mechanism. *Trends Biochem. Sci.* 25: 392-397.

- Eckhoff, M.K., Aidos, I., Hemre, G.I. and Lie, O. 1998. Collagen content in farmed Atlantic salmon (*Salmo salar*, L.) and subsequent changes in solubility during storage on ice. *Food Chem.* 62: 197-200.
- Ehira, S. 1976. Biochemical study on the freshness of fish. *Bull. Tokai Reg. Fish. Res. Lab.* 88: 1-5.
- Ellman, G.L. 1959. Tissue sulphydryl groups. *Arch. Biochem. Biophys.* 82: 70-77.
- Erdogdu, F., Balaban, M.O., Otwell, W.S. and Garrido, L. 2004. Cook-related yield loss for Pacific white (*Penaeus vannamei*) shrimp previously treated with phosphates: effects of shrimp size and internal temperature distribution. *J. Food Eng.* 64: 297-300.
- Espe, M., Ruohonen, K., Bjornevik, M., Froyland, L., Nortvedt, R. and Kiessling, A. 2004. Interactions between ice storage time, collagen composition, gaping and textural properties in farmed salmon muscle harvested at different times of the year. *Aquaculture*. 240: 498-504.
- Fatima, R., Farooqui, B. and Qadri, R.B. 1981. Inosine monophosphate and hypoxanthine as indices of quality of shrimp (*Penaeus morguensis*). *J. Food Sci.* 46: 1125-1127.
- Fatima, R., Khan, M.A. and Qadri, R.B. 1988. Shelf-life of shrimp (*Penaeus merguiensis*) stored in ice (0°C) and partially frozen (-3°C). *J. Sci. Food Agric.* 42: 235-247.
- Feliz, G.L.A., Gatlin, M.D., Lawrence, L.A. and Velazquez, P.M. 2002. Effect of dietary phospholipid on essential fatty acid requirements and tissue lipid composition of *Litopenaeus vannamei* juveniles. *Aquaculture*. 207: 151-167.
- Feng, P.C.S. and Hartman, P.A. 1982. Fluorogenic assays for immediate confirmation of *Escherichia coli*. *Appl. Envir. Microbiol.* 43: 1320– 1329.
- Fernandez-salguero, J. and Mackie, I.M. 1987. Comparative rates of spoilage of fillet and whole fish during storage of haddock (*Melanogrammus aeglefinus*) and herring (*Clupea harengus*) as determined by the formation of non-volatile and volatile amines. *Intern. J. Food Sci. Technol.* 22: 385-390.
- Fine, G. 1992. Non-protein nitrogen compounds in fish and shellfish. In *Advances in seafood biochemistry: composition and quality*. (Flick, G.J., Jr. and Martin, R.E eds.). p. 393-401. Technomic Pub. Co. Inc. Lancaster, Basel.

- Fiske, C.H. and Subbarow, Y. 1925. The colorimetric determination of phosphorus. *J. Biol. Chem.* 66: 375–400.
- Foegeding, E.A., Allen, C.E. and Dayton, W.R. 1986. Effect of heating rate on thermally formed myosin, fibrinogen and albumin gels. *J. Food Sci.* 52: 104-112.
- Foegeding, E.A., Lanier, T.C. and Hultin, H.O. 1996. Characteristics of edible muscle tissue. In *Food Chemistry*. 3rd (Fennema, O.R., ed.). p. 875-942. Marcel Dekker, Inc. New York.
- Fuke, S. 1994. Taste active components of seafood with special reference to umami substances. In *Seafoods: chemistry, processing technology and quality* (Shahidi, F. and Botta, J.R., eds.). p. 125-128. Blackie Academic & Professional, London.
- Fukuda, Y., Kakehata, K. and Arai, K. 1981. Denaturation of myofibrillar protein in deep-sea fish by freezing and storage. *Bull. Jpn. Soc. Sci. Fish.* 47: 663–672.
- Gardner, H.W. 1979. Lipid hydroperoxide reactivity with proteins and amino acid: a review. *J. Agric. Food Chem.* 27: 220-229.
- Gill, T.A. 1990. Objective analysis of seafood quality. *Food Rev. Intern.* 6: 681-714.
- Gill, T.A. 1992. Biochemical and chemical indices of seafood quality. In *Quality assurance in the fish industry. Proceeding of an international conference*. p. 377-388.
- Glencross, B.D., Smith, D.M., 2001. Optimising the essential fatty acids, eicosapentaenoic and docosahexaenoic acids in the diet of the prawn, *Penaeus monodon*. *Aquac. Nutr.* 7: 59-69.
- Glencross, B.D., Smith, D.M., Thomas, M.R. and Williams, K.C. 2002. Optimising the essential fatty acids in the diet for weight gain of the prawn, *Penaeus monodon*. *Aquaculture*. 204: 85–99.
- Gokodlu, N., Ozden, O. and Erkan, N. 1998. Physical, chemical and sensory analyses of freshly harvested sardines (*Sardina pilchardus*) stored at 4°C. *Can. J. Aquat. Food Prod. Technol.* 7: 5-15.
- Gómez-Guillén, M.C., Montero, P., Hurtado, O. and Borderias, A.J., 2000. Biological characteristics affect the quality of farmed Atlantic salmon and smoked muscle. *J. Food Sci.* 65: 53-60.

- Gong, H., Lawrence, A.L., Jiang, D.H. and Gatlin, III, D.M. 2000. Lipid nutrition of juvenile *Litopenaeus vannamei*: I. Dietary cholesterol and de-oiled soy lecithin requirements and their interaction. *Aquaculture*. 190: 307– 326.
- Gram, L., Huss, H.H. 1996. Microbial spoilage of fish and fish products. *Intern. J. Food Microbiol.* 33: 121–137.
- Gray, J.L. and Pearson, A.M. 1994. Lipid-derived off-flavours in meat formation and inhibition. In *Flavour of meat and meat products* (Shahidi, F., ed.). p. 116–143. London: Blackie Academic.
- Gray, J.L., Gomaa, E.A. and Buckley, D.J. 1996. Oxidative quality and shelf-life of meats. *Meat Sci.* 43: 11–123.
- Gruschczyk, B. and Kamp, G. 1989. The shift from glycogenolysis to glycogen resynthesis after escape swimming: studies on the abdominal muscle of the shrimp, *Crangon crangon*. *J. Comp. Physiol. B: Biochem. Syst. Envir. Physiol.* 753 – 760.
- Hale, M.B. and Waters, M.E. 1981. Frozen storage stability of whole and headless freshwater prawns *Machrobrachium rosenbergii*. *Mar. Fish. Rev.* 42: 18-21.
- Hamm, R. 1979. Delocalization of mitochondrial enzymes during freezing and thawing of skeletal muscle. In *Proteins at low temperatures. Advances in Chemistry Series* (vol. 180) (Fennema, O.R., ed.). pp. 191. American Chemical Society. Washington DC.
- Hansen M.E. and Llanes, A. 1994. Applications of crustacean wastes in biotechnology. In *Fisheries processing: biotechnological applications* (Martin A.M., ed.). p. 174-205. Chapman & Hall. London.
- Hashimoto, K., Watabe, S., Kono, M. And Skiro, K. 1979. Muscle protein composition of sardine and mackerel. *Bull. Jpn. Soc. Sci. Fish.* 45: 1435-1441.
- Hasting, R.J., Rodger, G.W., Park, R., Matthews, A.D. and Anderson, E.M. 1985. Differential scanning colorimetry of fish muscle: the effect of processing and species variation. *J. Food Sci.* 50: 1503-1510.
- Hatae, K., Tobimatsu, A., Takeyama, M. and Matsumoto, J.J. 1986. Contribution of the connective tissues on the texture difference of various fish species. *Bull. Jpn. Soc. Sci. Fish.* 52: 2001-2007.

- Hayakawa, S. and Nakai, S. 1895. Contribution of hydrophobicity, net charge and sulphydryl groups to thermal properties of ovalbumin. *Can. Inst. Food Sci. Technol.* 18: 190-195.
- Heymann, H., Hedrick, H.B., Karrasch, M.A., Eggeman, M.K. and Ellersieck, M.R. 1990. Sensory and chemical characteristics of fresh pork roasts cooked to different centre temperatures. *J. Food Sci.* 55: 613-617.
- Hsieh, R.J. and Kinsella, J.E. 1989. Lipoxygenase generates specific volatile carbonyl compounds in fish tissues. *J. Agric. Food Chem.* 37: 279-286.
- Hsieh, Y.L. and Regenstein, J.M. 1989. Texture changes of frozen stored cod and ocean perch minces. *J. Food Sci.* 54: 824-826.
- Huidobro, A. and Tejada, M. 1994. Emulsifying capacity of fish mince from several species during frozen storage. *J. Sci. Food Agric.* 61: 333-338.
- Huidobro, A., López-Caballero, E.M. and Mendes, R. 2002. Onboard processing of deepwater pink shrimp (*Parapenaeus longirostris*) with liquid ice: Effect on quality. *Eur. Food Res. Technol.* 214: 469-475.
- Huidobro, A., Mendes, R. and Nunes, M. 2001. Slaughtering of gilthead seabream (*Sparus aurata*) in liquid ice: influence of fish quality. *Eur. Food Res. Technol.* 213: 267-272.
- Hultin, H.O. 1985. Characteristics of Muscle Tissue. In *Food Chemistry*. (Fennema, O.R., ed.). p. 725-789. Marcel Dekker. New York.
- Hultman, L. and Rustad, T. 2002. Textural changes during iced storage of salmon (*Salmo salar*) and cod (*Gadus morhua*). *J. Aquat. Food Prod. Technol.* 11(3-4): 105-123.
- Hutin, H.O. 1984. Postmortem Biochemistry of Meat and Fish. *J. Chem. Educ.* 61: 289-298.
- Ichihashi, H., Kohno, H., Kannan, K., Tsumura, A. and Yamasaki, S. 2001. Multielement analysis of purple back flying squid using high resolution inductively coupled plasma-mass spectrometry (HR ICP-MS). *Envir. Sci. Technol.* 35: 3103-3108.
- Ikeda, S. 1979. Other organic components and inorganic components. In *Advances in Fish Science and Technology* (Connell, J.J., ed.). p. 111-124. Fishing News Books. Surrey.

- International Commission on Microbiological Specifications for Foods (ICMSF). 1986. Microorganism in foods. 6: Microbiology of Food Commodities. Blackie Academic & Professional. Baltimore.
- Ishikawa, W., Miyake, Y. and Yasuie, S. 1966. Chromatophores and carotenoids in cultured kuruma prawn, *Penaeus japonicus* Bate, of different body colours. Bull. Fish. Exp. Stn. Okayama Prefecture. 5: 18–24.
- Iwamoto, M., Yamanaka, H. and Abe, H. 1990. Rigor-mortis and its temperature dependency in several marine fishes. Bull. Jpn. Soc. Sci. Fish. 56: 93-99.
- Iwasaki, M. and Harada, R. 1985. Proximate and amino acid composition of the roe and muscle of selected marine species. J. Food Sci. 50: 1585–1587.
- Jadhav, S.J., Nimbalker, S.S., Kulkami, A.D. and Madhavi, D.L. 1996. Lipid oxidation in biological and food systems. In Food antioxidants. (Madhavi, D.L. *et al.*, eds.). p. 5-64. Marcel Dekker. New York.
- Jessen, F. 1996. Freezing induced protein changes in fish evaluated by two-dimensional gel electrophoresis. In Refrigeration and aquaculture. p. 363-369. Nantes France: IFREMER.
- Jeyasekaran, G. and Ayyappan, S. 2002. Postharvest microbiology of farm-reared, tropical freshwater prawn (*Macrobrachium rosenbergii*). J. Food Sci. 67: 1859-1861.
- Jeyasekaran, G., Ganesan, P., Anadaraj. R., Jeya Shakila, R. and Sukumar, D. 2006. Quantitative and qualitative studies on the bacteriological quality of Indian white shrimp (*Penaeus indicus*) stored in dry ice. Food Microbiol. 23: 526-533.
- Jeyasekaran., G., Ganesan, P., Jeya Shakila, R. Maheswari, K. and Sukumar, D. 2004. Dry ice as a novel chilling medium along with water ice for short-term preservation of fish Emperor breams, lethrinus (*Lethrinus miniatus*). Innov. Food Sci. Emerg. Technol. 5: 485-493.
- Jeyaweera, V. and Subasinghe, S. 1988. Some chemical and microbiological changes during chilled storage of prawns (*Penaeus indicus*). FAO Fisheries Report No. 401. p. 19-22.

- Jiang, S.T., Wang, F.J. and Chen, C.S. 1989. Properties of actin and stability of the actomyosin reconstituted from milkfish (*Chanos chanos*) actin and myosin. *J. Agric. Food Chem.* 37: 1232-1235.
- Johansson, L., Ruderus, H., Beilby, R.I., 1992. Optimum internal temperature established by sensory evaluation for fish prepared in conventional and microwave ovens. *Home Econ. Res. J.* 21: 192-205.
- John, A. and Shashtri, P.N. 1998. Studies on food macromolecules by differential scanning colorimetry: a critical appraisal. *J. Food Sci. Technol.* 35: 1-14.
- Johnston, I.A. and Frearson, N.G. 1973. The effects of environmental temperature on properties of myofibrillar adenosine triphosphate from various species of fish. *J. Biochem.* 133: 735-738.
- Johnston, J.J., Ghanbari, H.A., Wheeler, W.B. and Kirk, J.R. 1988. Characterization of shrimp lipid. *J. Food Sci.* 48: 33-35.
- Jones, K.W. and Mandigo, R.W. 1982. Effect of chopping temperature on the microstructure of meat emulsions. *J. Food Sci.* 47: 1930-1934.
- Kamal, M., Rahman, M.M., Yasmin, L., Istam, M.N., Nurullah, M. and Mazid, M.A. 2000. Studies on the post-mortem changes in shrimp and prawn during ice storage: II. Biochemical aspects of quality changes. *Bangladesh J. Fish. Res.* 4: 91-96.
- Kanazawa, A., Teshima, S. and Ono, K. 1979a. Relationship between essential fatty acid requirements of aquatic animals and the capacity for bioconversion of linolenic acid to highly unsaturated fatty acids. *Comp. Biochem. Physiol.* 63B: 295– 298.
- Kanazawa, A., Teshima, S. and Tokiwa, S. 1979b. Biosynthesis of fatty acids from palmitic acid in the prawn, *Penaeus japonicus*. *Mem. Fac. Fish. Kagoshima Univ.* 28: 17– 20.
- Karakoltsidis, P.A., Zotos, A. and Constantinides, S.M. 1995. Composition of the commercially important Mediterranean finfish, crustaceans, and molluscs. *J. Food Comp. Anal.* 8: 258-273.
- Karel, M., Fennema, O.R. and Lund, D.B. 1975. Principles of food science. Vol. II. p. 485. Marcel Dekker. New York.

- Katayama, T., Hirata, K. and Chichester, C.O. 1971. The biosynthesis of astaxanthin: IV. The carotenoids in the prawn, *Penaeus Japonicus* Bate (Part I). Bull. Jpn. Soc. Sci. Fish. 37: 614–620.
- Katayama, T., Katama, T. and Chichester, C.O. 1972. The biosynthesis of astaxanthin: VI. The carotenoids in the prawn *Penaeus japonicus* Bate (Part II). Int. J. Biochem. 3: 363–368.
- Ke, P.J., Ackman, R.G., Linke, B.A. and Nash, D. M. 1977. Differential lipid oxidation in various parts of frozen mackerel. J. Food Technol. 12: 37-47.
- Kim, J., Marshall, M.R. and Wei, C.I. 2000. Polyphenoloxidase. In Seafood enzyme utilization and influence on postharvest seafood quality (Haard, N.F. and Simpson B.K., eds.) p. 271–315. Marcel Dekker. New York.
- King, I., Childs, M.T, Dorsett, C., Ostrander, J.G. and Monsey, E.R. 1990. Shellfish: proximate composition, minerals, fatty acids, and sterols. J. Am. Diet. Assoc. 90(5): 677-685.
- Kolakowska, A. 2002. Lipid oxidation in food systems. In Chemical and functional properties of food lipids (Sikorski, Z.E. and Kolakowska, A., eds.). p. 133–160. CRC Press. USA.
- Konosu, S. and Yamaguchi, K. 1982. The flavour components in fish and shellfish. In Chemistry and biochemistry of marine food products (Martin R.E., Flick G.J., Hebard C.E. and Ward, D.R., eds) p. 367-404. AVI Publishing. Westport.
- Koutsoumanis, K., Lampropoulou, K. and Nychas, G.E. 1999. Biogenic amines and sensory changes associated with the microbial flora of Mediterranean gilt-head sea bream (*Sparus aurata*) stored aerobically at 0, 8, and 15 °C. J. Food Prot. 62: 398-402.
- Kubota, K. and Kobayashi, A. 1988. Identification of unknown methyl ketones in volatile flavor components from cooked small shrimp. J. Agric. Food Chem. 36: 121-123.
- Laemmli, U.K. 1970. Cleavage of structural proteins during a assembly of head of bacteriophage T4. Nature. 227: 680-685.
- Ladrat, C., Verrez-Bagnis, V., Noel, J. and Fleurence, J. 2003. *In vitro* proteolysis of myofibrillar and sarcoplasmic proteins of white muscle of sea bass (*Dicentrarchus labrax* L.): effects of cathepsins B, D and L. Food Chem. 81: 517-525.

- Lakshmanan, P.T., Antony, P.D. and Gopakumar, K. 1996. Nucleotide degradation and quality changes in mullet (*Liza corsula*) and pearlspot (*Etroplus suratensis*) in ice and at ambient temperatures. Food Controls. 7: 227-283.
- Lakshmanan, R., Shakila, R.J. and Jeyasekaran, G. 2002. Survival of amine-forming bacteria during the ice storage of fish and shrimp. Food Microbiol. 19: 617-625.
- Lalitha, K.V. and Surendran, P.K. 2006. Microbiological changes in farm reared freshwater prawn (*Macrobrachium rosenbergii* de Man) in ice. Food Controls. 17: 802-807.
- Lall, S.P., 1989. The minerals. In Fish Nutrition, 2nd ed. (Halver, J.E., ed.). p. 231. Academic Press. San Diego, CA.
- Liao, I.C. and Chien, Y.H. 1994. Culture of kuruma prawn (*Penaeus japonicus*) in Asia. World Aquac. 25: 18-33.
- Liao, W.L., Nur-E-Borhan, S.A., Okada, S., Matsui, T. and Yamaguchi, K. 1993. Pigmentation of cultured black tiger prawn by feeding with a spirulina supplemented diet. Nippon Suisan Gakkaishi. 59: 165-169.
- Lin, R.Y., Huang L.S. and Huang, H.C. 2003. Characteristics of NADH-dependent lipid peroxidation in sarcoplasmic reticulum of white shrimp, *Litopenaeus vannamei*, and freshwater prawn, *Macrobrachium rosenbergii*. Comp. Biochem. Physiol. 135B: 683-687.
- Liston, J. 1980. Microbiology in fishery science. In Advances in Fish Science and Technology (Connell, J.J., ed.). p. 138-157. Fishing News Book Limited. Surrey, Farnham.
- Liston, J. 1982. Chemistry and biochemistry of marine food products. (Martin, R., Flick, G., Herbard. C.E. and Ward, D.E. eds). p. 27-36. AVI Publ. Co. Connecticut, USA.
- Lopez, A.C. and Maragoni, A.G. 2000. Lipases. In Seafood enzymes: utilization and influence on postharvest seafood quality (Harrd, N.F. and Simpson, B.K., eds.). p. 121-146. Marcel Dekker, Inc. New York.
- Lou, S.N. 1997. Effect of thermal processing on purine contents of grass shrimp (*Penaeus monodon*). J. Food Sci. 24: 438-447.
- Lou, S.N. 1998. Purine content in grass shrimp during storage as related to freshness. J. Food Sci. 63: 442-444.

- Lou, S.N., Chen, T.Y., and Chen, H.H. 1996. Determination of purine contents in some selected fishery products. *J. Nutr. Sci.* 21: 433-444.
- Lou, S.N., Chen, T.Y., Lin, C.D., and Chen, H.H. 1997. Effect of cooking on purine contents of some fish. *J. Food Sci.* 24: 258-262.
- Lovell, R.T. 1989. Nutrition and feeding of fish. Van nostrand reinhold, New York, 260 pp.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193: 256-275.
- Luzia, A.L., Sampaio, R.G., Castellucci, M.N.C. and Torres, A.F.S.E. 2003. The influence of season on the lipid profiles of five commercially important species of Brazilian fish. *Food Chem.* 83: 93-97.
- Mackie, I.M. 1994. Fish protein. In *New and developing sources of food proteins* (Hubson, B.J.G., ed.). p. 95-137. Chapman & Hall. New York.
- Manthey, M., Karnop, G and Rehbein, H. 1988. Quality changes of European catfish (*Silurus glanis*) from warm-water aquaculture during storage on ice. *Intern. J. Food Sci. Technol.* 23: 1-9.
- Martens, H., Stabursvik, E., and Martens, M. 1982. Texture and colour changes in meat during cooking related to thermal denaturation of muscle proteins. *J. Text. Stud.* 13: 291-309.
- Martinez-Alvarez, O., Lopez-Caballero, M.E., Montero, P. and Gomez-Guillen, M.C. 2007. Spraying of 4-hexylresorcinol based formulations to prevent enzymatic browning in Norway lobsters (*Nephrops norvegicus*) during chilled storage. *Food Chem.* 100: 147-155.
- Martinez, C.L., Campana, T.A. and Porchas, C.M. 2003. Dietary protein level and natural food management in the culture of blue (*Litopenaeus stylirostris*) and white shrimp (*Litopenaeus vannamei*) in microcosms. *Aquac. Nutr.* 9: 155-160.
- Martinez, I., Friis, T.J. and Careche, M. 2001. Post mortem muscle protein degradation during ice-storage of Arctic (*Pandalus borealis*) and tropical (*Penaeus japonicus* and *Penaeus monodon*) shrimps: a comparative eletrophoretic and immunological study. *J. Sci. Food Agric.* 81: 1199-1208.

- Matsumoto, J.J. 1980. Chemical deterioration of muscle proteins during frozen storage. In Chemical deterioration of protein (Whitaker, J.K. and Fujimaki, J.R. eds.) p. 95. ACS Symp. Series 123. Washington, DC.
- Matsumoto, M. and Yamanaka, H. 1991. Influences of antibiotics chloramphenicol on post-mortem biochemical changes in the muscle of kuruma prawn during storage. Bull. Jpn. Soc. Sci. Fish. 57: 2291-2297.
- McCormick, R.J. 1994. Structure and properties of tissue. In Muscle foods meat poultry and seafood technology (Kinsman, D.M., Kotula, A.W. and Breidenstein, B.C., ed.). p. 25-62. Chapman & Hall. New York.
- Mendes, R, Huidobro, A. and López-Caballero, E. 2002. Indole levels in deepwater pink shrimp (*Parapenaeus longirostris*). Eur. Food Res. Technol. 214: 125-30.
- Mendes, R., Quinta, R. and Nunes, M.L. 2001. Changes in baseline levels of nucleotides during ice storage of fish and crustaceans from the Portuguese coast. Eur. Food Res. Technol. 212: 141-146.
- Mensaveta, P., Worawattanamateekul, W., Latscha, Y. and Clark, J.S. 1993. Correction of tiger prawn (*Penaeus monodon* Fabricius) coloration by astaxanthin. Aquac. Eng. 12: 203-213.
- Merican, Z.O. and Shim, K.F. 1996. Qualitative requirements of essential fatty acids for juvenile *Penaeus monodon*. Aquaculture. 147, 275-291.
- Meyers, S.P. and Latscha, T. 1997. Carotenoids. In Crustacean nutrition, advances in world Aquaculture, vol. 6. (D'Abromo, L.R., Conklin, D.E. and Akiyama, D.M., eds.). p. 164-193. World Aquaculture Society. Baton Rouge, LA.
- Middlebrooks, B. L., Toom, P. M., Douglas, W. L., Harrison, R. E and McDowell, S. 1988. Effects of storage time and temperature on the microflora and amine development in Spanish mackerel (*Scomberomorus maculatus*). J. Food Sci. 53: 1024-1029.
- Mietz, J.L. and Karmas, E. 1977. Chemical quality index of canned tuna as determined by high pressure liquid chromatography. J. Food Sci. 42: 155-158.
- Mietz, J.L. and Karmas, E. 1978. Polyamines and histamine content in rockfish, salmon, lobster and shrimp s an indicator of decomposition. J. Assoc. Off. Anal. Chem. 61: 139-145.

- Mizuta, S., Yamada, Y., Miyagi, T., and Yoshinaka, R. 1999. Histological changes in collagen related to textural development of prawn meat during heat processing. *J. Food Sci.* 64: 991-995.
- Mizuta, S., Yoshinaka, R., Sato, M., and Sakaguchi, M. 1994. Cahracterization of collagen in muscle of several crustacean species in association with raw meat texture. *Fish. Sci.* 60: 323-328.
- Mizuta, S., Yoshinaka, R., Sato, M., and Sakaguchi, M. 1997. Histological and biochemical changes of muscle collagen during chilled storage of the kuruma prawn *Penaeus japonicus*. *Fish. Sci.* 63: 784-793.
- Mohan, C.O., Ravishankar, C.N., Bindu, J. Geethalakshmi, V. and Srinivasn Gopal, T.K. 2006. Effect of themal process time on quality of “shrimp kuruma” in retortable pouches and aluminum cans. *J. Food Sci.* In press.
- Montero, P. and Borderias, J. 1990. Effect of rigor mortis and ageing on collagen in trout (*Salmo irideus*) muscle. *J. Sci. Food Agric.* 52: 141-146.
- Montero, P., Avalosb, A. and Perez-Mateosa, M. 2001. Characterization of polyphenoloxidase of prawns (*Penaeus japonicus*). Alternatives to inhibition: additives and high-pressure treatment. *Food Chem.* 75: 317–324.
- Moori, H., Cann, L. Y. and Taylor, L. Y. 1988. Histamine formation by luminous bacteria in mackerel stored at low temperatures. *Nippon Suisan Gakkaishi.* 54: 299-305.
- Morita, K., Kubota, K. and Aishima, T. 2001. Sensory characteristics and volatile components in aromas of boiled prawns prepared according to experimental designs. *Food Res. Intern.* 34: 473-481.
- Murakami, E.G. 1994. Thermal processing affects properties of commercial shrimp and scallops. *J. Food Sci.* 59: 237-241.
- Nakai, S. and Li-Chan, E. 1988. Hydrophobic interaction in muscle protein and their role in comminuted meat systems. *In Hydrophobic interaction in food systems.* (Nakai, S. and Li-Chan, E., eds.). p 63-127. Boca Raton, FL CRS press.
- Namiki, M. 1990. Antioxidant/antimutagens in food. *Crit. Rev. Food Sci. Nutr.* 29: 273-300.
- Nawar, W.W. 1996. Lipids. *In Food chemistry* (Fennema, O.R., ed.). p. 225-314. Marcel Dekker, Inc. New York.

- Nilsson, L.B. and Ekstrand, B. 1995. Frozen storage and thawing methods affect biochemical and sensory attributes of rainbow trout. *J. Food Sci.* 63: 627-635.
- Nip, W.K. and Moy, J.H. 1988. Microstructural changes of ice-chilled and cooked freshwater prawn, *Macrobrachium rosenbergii*. *J. Food Sci.* 53: 319-322.
- Nip, W.K., Lan, C.Y. and Moy, J.H. 1985. Partial characterization of a collagenolytic enzyme fraction from the hepatopancreas of the freshwater prawn, *Macrobracnium rosenbergii*. *J. Food Sci.* 50: 1187.
- Nishimoto, J., Suwetja, I.K. and Miki, H. 1985. Estimation of keeping freshness period and practical storage life of mackerel muscle during storage at low temperatures. *Mem. Fac. Fish. Kagoshima Univ.* 34: 89-96.
- Ochiai, Y. and Chow, C.J. 2000. Myosin ATPase. In Seafood enzymes. (Haard, N.F. and Simpson, B.K., eds.) p. 69-90. Marcel Dekker. New York.
- Oehlenschlager, J. 1997. Volatile amines as freshness/spoilage indicators, a literature review: 571-578.
- Offer, G. and Knight, P. 1988. The structural basis of water-holding in meat. In Developments in meat science-4 (Lawrie, R., ed.). p. 63–243. Elsevier Applied Science. London.
- Ogawa, M., Ehara, T., Tamiya, T. and Tsuchiya, T. 1993. Thermal stability of fish myosin. *Comp. Biochem. Physiol.* 106: 517-523.
- Ogawa, M., Magalhaes-Neto, E. de O., Aguiar-Junior, O. and Kozima, T.T. 1984. Incidence of melanosis in the integumentary tissue. *Bull. Jpn. Soc. Sci. Fish.* 50: 471-475.
- Ogawa, M., Tamiya, T. and Tsuchiya, T. 1994. Structural changes of carp myosin during heating. *Fish. Sci.* 60: 723-727.
- Okada, S., Nur-E-Borhan, S.A. and Yamaguchi, K. 1994. Carotenoid composition in the commercial black tiger prawns. *Fish. Sci.* 60: 213-215.
- Okada, S., Nur-E-Borhan, S.A., Watabe, S. and Yamaguchi, K. 1995. Changes in body color appearance of the black tiger prawn *Penaeus monodon* by the varied composition of carotenoids soluble as carotenoprotein and remaining insoluble after collagenase treatment for the muscular epithelium. *Fish. Sci.* 61: 964-967.

- Okada, T., Inoue, N. and Akiba, M. 1986. Electron microscopic observation and biochemical properties of carp myosin B during frozen storage. Bull. Japan. Soc. Sci. Fish. 52: 345-353.
- Ozogul, F., Gokbulut, C., Ozogul, Y. and Ozyurt, G. 2006. Biogenic amine production and nucleotide ratios in gutted wild sea bass (*Dicentrarchus labrax*) stored in ice, wrapped in aluminium foil and wrapped in cling film at 4 °C. Food Chem. 98: 76-84.
- Ozogul, F., Gokbulut, C., Ozyurt, G., Ozogul, Y. and Dural, M. 2005. Quality assessment of gutted wild sea bass (*Dicentrarchus labrax*) stored in ice, cling film and aluminium foil. Eur. Food Res. Technol. 220: 292-298.
- Pacheco-Aguilar, R., Lugo-Sanchez, M.E. and Robles-Burgueno, M.R. 2000. Postmortem biochemical characteristic of Monterey sardine muscle stored at 0°C. J. Food. Sci. 65: 40-47.
- Palka, K. and Daun, K. 1999. Changes in texture, cooking losses, and myofibirellar structure of bovine *M. semitendinosus* during heating. Meat Sci.. 51: 237-243.
- Pearsons, A.M. and Young, R.B. 1989. Muscle and meat biochemistry. Academic Press, Inc. San Diego. CA.
- Peleologos, E.K., Savvaidis, I.N. and Kontominas, M.G. 2004. Biogenic amines formation and its relation to microbiological and sensory attributes in ice-stored whole, gutted and filleted Mediterranean sea bass (*Dicentrarchus labrax*). Food Microbiol. 21: 549-557.
- Penafiorida, V.D. 1999. Interaction between dietary levels of calcium and phosphorus on growth of juvenile shrimp, *Penaeus monodon*. Aquac. 172: 281- 289.
- Poulter, R.G., Ledward, D.A., Godber, S., Hall, G. and Rowlands, B. 1985. Heat stability of fish muscle protein. J. Food Technol. 20: 203-217.
- Ragnarsson, K. and Regenstein, J.M. 1989. Changes in electrophoretic patterns of gadoid and non-gadoid fish muscle during frozen storage. J. Food Sci. 54: 819-823.
- Rahaman, M., Yasmin, L., Kamal, M.D., Mazid, M.A. and Islam, N. M.D. 2001. Effect of delayed icing on the quality changes in brackish water shrimp *Penaeus monodon* during ice storage. J. Biol. Sci. 11: 1390-1394.

- Rawles, D.D. and Flick, G.J. 1996. Biogenic amines in fish and shellfish. *Adv. Food Nutr. Res.* 39: 329-365.
- Rehbein, H. 1979. Development of an enzymatic method to differentiate fresh and sea-frozen and thawed fish fillets I. Comparison of the applicability of some enzymes of fish muscle. *Z. Lebensmittel-Untersuchung und-Forsc.* 169: 263–265.
- Rehbein, H. 1988. Relevance of trimethylamine oxide dimethylase activity and haemoglobin content to formaldehyde production and texture deterioration in frozen stored minced fish muscle. *J. Sci. Food Agric.* 43: 261.
- Riaz, F. and Qadri, R. B. 1985. Quality changes in lobster (*Panulirus poliphagus*) muscle during storage in ice. *J. Agric. Food. Chem.* 33: 117-122.
- Robinson, H.W. and Hodgen, C.G. 1940. The biuret reaction in the determination of serum protein I. A study of the condition necessary for the production of the stable color which bears a quantitative relationship to the protein concentration. *J. Biol. Chem.* 135: 707–725.
- Rolle, R.S., Guizani, N., Chen, J.S., Marshall, M.R, Yan, J.S. and Wei, C.I. 1991. Purification and characterization of phenoloxidase isoforms from Taiwanese black tiger shrimp (*Penaeus monodon*). *J. Food Biochem.* 15: 17-32.
- Rosa, R. and Nunes, L.M. 2003. Nutritional quality of red shrimp, *Aristeus antennatus* (Risso), pink shrimp, *Parapenaeus longirostris* (Lucas), and Norway lobster, *Nephrops norvegicus* (Linnaeus). *J. Sci. Food Agric.* 84: 89 – 94.
- Rowland, B., Finne, G. and Tillman, R. 1982. A morphological study of muscle proteolysis in the tails of *Macrobrachium rosenbergii*. *Proc. 7th Annual Trop. Subtrop. Fish. Technol. Conf. Amer. TAMU-SG-82-10.* p. 105.
- Saito, T., Arai, K. and Matsuyoshi, M. 1959. A new method for estimating the freshness of fish. *Bull. Jpn. Soc. Sci. Fish.* 24: 739-740.
- Salvato, B., Santamaria, M., Beltramini, M., Alzuet, G. and Casella, L. 1998. The enzymatic properties of *Octopus vulgaris* hemocyanin: o-diphenol oxidase. *Biochem.* 37: 14065-14077.
- Sasi, M., Jeyasekaran, G., Shanmugam, S.A., and Jeya Shakila, R. 2000. Chilling fresh fish in dry and wet ice. *Asian Fish. Sci.* 13: 375-382.

- Sato, K., Ohashi, C., Ohtsuki, K. and Kawabata, M. 1991. Type V collagen in trout (*Salmo gairdneri*) muscle and its solubility change during chilled storage of muscle. J. Agric. Food Chem. 39: 1222-1225.
- Sato, K., Yoshinaka, R., Sato, M. and Shimizu, Y. 1986. Collagen content in the muscle of fishes in association with their swimming movement and meat texture. Bull. Jpn. Soc. Sci. Fish. 52: 1595-1598.
- Sato, K., Yoshinaka, R., Sato, M., Itoh, H. and Shimizu, Y. 1988. Isolation of type I and V collagens from carp muscle. Comp. Biochem. Physiol. 90B: 155-158.
- Schiedt, K., Bischof, S. and Glinz, E. 1993. Metabolism of carotenoids and *in vivo* racemization of (3S, 3'S)-astaxanthin in the crustacean *Penaeus*. Meth. Enzymol. 214: 148-168.
- Senoz, B., Isykly, N. and Coksoyler, N. 2000. Biogenic amines in Turkish sausages (sucuk). J. Food Sci. 65: 764-767.
- Seo, H., Endo, Y., Fujimoto, K., Moku, M. and Kawaguchi, K. 1997. Denaturation of myofibrillar protein in myctophid fish during refrigeration and freezing storage. Fish. Sci. 63: 839-840.
- Shahidi, F. 1994. Seafood proteins and preparation of protein concentrates. In Seafoods chemistry, processing technology and quality (Shahidi, F. and Botta, J.R., eds.). p. 3-9. Chapman & Hall. Great Britain.
- Shakila, R.J., Vasundhara, T.S. and Rao, D.V. 1995. Rapid quality assessment of shrimps during storage by monitoring amines. J. Food Sci. Technol. 32: 310-314.
- Shamshad, S.I., Kher-Un-Nisa, Riaz, M., Zuberi, R. and Qadri, R.B. 1900. Shelf-life of shrimp (*Penaeus merguiensis*) stored at different temperatures. J. Food Sci. 55: 1201.
- Shenouda, S.Y.K. 1980. Theories of protein denaturation during frozen storage of fish flesh. Adv. Food Res. 26: 275-311.
- Shimomura, H., Takahashi, T., Morishita, T. and Ueno, R. 1987. Investigation of the differentiation of frozen-thawed fish from unfrozen fish by comparison of lysosomal enzyme activity. Nippon Suisan Gakkaishi. 53: 1841-1845.
- Sikorski, Z.E., 1994. The myofibrillar proteins in seafoods. In Seafood proteins (Sikorski, Z.E., Pan, B.S. and Shahidi, F., eds.). p. 40-57. Chapman & Hall. New York.

- Sikorski, Z.E. Kolokowska, A. and Burt, Jr. 1990. Postharvest biochemical and microbial Changes. In Seafood: resource, nutritional composition and preservation. (Sikorski, A.E., ed.). p. 55-75. CRC Press, Inc. Florida.
- Sikorski, Z.E. Kolokowska, A. and Sun Pan, B. 1990. The nutritive composition of the major groups of marine food organisms. In Seafood: resources, nutritional composition and preservation (Sikorski, Z.E., ed.). p. 29-54. CRC press, Inc. Florida.
- Sikorski, Z.E, Olley, J. and Kostuch, S. 1976. Protein changes in frozen fish. CRC. Crit. Rev. Food Sci. Nutr. 8: 97-129.
- Silla-Santos, S.M.H. (1996). Biogenic amines: their importance in foods. Intern. J. Food Microbiol. 29: 213-231.
- Sims, G.G., Farn, G. and York, R.K. 1992. Quality index for tuna: correlation of sensory attributes with chemical indices. J. Food Sci. 57: 1112-1115.
- Sivakumar, P., Suguna, L. and Chandrasekaran, G. 1997. Purification and partial characterization of a type V like collagen from the muscle of marine prawn, *Penaeus indicus*. J. Biosci. 22: 131-141.
- Sivakumar, P., Suguna, L. and Chandrasekaran, G. 2000. Molecular species of collagen in the intramuscular connective tissues of the marine crab, *Scylla serrata*. Comp. Biochem. Physiol. 125B: 555-562.
- Söderhäll, K. and Cerenius, L. 1992. Crustacean immunity. Annu. Rev. Fish Diseases. 2: 3-23.
- Söderhäll, K. and Smith, V.J. 1986. The prophenoloxidase activating system: the biochemistry of its activation and role in cellular immunity, with special reference to crustaceans. In Immunity in invertebrates: cells, molecules, and defense reactions (Brehelin M, ed.). p. 208-23. Springer-Verlag. Berlin.
- Srinivasan, S. and Hultin, H.O. 1997. Chemical, physical and functional properties of cod proteins modified by a nonenzyme free-radical-generating system. J. Agric. Food Chem. 45: 310-320.
- Srinivasan, S., Xiong, Y.L. and Blanchard, S.P. 1997. Effects of freezing and thawing methods and storage time on thermal properties of freshwater prawns (*Machrobrachium rosenbergii*). J. Sci. Food Agric. 75: 37-44.

- Srinivasan, S., Xiong, Y.L., Blanchard, S.P. and Tidwell, J.H. 1997. Physicochemical changes in prawns (*Machrobrachium rosenbergii*) subjected to multiple freeze-thaw cycles. *J. Food Sci.* 62: 123-127.
- St. Angelo, A.J. 1996. Lipid oxidation in foods. *CRC. Crit. Rev. Food Sci. Nutr.* 36: 175-224.
- Steel, R.G.D. and Torrie, J.H. 1980. Principle and procedures of statistics: a biometrical approach. McGraw-Hill. New York.
- Stepnowski, P., O' lafsson, G., Helgason, H. and Jastor, B. 2004. Recovery of astaxanthin from seafood wastewater utilizing fish scales waste. *Chemosphere.* 54: 413– 417.
- Strachan, N.J.C. and Nicholson, F. 1992. Gill air a analysis as an indicator of cod freshness and spoilage. *Intern. J. Food Sci. Technol.* 27: 261-269.
- Suphamongkhon, K. 2002. Office of the Thai trade representative. *Thai Trade Representative Thailand's Policy on International Trade and Investment after the 1997 Economic Crisis for the International Boston Seafood Show, Massachusetts Boston,* 12 March.
- Takama, K., Suzuki, T., Yoshida, K., Hirofumi, A. and Mitsui, T. 1999. Phosphatidyl choline levels and their fatty acid composition in teleost tissues and squid muscle. *Comp. Biochem. Physiol.* 124B: 109-116.
- Taylor, S.L. 1986. Histamine food poisoning: toxicology and clinical aspects. *Crit. Rev. Toxicol.* 17: 91-128.
- Thannhauser, T.W., Konishi, Y. and Scheraga, H.A. 1987. Analysis for disulfide bonds in peptides and proteins. *Meth. Enzymol.* 143: **115-118**.
- Thanonkaew, A., Benjakul, S. and Visessanguan, W. 2006. Chemical composition and thermal property of cuttlefish (*Sepia pharaonis*) muscle. *J. Food Comp. Anal.* 19: 127-133.
- Thanonkaew, A., Benjakul, S. and Visessanguan, W. 2006. The effect of metal ions on lipid oxidation, colour and physicochemical properties of cuttlefish (*Sepia pharaonis*) subjected to multiple freeze-thaw cycles. *Food Chem.* 95: 591–599.
- Tichivangana, J.Z. and Morrissey, P.A. 1985. Metmyoglobin and inorganic metals as prooxidants in raw and cooked muscle systems. *Meat Sci.* 15: 107-116.
- Tsai, I.H., Chuang, K.L. and Chuang, J.L. 1986. Chymotrypsins in digestive tracts of crustacean decapods (shrimps). *Comp. Biochem. Physiol.* 85B: 235-239.

- U.S. Food and Drug Administration. 2005. Frozen shrimp sensory evaluation short course. National Center for Genetic Engineering and Biotechnology (BIOTEC).
- Van Holde, K.E. and Miller, K.I. 1995. Hemocyanins. In Advances in protein chemistry. (Anfinsen, C.B., Edsall, J.T., Eisenberg, D.S. and Richards F.M., eds.). p. 1-81. Academic. Press, Inc. New York.
- Veiseth, E., Fjaera , S.O., Bjerkeng, B. and Skjervold, P.O. 2006. Accelerated recovery of Atlantic salmon (*Salmo salar*) from effects of crowding by swimming. Comp. Biochem. Physiol. 144 B: 351–358.
- Vogt, W. 1995. Oxidation of methionyl residues in protein: tool, target, and reversal. Free Rad. Biol. Med. 18: 93-105.
- Vojdani, F. 1996. Solubility. In Method of testing protein functionality (Hall, G.M., ed.). p. 11-60. Chapman & Hall. London.
- Wangner, J.R. and Anon, M.C. 1986. Effect of frozen storage on protein denaturation in bovine muscle. II. Influence on solubility, viscosity and electrophoretic behavior of myofibrillar proteins. J. Food Technol. 2: 547-553.
- Wright, D.J. and Wilding, P. 1984. Differential scanning calorimetric study of muscle and its proteins: myosin and subfragments. J. Sci. Food Agric. 35: 357-372.
- Xiong, Y.L. 1997. Structure-function relationships of muscle protein. In Food protein and their applications. (Damodaran, S. and Paraf, A., eds.). p. 341-392. Marcel Dekker. New York.
- Xiong, Y.L. 2000. Protein oxidation and implication for muscle food quality. In Antioxidants in muscle food (Decker, E.A. Faustman, C. and Lopez-Bote, C.J., eds.). p. 85-111. John: Wiley & Sons, Inc. New York.
- Xiong, Y.L. and Decker, E.A. 1995. Alteration of muscle protein functionality by oxidative and antioxidative process. J. Muscle Food. 6: 139-160.
- Yamagata, M. and Low, L.K. 1995. Banana shrimp, *Peneaus merguiensis*, quality changes during iced and frozen storage. J. Food Sci. 60: 721-726.

- Yanar, Y. and Celik, M. 2005. Seasonal variations of fatty acid composition in wild marine shrimps (*Penaeus semisulcatus* De Haan, 1844 and *Metapenaeus monoceros* Fabricius, 1789) from the Eastern Mediterranean Sea. Food Sci. Technol. Intern. 11: 391-395.
- Yanar, Y. and Celik, M. 2006. Seasonal amino acid profiles and mineral contents of green tiger shrimp (*Penaeus semisulcatus* De Haan, 1844) and speckled shrimp (*Metapenaeus monoceros* Fabricius, 1789) from the Eastern Mediterranean. Food Chem. 94: 33-36.
- Yanar, Y., Celik, M. and Yanar, M. 2004. Seasonal changes in total carotenoid contents of wild marine shrimps (*Penaeus semisulcatus* and *Metapenaeus monoceros*) inhabitating the Eastern Mediterranean. Food Chem. 88: 267-269.
- Yepiz-Plascencia, G., Vargas-Albores, F. and Higuera-Ciapara, I. 2000. Penaed shrimp hemolymph lipoproteins. Aquaculture. 191: 177-189.
- Yokoyama, Y., Sakaguchi, M., Kawai, F. and Kanamori, M. 1992. Changes of the solubility and ATP-related compounds in various tissues of oyster during ice storage. Bull. Jpn. Soc. Sci. Fish. 58: 2125-2136.
- Yongswawatdigul, J. and Park, J.W. 2002. Biochemical and conformation changes of actomyosin from threadfin bream stored in ice. J. Food Sci. 67: 895-990.
- Yoshinaka, R., Mizuta, S., Itoh, Y. and Sato, M. 1990. Two genetically distinct types of collagen in kuruma prawn, *Penaeus japonicus*. Comp. Biochem. Physiol. 96B: 451-456.
- Yoshinaka, R., Sato, K., Itoh, Y., Nakajima, S. and Sato, M. 1989. Content and partial characterization of collagen in crustacean muscle. Comp. Biochem. Physiol. 94B: 219-233.
- Young, O.A. and West, J. 2001. Meat color. In Meat science and applications (Hui, Y.H., Nip, Wai-Kit Rogers, R.W. and Young, O.A., eds.). p. 39-69. Marcel Dekker, Inc. New York.
- Zayas, J.F. 1997. The Effect of freezing and storage; whey frozen on protein solubility. In Functionality of proteins in food. p. 18-21. Springer. New York.
- Zeng, Z.Q., Thorarinsdottir, A.K. and Olafsdottir, G. 2005. Quality changes of shrimp (*Pandalus borealis*) stored under different cooling conditions. J. Food Sci. 70: 459-466.