

## REFERENCES

- Ababouch, L.H., Souibri, L., Rhaliby, K., Ouahdi, O., Battal, M. and Busta, F.F. 1996. Quality changes in sardines (*Sardina pilchardus*) stored in ice and at ambient temperature. *Food Microbiol.* 13: 123-132.
- Adams, M.R. and Marteau, P. 1995. On the safety of the lactic acid bacteria. *Int. J. Food Microbiol.* 27: 263-264.
- Adams, M.R. and Moss, M.O. 1995. *Food Microbiology*. pp. 192-203. The Royal Society of Chemistry, Cambridge.
- Adams, M.R. and Moss, M.O. 2000. *Food Microbiology*. pp. 156-251. The Royal Society of Chemistry, Cambridge.
- Aickin, C.C. and Thomas, R.C. 1975. Micro-electrode measurement of the internal pH of carp muscle fiber. *J. Physiol.* 52: 803-807.
- Aktas, N. and Kaya, M. 2001. The influence of marinating with weak organic acids and salts on the intramuscular connective tissue and sensory properties of beef. *Eur. Food Res. Technol.* 231: 88-94.
- Ali, M.T., Gleeson, R.A., Wei, C.I. and Marshall, M.R. 1994. Activation mechanisms of prophenoloxidase on melanosis development in Florida spiny lobster (*Pandalirus argus*) cuticle. *J. Food Sci.* 59: 1024-1030.
- Alvarez, J.A., Pozo, R. and Pastoriza, L. 1996. Effect of a cryoprotectant agent (sodium tripolyphosphate) on hake slices preserved in modified atmosphere packing. *Food Sci. Technol. Int.* 2: 177-181.
- Amano, K. and Yamada, K. 1964. A biological formation of FA in the muscle tissue of gadoid fish. *Nippon Suisan Gakkaishi.* 30: 430-435.
- An, H., Weewasinghe, V., Seymour, T.A. and Morrissey, M.T. 1994. Cathepsin degradation of Pacific whiting surimi proteins. *J. Food Sci.* 59: 1013-1017, 1033.
- An, H., Seymour, T.A., Wu, J. and Morrissey, M.T. 1994. Assay systems and characterization of Pacific whiting (*Merluccius productus*) protease. *J. Food Sci.* 59: 277-281.
- Ando, S., Osada, K., Hatano, M. and Saneyoshi, M. 1992. Metabolism of astaxanthin in muscle and ovary from brook trout *Salvelinus fontinalis*. *Comp. Biochem. Physiol.* 96B: 355-359.

- Anon. 1992. Code of Federal Regulations, Food and Drugs 21 Parts 170.199. Office of the Federal Register National Archives and Records Administration: Washington, DC.
- Anonymous. 1998. Aquaculture drugs. *In* Fish and Fishery Products Hazards and Control Guide, FDA, Center for Food Safety and Applied Nutrition, Washington, DC: Office of Seafood. pp. 115-132.
- Anonymous. 2000. Proposed draft code of practice for fish and fishery products. Joint FAO/WHO food standards programme. Codex committee on fish and fishery products. 24<sup>th</sup> Session. Alesund: Norway. pp. 108. 5-9.
- Aoki, H., Ahsan, M.N., Matsuo, K., Hagiwara, T. and Watabe, S. 2003. Purification and characterization of collagenolytic proteases from the hepatopancreas of Northern shrimp (*Pandalus eous*). *J. Agric. Food Chem.* 51: 777-783.
- Aoki, H., Ahsan, M.N., Matsuo, K., Hagiwara, T. and Watabe, S. 2004. Partial purification of proteases that are generated by processing of the Northern shrimp (*Pandalus borealis*) and which can tenderize beef. *Int. J. Food Sci.* 39: 471-480.
- Applewhite, L., Otwell, W.S. and Marshall, M. 1991. Effect of kojic acid on pink shrimp phenoloxidase. Report of the Florida Sea Grant College Program. pp. 141-146.
- Applewhite, L., Otwell, W.S. and Marshall, M.R. 1990. Effect of kojic acid on pink shrimp phenoloxidase. Proceedings of the Fifteenth Annual Conference Tropical and Subtropical Fisheries Technological Conference of the Americas. pp. 41-146. Orlando, FL.
- Aranishi, F., Hara, K., Osatomi, K. and Ishihara, T. 1997. Substrate specificity of carp (*Cyprinus carpio*) cathepsin H with methylcoumarylamide substrates. *Comp. Biochem. Physiol.* 116B: 203-208.
- Asghar, A. and Bhatti, A.R. 1987. Endogenous proteolytic enzymes in skeletal muscle: their significance in muscle physiology and during postmortem aging events in carcasses. *Adv. Food Res.* 343-451.
- Ashie, I.N.A., Simpson, B.K. and Ramaswamy, H.S. 1997. Changes in texture and microstructure of pressure-treated fish muscle tissue during chilled storage. *J. Muscle Foods.* 8: 13-32.

- Ashie, I.N.A., Smith, J.P. and Simpson, B.K. 1996. Spoilage and shelf-life extension of fresh fish and shellfish. *Crit. Rev. Food Sci. Technol.* 36: 87-121.
- Bak, L.S., Andersen, A.B., Andersen, E.M. and Bertelsen, G. 1999. Effect of modified atmosphere packaging on oxidative changes in frozen stored cold water shrimp (*Pandalus borealis*). *Food Chem.* 64: 169-175.
- Balamatsia, C.C., Paleologos, E.K., Kontominas, M.G. and Savvaidis, I.N. 2006. Correlation between microbial flora, sensory changes and biogenic amines formation in fresh chicken meat stored aerobically or under modified atmosphere packaging at 4°C: possible role of biogenic amines as spoilage indicators. *Antonie van Leeuwenhoek.* 89(1): 9-17.
- Banks, H., Nickelson, H.R. and Finne, G. 1980. Shelf-life studies on carbon dioxide packaged finfish from the gulf of Mexico. *J. Food Sci.* 45: 157-162.
- Baranoski, E.S., Nip, W.K. and Moy, J.H. 1984. Partial characterization of a crude enzyme extract from the freshwater prawn, *Macrobrachium rosenbergii*. *J. Food Sci.* 49: 1494.
- Barnett, H.J., Nelson, R.W., Hunter, P.J. and Groninger, H. 1978. Use of carbon dioxide dissolved in refrigerated brine for the preservation of pink shrimp (*Pandalus* spp.) *Marine Fish. Rev.* 40: 24-28.
- Barrett, A.J. 1997. Cathepsin B and other thiol proteinases. *In* Proteinase in Mammalian Cells and Tissues. (Barrett, A.J. ed.), pp 181-208. North-Holland: Amsterdam.
- Barrett, A.J. and Kirschke, H. 1981. Cathepsin B, cathepsin H and cathepsin L. *Methods Enzymol.* 80: 535-561.
- Bartolo, I. and Birk, E.O. 1998. Some factors affecting Norway lobster (*Nephrops norvegicus*) cuticle polyphenol oxidase activity and blackspot development. *Int. J. Food Sci. Technol.* 33: 329-336.
- Bendall, J.R. 1954. The swelling effect of polyphosphates on lean meat. *J. Sci. Food Agric.* 5: 468-475.
- Benjakul, S. and Morrissey, M.T. and An, H. 1997. Physicochemical changes in Pacific whiting muscle proteins during iced storage. *J. Food Sci.* 62: 729-733.

- Benjakul, S., Visessanguan, W., Riebroy, S., Ishizaki, S. and Tanaka, M. 2002. Gel-forming properties of surimi produced from bigeye snapper, *Priacanthus tayenus* and *P. macracanthus*, stored in ice. *J. Sci. Food Agric.* 82: 1442-1451.
- Benjakul, S., Visessanguan, W. and Leelapongwattana, K. 2003a. Purification and characterization of heat-stable alkaline proteinase from bigeye snapper (*Priacanthus macracanthus*) muscle. *Comp. Biochem. Physiol.* 134B: 579-591.
- Benjakul, S., Visessanguan, W. and Tueksuban, J. 2003b. Changes in physico-chemical properties and gel-forming ability of lizardfish (*Saurida tumbil*) during post-mortem storage in ice *Food Chem.* 80: 535-544.
- Benjakul, S., Visessanguan, W. and Tueksuban, J. 2003c. Heat-activated proteolysis in lizardfish (*Saurida tumbil*) muscle. *Food Res. Int.* 36: 1021-1028.
- Benjakul, S., Visessanguan, W. and Tanaka., M. 2005. Properties of phenoloxidase isolated from the cephalothorax of kuruma prawn (*Penaeus japonicus*). *J. Food Biochem.* 29: 470-485.
- Benner, R.A., Miget, R., Finne, G. and Acuff, G.R. 1994. Lactic acid/melanosis inhibitors to improve shelf life of brown shrimp (*Penaeus aztecus*). *J. Food Sci.* 59: 242-250.
- Bennett, R.W., Yeterian, M., Smith, W., Coles, C.M., Sassaman, M. and McClure, F.D. 1986. *Staphylococcus aureus* identification characteristics and enterotoxigenicity. *J. Food Sci.* 51:1337-1339.
- Boone, B.B. 1982. Modified atmosphere packaging of fresh fish. M.S. Thesis, University of Washington, Seattle, W.A.
- Borch, E., Muermans, M.L.K. and Blixt, Y. 1996. Bacterial spoilage of meat and cured meat products. *Int. J. Food Microbiol.* 33: 103-120.
- Boskou, G. and Debevere, J. 2000. Shelf-life extension of cod fillets with acetate buffer spray prior to packaging under modified atmospheres. *Food Add. Contam.* 17: 17-25.
- Bradshaw, J.G., Francis, D.W. and Twedt, R.M. 1974. Survival of *Vibrio parahaemolyticus* in cooked seafood at refrigeration temperatures. *Appl. Microbiol.* 27: 657-661.
- Branen, A.L. 1978. Interaction of fat oxidation and microbial spoilage in muscle foods. Proceeding of the 31st reciprocal meat conference. Chicago, IL: National Live Stock and Meat Board.

- Brashears, M.M. and Durre, W.A. 1999. Antagonistic action of *Lactobacillus lactis* toward *Salmonella spp.* and *Escherichia coli* O157: H7 during growth and refrigerated storage. *J. Food Prot.* 62: 1336-1340.
- Brauer, J.M.E., Leyva, J.A.S., Alvarado, L.B. and Sandez, O.R. 2003. Effect of dietary protein on muscle collagen, collagenase and shear force of farmed white shrimp (*Litopenaeus vannamei*). *Eur. Food Res. Technol.* 217: 277-280.
- Bremner, A. 1998. If freshness is lost, where does it go? *In* Methods to determine the freshness of fish in research and industry. (Ólafsdóttir, G., Luten, J., Dalgaard, P., Careche, M., Verrez, Bagnis. V., Martinsdóttir. E. and Heia, K. eds.), pp 36-51. *Int. Inst. of Refrigeration: Paris.*
- Britton, G., Armit, G.M. Lau, S.Y.M. Patel, A.K. and Shone, C.C. 1981. Carotenoproteins *In* Carotenoid Chemistry and Biochemistry (Britton, G. and Goodwin, T.W. eds), pp. 237-251. Pergamon Press: Oxford
- Browdy, C.L., 1998. Recent developments in penaeid broodstock and seed production technologies: improving the outlook for superior captive stocks. *Aquaculture.* 164: 3-21.
- Buchanan, R.L. and Klawitter, L.A. 1991. Effect of temperature history on the growth of *Listeria monocytogenes* Scott A at refrigeration temperatures. *Int. J. Food Microbiol.* 12: 235-245.
- Buege, J.A. and Aust, S.D. 1978. Microsomal lipid peroxidation. *Method Enzymol.* 52: 302-304.
- Bullard, F.A. and Collins, J. 1978. Physical and chemical changes of pink shrimp, *Pandalus borealis*, held in carbon dioxide modified refrigerated seawater compared with pink shrimp held in ice. *Fish. Bull.* 76: 73-78.
- Buta, J.G. and Moline, H.E. 2001. Prevention of browning of potato slices using polyphenoloxidase inhibitors and organic acids. *J. Food Quali.* 24: 271-282.
- Cao, M.J., Osatomi, K., Suzuki, M., Hara, K., Tachibana, K. and Ishihara, T. 2000. Purification and characterization of two anionic trypsin from the hepatopancreas of crap. *Fish. Sci.* 66: 1172-1179.
- Cann, D.C. 1988. Modified atmosphere packaging of fishery products. *INFOFISH Int.* 1: 37-39.

- Cann, D.C., Smith, G.L. and Huston, N.C. 1983. Further studies on marine fish stored under modified atmosphere packaging. Torry Research Station: Aberdeen. 61p.
- Careche, M., Mazo, M.L., Torrejon, P. and Tejada, M. 1998. Importance of frozen storage temperature in the type of aggregation of myofibrillar proteins in cod (*Gadus morhua*) fillets. *J. Agric. Food Chem.* 46: 1539-1546.
- Chaitiamvong, S. and Supongpan, M. 1992. A guide to penaeoid shrimps found in thai waters. Australian Institute of Marine Science. Townsville: Australia.
- Chen, J.S., Balaban, M., Wei, C., Gleeson, R.A. and Marshall, M.R. 1993. Effect of carbon dioxide on the inactivation of Florida spiny lobster polyphenol oxidase. *J. Sci. Food Agric.* 61: 253-259.
- Chen, J.S., Balaban, M., Wei, C., Marshall, M.R. and Hsu, W.Y. 1992. Inactivation of polyphenol oxidase by high pressure carbon dioxide. *J. Agric. Food Chem.* 40: 2345-2349.
- Chen, J.S., Charest, D.J., Marshall, M.R. and Wei, C.I. 1997. Comparison of two treatment methods on the purification of shrimp polyphenol oxidase. *J. Sci. Food Agric.* 75: 12-18.
- Chen, J.S., Rolle, R.S., Marshall, M.R. and Wei, C.I. 1991a. Comparison of phenoloxidase activity from Florida spiny lobster and Western Australian lobster. *J. Food Sci.* 56: 154-157.
- Chen, J.S., Wei, C.I. and Marshall, M.R. 1991b. Inhibitory mechanism of kojic acid on polyphenol oxidase. *J. Agric. Food Chem.* 39: 1897-1901.
- Chen, T.C. and Waimaleongora-Ek, C. 1981. Effect of pH on TBA values of ground raw poultry meat. *J. Food Sci.* 46: 1946-1947.
- Chewale, J.G., Takahasi, T. and Tang, J. 1985. The primary structure of cathepsin D and the implications for its biological functions. *In Aspartic proteinases and their inhibitors* (Kostka, V. ed.), pp 101-116. Berlinide Gruyter.
- Chinivasagam, H.N., Bremner, H.A. and Reeves, R. 1998. Can spoilage bacteria cause blackspot (melanosis) in stored prawns? *Lett. Appl. Microbiol.* 27(1): 5-8.
- Chukiatwattana, P. 2002. Partial purification and characterization of protease from shrimp (*Penaeus monodon*) hepatopancreases. Master of Science Thesis in Food Science and Technology. Kasetsart University. 77p.

- Church, N. 1994. Developments in modified atmosphere packaging and related technologies. *Trends Food Sci. Technol.* 5: 345-352.
- Church, N., 1998. MAP fish and crustaceans-sensory enhancement. *Food Sci. Technol. Today.* 12(2): 73-83.
- Chytiri, S., Chouliara, I., Savvaidis, I.N. and Kontominas, M.G. 2004. Microbiological, chemical and sensory assessment of iced whole and filleted aquacultured rainbow trout. *Food Microbiol.* 21: 157-165.
- Cobb, B.F., Alaniz, I. and Thompson, C.A. 1973. Biochemical and microbial studies on shrimp: volatile nitrogen and amino nitrogen analysis. *J. Food Sci.* 38: 431-436.
- Cobb, B.F., Vanderzant, C., Hanna, M.O. and Yeh, C.P. 1976. Effect of ice storage on microbiological and chemical changes in shrimp and melting ice in a model system. *J. Food Sci.* 41: 29-34.
- Cobb, B.F.III. 1977. Biochemistry and physiology of shrimp: effect on use as food. *In Tropical Products Inst. (Proceedings of the Conference on Handling, Processing, Marketing Tropical Fish. Ed.),* pp 405-411. 5-9 July 1976. Tropical Products Inst: London.
- Connor, W.E. and Lin, D.S. 1982. The effects of shellfish in the diet upon the plasma lipid levels in humans. *Metabolism.* 31: 1046-1051.
- Conway, E.T. 1950. *Microdiffusion and volumetric error.* Crosby Lockwood and Son Ltd: London.
- Cousin, M.A., Jay, I.M. and Vasavada, P.C. 1992. Psychrotrophic microorganism. *In Compendium of Methods for the Microbiological Examination of Food (Vanderzand, C. and Splittstoesser, D.F. eds.),* pp. 153-168.
- Daeschel, M.A. 1989. Antimicrobial substances from lactic acid bacteria for the use as food preservatives. *Food Technol.* 43(4): 164-167.
- Dalet-Fumeron, V., Guinec, N. and Pagano, M. 1993. *In vitro* activation of pro-cathepsin B by three serine proteinases: leucocyte elastase, cathepsin G, and the urokinase-type plasminogen activator. *FEBS.* 323: 251-254.
- Dalgaard, P. 1995. Qualitative and quantitative characterization of spoilage bacteria from packed fish. *Int. J. Food Microbiol.* 26: 319-333.

- Dalgaard, P. and Jørgensen, L.V. 2000. Cooked and brined shrimps packed in a modified atmosphere have a shelf-life of >7 months at 0°C, but spoil in 4-6 days at 25°C. *Int. J. Food Sci. Technol.* 35: 431-442.
- Daniels, J.A., Krishnamurthi, R. and Rizvi, S.H. 1985. A review of the effects of carbon dioxide on microbial growth and food quality. *J. Food Protect.* 48: 532-537.
- Davidson, P.M. and Juneja, V. 1990. Antimicrobial agents. *In* Food additives. (Branen, A.L, Davidson, P.M. and Salminen, S. eds.) pp. 83-138. Marcel Dekker: New York.
- Davis, A.R. and Slade, A. 1995. Fate *Aeromonas* and *Yersinia* on modified-atmosphere-packaged (MAP) cod and trout. *Lett. Appl. Microbiol.* 21: 354-358.
- Debevere, J. and Boskou, G. 1996. Effect of modified atmosphere packaging on the TVB/TMA producing microflora of cod fillets. *Int. J. Food Microbiol.* 31: 221-229.
- Debevere, J., Devlieghere, F., Van Sprundel, P. and De Meulenaer, B. 2001. Influence of acetate and CO<sub>2</sub> on the TMAO-reduction reaction by *Shewanella baltica*. *Int. J. Food Microbiol.* 68: 115-123.
- de la Hoz, L., Lopez-Galvez, D.E., Fernandez, M., Hierro, E. and Ordonez, J.A. 2000. Use of carbon dioxide enriched atmospheres in the refrigerated storage (2°C) of salmon (*Salmo salar*) steaks. *Eur. Food Res. Technol.* 210: 179-188.
- Dendinger, J.E. and O'Connor, K.L. 1990. Purification and characterization of a trypsin-like enzyme from the midgut gland of the Atlantic blue crab, *Callinectes sapidus*. *Comp. Biochem. Physiol.* 95B: 525-530.
- Dhananjaya, S. and Stroud, G.D. 1994. Chemical and sensory changes in haddock and herring stored under modified atmosphere. *Int. J. Food Sci. Technol.* 29: 575-583.
- Dionysius, D.A., Hoek, K.S., Milne, J.M. and Slattery, S.L. 1993. Trypsin-like enzyme from sand crab (*Portunus pelagicus*): Purification and characterization. *J. Food Sci.* 58: 780-784, 792.
- Donato, M.D., Manrique, R., Ramirez, R., Mayer, L. and Howell, C., 2005. Mass selection and inbreeding effects on a cultivated strain of *Penaeus (Litopenaeus vannamei)* in Venezuela. *Aquaculture* 247: 159-167.
- Doyle, N.P., Beuchat, L.R. and Montville, T.J. 2001. Food microbiology. (2<sup>nd</sup> ed.). ASM Press: Washington, DC, USA. pp. 383-410.



- Dziezak, J.D. 1986. Preservatives: Antioxidants - The ultimate answer to oxidation. *Food Technol.* 40(9): 94-102.
- Dziezak, J.D. 1990. Phosphates improve many food. *Food Technol.* 44(14): 80-86.
- Eisen, A.Z., Henderson, K.O., Jeffrey, J.J. and Bradshaw, R.A. 1973. A collagenolytic protease from the hepatopancreas of the fiddler crab, *Uca pugilator*, purification and properties. *Biochemistry.* 12: 1814-1822.
- Erickson, M.C., Gordon, D.T. and Anglemier, A.F. 1983. Proteolytic activity in the sarcoplasmic fluids of parasitized Pacific whiting (*Merluccius productus*) and unparasitized ture cod (*Gadus macrocephalus*). *J. Food Sci.* 48: 1315-1319.
- Erkan, N., Ozden, O. and Inugur, M. 2007. The effects of modified atmosphere and vacuum packaging on quality of chub mackerel. *Int. J. Food Sci. Technol.* (Inpress online).
- Etherington, D.J. 1984. The contribution of proteolytic enzymes to postmortem changes in muscle. *J. Anim. Sci.* 59: 1644-1649.
- European Commission. 2003. Opinion of the Scientific Committee on Food on 4-Hexylresorcinol. Health and Consumer Protection Directorate-General (Online). Available [http://europa.eu.int/comm/food/fs/sc/scf/out170\\_en.pdf](http://europa.eu.int/comm/food/fs/sc/scf/out170_en.pdf). (2005. Nov 17).
- FAO. 2007. Fisheries global information system (Online). Available <http://www.fao.org> (2007. June 6).
- Farber, J.M. 1991. Microbiological aspects of modified-atmosphere packaging technology-a review. *J. Food Protect.* 54: 58-70.
- Fatima, R., Khan, M.A. and Qadri, R.B. 1988. Shelf life of shrimp (*Penaeus merguensis*) stored in ice (0°C) and partially frozen (-3°C). *J. Sci. Food Agric.* 42: 235-247.
- Federal Register. 1985. Notice to shippers, distributors, packers, and importers of shrimp containing sulfites. 50: 2957-2958.
- Feng, P.C.S. and Hartman, P.A. 1982. Fluorogenic assays for immediate confirmation of *Escherichia coli*. *Appl. Environ. Microbiol.* 43: 1320-1329.

- Fennema, O.R. 1990. Comparative water-holding properties of various muscle foods. A critical review relating to definitions, methods of measurement, governing factors, comparative data and mechanistic matters. *J. Muscle Foods*. 1: 363-381.
- Ferrer, O.B., Koburger, J.A., Otwell, W.S., Gleeson, R.A., Simpson, B.K. and Marshall, M. 1989. Phenoloxidase from the cuticle of Florida spiny lobster (*Panulirus argus*): Mode of activation and characterization. *J. Food Sci.* 54: 63-67.
- Finne, G. 1982. Modified and controlled atmosphere storage of muscle foods. *Food Technol.* 36(2): 128-133.
- Flores, S.C. and Crawford, D.L. 1973. Postmortem quality changes in iced pacific shrimp, (*Pandalus jordani*). *J. Food Sci.* 38: 575-579.
- Foegeding, E.A., Laneir, T.C. and Hultin, H.O. 1996. Characteristics of edible muscle tissues. *In* Food Comistry. (Fennema, O.R. ed.), pp. 879-942. Marcel Dekker: New York, USA.
- Galgani, F., Benyamin, Y. and Ceccaldi, H.J. 1984. Identification of digestive proteinases of *Penaeus kerathurus* (Forskal): A comparison with *Penaeus japonicus* bate. *Comp. Biochem. Physiol.* 78B: 355-361.
- Galgani, F. and Nagayama, F. 1987. Digestive proteinases in the Japanese lobster *Panulirus japonicus*. *Comp. Biochem. Physiol.* 87B: 889-893.
- Gardner, J., Manohar, S. and Borisenok, W.S. 1991. Method and composition for preserving fresh peeled fruits and vegetables. U.S. Patent 4: 988,523.
- Gates, B.J. and Travis, J. 1969. Isolation and comparative properties of shrimp trypsin. *Biochem.* 8: 4483-4489.
- Geissler, K., Manafi, M., Amoros I. and Alonso, J.L. 2000. Quantitative determination of total coliforms and *Escherichia coli* in marine waters with chromogenic and fluorogenic media. *J. Appl. Microbiol.* 88: 280-285.
- Gibson, D.M. and Davis, H.K. 1995. Fish and shellfish products in sous vide and modified atmosphere packs. *In* Principles of modified atmosphere and sous vide product packaging. (Farber, J.M. and Dodds, K.L. eds.), pp 153-174. Technomic Publishing Co. Inc: Lancaster PA.

- Gildberg, A. 1988. Aspartic proteinase in fishes and aquatic invertebrates. *Comp. Biochem. Physiol.* 91B: 425-435.
- Gill, C.O. and Penney, N. 1986. Packaging conditions for extended storage of chilled dark, firm, dry beef. *Meat Sci.* 18: 41-54.
- Gimenez, B., Roncales, P. and Beltran, J.A. 2004. The effects of natural antioxidants and lighting conditions on the quality characteristics of gilt-head sea bream fillets (*Sparus aurata*) packaged in a modified atmosphere. *J. Sci. Food Agric.* 84: 1053-1060.
- Gokodlu, N., Ozden, O. and Erkan, N. 1998. Physical, chemical and sensory analysis of freshly harvested sardines (*Sardina pilchardus*) stored at 4°C. *J. Aquat. Food Prod. Technol.* 7(2): 5-15.
- Gomez-Guillen, M.C. and Batista, I. 1997. Seasonal changes and preliminary characterization of cathepsin D-like activity in sardine (*Sardina pilchardus*) muscle. *Int. J. Food Sci. Technol.* 32: 255-260.
- Gomez-Guillen, M.C., Martinez-Alvarez, O., Lamas, A. and Montero, P. 2005. Melanosis inhibition and SO<sub>2</sub> residual levels in shrimps (*Parapenaeus longirostris*) after different sulfite-based treatments. *J. Food Sci. Agric.* 85: 14-143-1148.
- Gonçalves, A.C., López-Caballero, M.E. and Nunes, M.L. 2003. Quality changes of deep pink shrimp (*Parapenaeus longirostris*) packed in modified atmosphere. *J. Food Sci.* 68: 2586-2590.
- González, C.J., López-Díaz, T.M., García-López, M.L., Prieto, M. and Otero, A. 1999. Bacterial microflora of wild brown trout (*Salmo trutta*), wild pike (*Esox lucius*) and aquacultured rainbow trout (*Oncorhynchus mykiss*). *J. Food Prot.* 62: 1270-1277.
- Gould, G.W. 2000. Strategies for Food Preservation. *In* The Microbiological Safety and Quality of Food (Lund, B.M., Baird-Parker, T.C. and Gould, G.W. eds.), pp. 19-35. Aspen Publishers: Gaithersburg, M.D.
- Gram, L. and Huss, H.H. 1996. Microbiological spoilage of fish and fish products. *Int. J. Food Microbiol.* 33: 121-137.

- Granicher, D. and Portzehl, H. 1964. The influence of magnesium and calcium pyrophosphate chelates, of free magnesium ions, free calcium ions, and free pyrophosphate ions on the dissociation of actomyosin in solution. *Biochim et Biophys Acta*. 86: 567-578.
- Gray, R.J.H., Hoover, D.G. and Muir, A.M. 1983. Attenuation of microbial growth on modified atmosphere packaged fish. *J. Food Prot.* 46: 610-613.
- Guandalini, E., Ioppolo, A., Mantovani, A., Stacchini, P. and Giovannini, C. 1998. 4-Hexylresorcinol as inhibitor of shrimp melanosis: efficacy and residues studies; evaluation of possible toxic effect in a human intestinal in vitro model (Caco<sup>-2</sup>); preliminary safety assessment. *Food Addit. Contam.* 15: 171-180.
- Hamm, R. 1986. Functional properties of the myofibrillar system and their measurements *In* Muscle as food (Bechtel, P.J. ed), pp. 135-199. Academic Press Inc: San Diego, CA.
- Hammack, T.S., Amaguana, R.M., June, G.A., Sherrod, P.S. and Andrews, W.H. 1999. Relative effectiveness of selenite cystine broth, tetrathionate broth, and Rappaport-Vassiliadis medium for the recovery of *Salmonella* from foods with a low microbial load. *J. Food Prot.* 62: 16-21.
- Hamre, K., Øfsti, A., Naess, T., Nortvedt, R. and Holm, J.C. 2003. Macronutrient composition in formulated diets for Atlantic halibut (*Hippoglossus hippoglossus*, L.) juveniles a multivariate approach. *Aquaculture*. 227: 233-244.
- Hanada, K., Tamai, M., Tamagishi, M., Ohmura, S., Sawada, J., and Tanaka, I. 1978. Isolation and characterization of E-64, a new thiol protease inhibitor. *Agr. Biol. Chem.* 42: 523-528.
- Hebard, C.E., Flick, G.J. and Martin, R.E. 1982. Occurrence and significance of trimethylamine oxide and its derivatives in fish and shellfish. *In* Chemistry and Biochemistry of Marine Food Products (Martin, R.E., Flick, G.J., Hebard, C.E. and Ward, D.R. eds.), pp. 149-304. AVI Publishing Company: Westport, CT.
- Hernandez-Herrero, M.M., Duflos, G., Malle, P. and Bouquelet, S. 2003. Collagenase activity and protein hydrolysis as related to spoilage of iced cod (*Gadus morhua*). *Food Res. Int.* 36: 141-147.

- Heu, M.S., Kim, H.R. and Pyeun, J.H. 1995. Comparison of trypsin and chymotrypsin from the viscera of anchovy, *Engraulis japonica*. *Comp. Biochem. Physiol.* 112B: 557-567.
- Hjelmeland, K. and Raa, J. 1982. Characteristics of two trypsin type isozymes isolated from the Arctic fish capelin (*Mallotus villosus*). *Comp. Biochem. Physiol.* 71B: 557-562.
- Honjo, I., Kimura, S. and Nonaka, M. 1990. Purification and characterization of trypsin-like enzyme from shrimp *Penaeus indicus*. *Nippon Suisan Gakkaishi.* 56: 1627-1634.
- Hoz, L., Lopez-Galez, D. E., Fernandez, M., Hierro, E. and Ordonez, J. A. 2000. Use of carbon dioxide enriched atmospheres in the refrigerated storage (2°C) of salmon (*Salmo salar*) steaks. *Eur. Food Res. Technol.* 210: 179-188.
- Huidobro, A., Lopez-Caballero, M.E. and Mendes, R. 2002. Onboard processing of deepwater pink shrimp (*Parapenaeus longirostris*) with liquid ice: effect on quality. *Eur. Food Res. Technol.* 214: 469-475.
- Huss, H.H. 1971. Prepackaged Fresh Fish. *In* Fish inspection and quality control. (Kreuzer, R. ed.), pp. 60-65. Fishing News (Books) Limited: London.
- Huss, H.H. 1995. Quality and quality changes in fresh fish. *FAO Fishery Technical Paper*, No. 384: 1-195.
- International commission on microbiological specifications for foods (ICMSF). 1986. Sampling plans for fish and shellfish. *In* ICMSF, *Microorganisms in Foods. Sampling for Microbiological Analysis: Principles and Scientific Applications* Vol. 2. 2nd edition. (ICMSF, ed.). University of Toronto Press: Toronto, Canada.
- Ishida, M., Niizelei, S. and Nagayama, F. 1994. Thermostable proteinase in salted anchovy muscle. *J. Food Sci.* 59: 781-785.
- Ita, P.S. and Hutkins, R.W. 1991. Intracellular pH and survival of *Listeria monocytogenes* Scott A in tryptic soy broth containing acetic, lactic, citric, and hydrochloric acids. *J. Food Prot.* 54: 15-19.
- Jiang, S.T., Hev, Y.H., Lee, J.J. and Wang, J.H. 1993. Comparison of the cathepsin D from mackerel (*Scomber australasicus*) and milkfish (*Chanos chanos*) muscle. *Biosci. Biotech. Biochem.* 57: 571-577.

- Jiang, S.T. and Lee, T.C. 1988. Effect of modified ice storage on the quality and prevention of darkening discoloration of shrimp (*Solenocera prominentis*). Bull. Jpn. Soc. Sci. Fish. 54: 1415-1422.
- Jiang, S.T., Lee, B.L., Tsao, C.Y. and Lee, J.J. 1997. Mackerel cathepsin B and L effects on thermal degradation of surimi. J. Food Sci. 62: 310-315.
- Jiang, S.T., Lee, J.J. and Chen, H.C. 1994. Purification and characterization of cathepsin B from ordinary muscle of mackerel (*Scomber australasicus*). J. Agric. Food Chem. 42: 1073-1079.
- Jiang, S.T., Moody, M.W. and Chen, H.C. 1991. Purification and characterization of proteases from digestive tract of grass shrimp (*Penaeus monodon*). J. Food Sci. 56: 322-326.
- Jiang, S.T., Nei, F.P., Chen, H.C. and Wang, J.H. 1992a. Comparative study on the cathepsin D from the banded shrimp (*Penaeus japonicus*) and grass shrimp (*Penaeus monodon*). J. Agric. Food Chem. 40: 961-966.
- Jiang, S.T., Wang, J.H. and Chen, H.C. 1992b. Lysosomal enzyme effects on the postmortem changes in Tilapia (*Tilapia × Tilapia aurea*) muscle myofibrils. J. Food Sci. 57: 277-282.
- Joffraud, J.J., Leroi, F., Roy, C. and Berdague, J.L., 2001. Characterisation of volatile compounds produced by bacteria isolated from the spoilage flora of cold-smoked salmon. Int. J. Food Microbiol. 66: 175-184.
- Kawabata, T., Nishimura, Y., Masahide, H. and Kato, K. 1993. Purification and processing of rat liver procathepsin B. J. Biochem. 133: 389-394.
- Kawamura, Y., Nishimura, K., Igrashi, S., Doi, E. and Yonezawa, D. 1981. Characteristics of autolysis of Antarctic krill. Agric. Biol. Chem. 45: 93-100.
- Kim, C.R., Hearnberger, J.O., Vickery, A.R., White, C.H. and Marshall, D.L. 1995. Extending shelf-life of refrigerated catfish fillets using sodium acetate and monopotassium phosphate. J. Food Prot. 58: 644-647.
- Kim, C.R. and Hearnberger, J.O. 1994. Gram negative bacteria inhibition by lactic acid culture and food preservations on catfish fillets during refrigerated storage. J. Food Sci. 59: 513-516.

- Kim, B.G. 1999. Developmental biology of larval moi, *Polydactylus sexfilis*, and implications for aquaculture. PhD. Thesis, University of Hawaii. 186 pp.
- Kim, H.R., Meyers, S.P. and Godber, J.S. 1992. Purification and characterization of anionic trypsins from hepatopancreas of crayfish, *Procambarus clarkii*. *Comp. Biochem. Physiol.* 103B: 391-398.
- Kim, H.R., Meyers, S.P. and Godber, J.S. 1996a. Anionic trypsins from crayfish hepatopancreas: Effects on protein degradation of tail meat. *J. Food Sci.* 61: 78-80, 96.
- Kim, H.R., Kim, D.S., Ahn, C.B. and Pyeun, J.H. 1996b. Purification and characterization of trypsins affecting on autolysis of shrimp, *Penaeus japonicus*. *J. Korean Fish. Soc.* 29: 797-804.
- Kim, S.K., Jeon, Y.J., Byun, H.G., Kim, Y.T. and Lee, C.K. 1997. Enzymatic recovery of cod frame proteins with crude proteinase from tuna pyloric caeca. *Fish. Sci.* 63: 421-427.
- Kim, Y.T. and Kim S.K. 1991. Purification and characterization of the collagenase from the tissue of filefish, *Novoden modestrus*. *Korean Biochem. J.* 24: 401-409.
- Kimura, B. and Murakami, M. 1993. Fate of food pathogens in gas-packaged jack mackerel fillets. *Nippon Suisan Gakkaishi.* 59: 1163-1169.
- Kirschke, H. and Barrett, A.J. 1987. Chemistry of lysosomal proteinases. *In* Lysosomes: their role in protein breakdown (Glaumann, H. and Ballard, F.J. eds.), pp 193-240. Academic Press: London.
- Kirschke, H., Wood, L., Roisen, F.J. and Bird, J.W. 1983. Activity of lysosomal cysteine proteinase during differentiation of rat skeletal muscle. *J. Biochem.* 214: 871-877.
- Klimova, O.A., Borukhov, S.I., Solovyeva, N.I., Balaevskaya, T.O. and Strongin, A.Y. 1990. The isolation and properties of collagenolytic proteases from crab hepatopancreas. *Biochem. Biophys. Res. Commun.* 166: 1411-1420.
- Kolakowska, A., Czerniejewska-Surma, B. and Gajowiecki-Lienkiewicz, L. 1992. Effect of fishing season on shelf life of iced Baltic herring. *In* Quality Assurance in the Fish Industry. (Huss, H.H., Jakobsen, M. and Liston, J. eds.), pp. 81-91. Elsevier Science Publisher: Amsterdam, The Netherlands.

- Kolodziejska, I. and Sikorski, Z.E. 1996. Neutral and alkaline muscle proteases of marine fish and invertebrates. A review. *J. Food Biochem.* 20: 349-363.
- Konno, K. and Fukazawa, C. 1993. Autolysis of squid mantle muscle protein as affected by storage conditions and inhibitors. *J. Food Sci.* 58: 1198-1202.
- Konosu, S. and Yamaguchi, K. 1982. The flavor components in fish and shellfish. *In Chemistry and biochemistry of marine food products.* (Martin, R.E., Flick, G.J., Hebard, C.E. and Ward, D.R. eds.), pp 367-404. AVI Publishing Co: Westport, Conn.
- Kristjansson, M.M., Guthmundsdottir, S., Fox, J.W. and Bjarnason, J.B. 1995. Characterization of collagenolytic serine proteinase from the Atlantic cod (*Gadus morhua*). *Comp. Biochem. Physiol.* 110B: 707-717.
- Kubo, I. and Kinst-Hori, I. 1998. Tyrosinase inhibitors from cumin. *J. Agric. Food Chem.* 46: 5338-5341.
- Laakonen, E., Sherbon, J.W. and Wellington, G.H. 1970. Low temperature, long-term heating of bovine muscle. *J. Food Sci.* 35: 181-183.
- Lakshmanan, R., Shakila, R.J. and Jeyasekaran, G. 2002. Survival of amine forming bacteria during the ice storage of fish and shrimp. *Food Microbiol.* 19: 617-625.
- Lalitha, K.V. and Surendran, P.K. 2006. Microbiological changes in farm reared freshwater prawn (*Macrobrachium rosenbergii* de Man) in ice. *Food Control.* 17: 802-807.
- Lannelongue, M., Finne, G., Hanna, M., Nickelson, R. and Vanderzant, G. 1982. Storage characteristics of brown shrimp (*Penaeus aztecus*) stored in retail packages containing CO<sub>2</sub>-enriched atmospheres. *J. Food Sci.* 47: 911-913, 923.
- Layrisse, M. and Matches, J.R. 1984. Microbiological and chemical changes of spotted shrimp (*Pandalus platyceros*) stored under modified atmospheres. *J. Food Prot.* 47: 453-437.
- Lee, Y.C. and Um, Y.S. 1995. Quality determination of shrimp (*Penaeus japonicus*) during iced and frozen storage. *Korean J. Food Sci. Technol.* 27: 520-524.
- Leitao, M.F.F. and Rios, D.P.A. 2000. Microbiological and chemical changes in freshwater prawn (*Macrobrachium rosenbergii*) stored under refrigeration. *Brazilian J. Microbiol.* 31: 178-183.



- Liao, M.L. and Seib, P.A. 1988. Chemistry of L-ascorbic acid related to foods. *Food Chem.* 30: 289-312.
- Lin, K.W. and Chuang, C.H. 2001. Effectiveness of dipping with phosphate, lactate and acetic acid solutions on the quality and shelf-life of pork loin chop. *J. Food Sci.* 66:494-498.
- Lindner, P., Angel, S., Weinberg, Z.G. and Granit, R. 1988. Factors inducing mushiness in stored prawns. *Food Chem.* 29: 119-132.
- Lindgren, S.E. and Dobrogosz, W.J. 1990. Antagonistic activities of lactic acid bacteria in food and feed fermentations. *FEMS Microbiol. Rev.* 87: 149-164.
- Lopez-Galvez, D.E., Dela-Hoz, L., Blanco, M. and Ordonez, A., 1998. Refrigerated storage (2°C) of sole (*Solea solea*) fillets under CO<sub>2</sub> enriched atmospheres. *J. Agric. Food Chem.* 46: 1143-1149.
- Lopez-Galvez, D.E., Dela-Hoz, L. and Ordonez, J.A. 1995. Effect of carbon-dioxide and oxygen-enriched atmospheres on microbiological changes and chemical changes in refrigerated tuna (*thunnus-alalunga*) steaks. *J. Agric. Food Chem.* 43: 483- 490.
- López-Caballero, M.E., Gonçalves, A.C. and Nunes, M.L. 2002. Effect of CO<sub>2</sub>/O<sub>2</sub>-containing modified atmospheres on packed deepwater pink shrimp (*Parapenaeus longirostris*). *Eur. Food Res. Technol.* 214: 192-197.
- Lopez-Caballero, M.E., Perez-Mateos, M., Borderias, J.E. and Montero, P. 2000. Extension of the shelf life of prawns (*Penaeus japonicus*) vacuum packaging and high-pressure treatment. *J. Food Prot.* 63: 1381-1388.
- Lopez-Caballero, M.E., Perez-Mateos, M., Borderias, J.E. and Montero, P. 2006. Quality of Norway lobster (*Nephrops norvegicus*) treated with a 4-hexylresorcinol-based formulation. *Eur. Food Res. Technol.* 222: 425-431.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193: 256-275.
- Lu, P.J., Liu, H.Ch. and Tsai, I.H. 1990. The midgut trypsins of shrimp (*Penaeus monodon*). High efficiency toward native protein substrates including collagens. *Biol. Chem. Hoppe-Seyler.* 371: 851-859.

- Mach, L., Schwihla, H., Stuwe, K., Rowan, A.D., Mort, J.S. and Glossl, J. 1993. Activation of procathepsin B in human hepatoma cells: its conversion into the mature enzyme relies on the action of cathepsin B itself. *J. Biochem.* 293: 437-442.
- Madero, C.F. and Finne, G. 1982. Properties of phenoloxidase isolated from gulf shrimp. *Proceedings of the Seventh Annual Tropical and Subtropical Fisheries Technological Conference of the Americas*, p. 328-339. New Orleans: LA.
- Majchrzak, D., Mitter, S. and Elmadfa, I. 2004. The effect of ascorbic acid on total antioxidant activity of black and green teas. *Food Chem.* 88: 447-451.
- Maldhavi, D.L., Deshpande, S.S. and Salunkhe, D.K. 1995. *Food Antioxidants*. Marcel Dekker: Inc. New York.
- Markinodan, Y. and Ikeda, S. 1969. Studies on fish muscle protease-III. Purification and properties of a proteinase active in acid pH range. *Bull. Jpn. Soc. Sci. Fish* 35: 758-766.
- Markis, P. and Embs, R.J. 1966. Effect of sulfite and ascorbic on mushroom phenol oxidase. *J. Food Sci.* 31: 807-811.
- Marshall, D.L., Wiese-Lehigh, P.L., Well, J.H. and Farr, A.J. 1991. Comparative growth of *Listeria monocytogenes* and *Pseudomonas fluorescens* on precooked chicken nuggets stored under modified atmosphere. *J. Food Prot.* 54: 841-843.
- Marshall, D.L. and Jindal, V. 1997. Microbiological quality of catfish frames treated with selected phosphate. *J. Food Prot.* 60: 1081-1083.
- Marshall, M.R., Simpson, B.K., Ferrer, O. and Otwell, W.S. 1988. Purification and characterization of phenoloxidases in crustaceans: Shrimp and lobster. Report of the Florida Sea Grant College Program. pp. 1-12.
- Marshall, M., Kim, J.R. and Wei, C.I. 2000. Enzymatic Browning in Fruits, Vegetables and Seafoods. FAO (Online). Available [www.fao.org](http://www.fao.org) (2007. June 6).
- Martínez-Alvarez, O., Gomez-Guillen, M.C. and Montero, P. 2005a. Role of sulfites and 4-hexylresorcinol on microbial growth and melanosis prevention in shrimps using controlled atmosphere. *J. Food Prot.* 68: 103-110.

- Martiez-Álvarez, O., Lopez-Caballero, M.E., Montero, P. and Gomez-Guillen, M.C. 2005b. A 4-hexylresorcinol-based formulation to prevent melanosis and microbial growth in chilled tiger prawns (*Marsupenaeus japonicus*) from Aquaculture. *J. Food Sci.* 70: 415-422.
- Martiez-Álvarez, O., Montero, P. and Gomez-Guillen, M.C. 2004. Controlled atmosphere as coadjuvant to chilled storage for prevention of melanosis in shrimp (*Parapenaeus longirostris*). *Eur. Food Res. Technol.* 220: 125-130.
- Martinez, M.V. and Whitaker, J.R. 1995. The biochemistry and control of enzymatic browning. *Trends in Food Sci. Technol.* 6(6): 195-200.
- Masniyom, P., Benjakul, S. and Visessanguan, W. 2002. Shelf-life extension of refrigerated seabass slices under modified atmosphere packaging. *J. Sci. Food Agric.* 82: 873-880.
- Masniyom, P., Benjakul, S. and Vissanguan, W. 2004. ATPase activity, surface hydrophobicity, sulfhydryl content and protein degradation in refrigerated seabass muscle in modified atmosphere packaging. *J. Food Chem.* 28: 43-60.
- Masniyom, P., Benjakul, S. and Visessanguan, W. 2005. Combination effect of phosphate and modified atmosphere on quality and shelf-life extension of refrigerated seabass slices. *LWT-Food Sci. Technol.* 38: 745-756.
- Masniyom, P., Benjakul, S. and Visessanguan, W. 2006. Synergistic antimicrobial effect of pyrophosphate on *Listeria monocytogenes* and *Escherichia coli* O157 in modified atmosphere packaged and refrigerated seabass slices. *LWT-Food Sci. Technol.* 39: 302-307.
- Matches, J.R. and Layrisse, M.E. 1985. Controlled atmosphere storage of spotted shrimp (*Pandalus platyceros*). *J. Food Prot.* 48: 709-711.
- McEvily, A.J., Iyengar, R. and Otwell, W.S. 1991. 4-Hexylresorcinol controls enzymatic browning in shrimp and has potential for application in a variety of other foods and beverages. *Food Technol.* 45(9): 80-86.
- McEvily, A.J., Iyengar, R. and Otwell, W.S. 1992. Inhibition of enzymatic browning in foods and beverages. *Crit. Rev. Food Sci. Nutr.* 32: 253-273.

- Mendes, R., Goncalves, A. and Nunes, M.L. 1999. Changes in free amino acids and biogenic amines during ripening of fresh and frozen sardine. *J. Food Biochem.* 23: 295-306.
- Mendes, R., Huidobro, A. and López-Caballero, E. 2002. Indole levels in deepwater pink shrimp (*Parapenaeus longirostris*). *Eur. Food Res. Technol.* 214: 125-130.
- Mihalyi, E. 1978. Application of proteolytic enzymes to protein structure studies. CRC Press, Inc: West Palm Beach, Fla.
- Ming-Lang, H., Hsiu-Ho, C. and Shann-Tzong, J. 1986. Effect of modified ice storage on the shelf-life of shrimp. *Bull. Jap. Soc. Sci. Fish.* 52: 479-488.
- Modderman, J.P. 1986. Technological aspects of use of sulfiting agents in food. *J. Assoc. Off. Anal. Chem.* 69: 1-3.
- Molins, R.A. 1991. Phosphates in Food. CRC Press, Boca Raton, FL.
- Montero, P., Avalos, A. and Perez-Mateos, M. 2001a. Characterization of polyphenoloxidase of prawns (*Penaeus japonicus*). Alternatives to inhibition: additives and high-pressure treatment. *Food Chem.* 75: 317-324.
- Montero, P., Lopez-Caballero, M.E. and Pérez-Mateos, M. 2001b. The effect of inhibitors and high pressure treatment to prevent melanosis and microbial growth on chilled prawns (*Penaeus japonicus*). *J. Food Sci.* 66: 1201-1206.
- Montero, P., Martínez-Alvarez, O. and Gómez-Guillén, M.C. 2004. Effectiveness of onboard application of 4-hexylresorcinol in inhibiting melanosis in shrimp (*Parapenaeus longirostris*). *J. Food Sci.* 68: 643-647.
- Montero, P., Martínez-Alvarez, O., Zamorano, J.P., Alique, R. and Gómez-Guillén, M.C. 2006. Melanosis inhibition and 4-hexylresorcinol residual levels in deepwater pink shrimp (*Parapenaeus longirostris*) following various treatments. *Eur. Food Res. Technol.* 223: 16-21.
- Mort, J.S. and Buttle, D.J. 1997. Molecules in focus. Cathepsin B. *Int. J. Biochem. Cell Biol.* 29(5): 715-720.
- Muncy, R.J. 1984. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (south Atlantic): white shrimp. U.S. Fish and Wildl. Serv. FWS/OBS-82/11.27. U.S. Army Corps of Engineers, TR EL-82-4. 19 pp.

- Murakami, K. and Noda, M. 1981. Study on proteases from the digestive organs of sardine. I. Purification and characterization of three alkaline proteinases from the pyloric caeca. *Biochim. Biophys. Acta.* 658: 17-26.
- Muramatsu, T. and Kariuchi, H. 1978. Proteolytic activity of extracts from the hepatopancreas of shrimp, *Parahaliporus sibogae* (De Man). *Bull. Jap. Soc. Sci. Fish.* 44: 171-174.
- Muriana, F.J.G., Ruiz-Gaterrez, V., Gallardo, M.L. and Minguez-Mosquera, M.I. 1993. A study of the lipids and carotenoprotein in the prawn *Penaeus japonicus*. *J. Biochem.* 114: 223-229.
- Nash, T. 1953. The colorimetric estimation of formaldehyde by means of the Hantzsch reaction. *J. Biochem.* 55: 416-419.
- Nawar, W.W. 1996. Lipids. *In Food Chemistry* (Fennema, O. ed.), pp. 225-319. Marcel Dekker: New York.
- Nip, W.K., Lan, C.Y. and Moy, J.H. 1985. Partial characterization of a collagenolytic enzyme fraction from the hepatopancreas of the freshwater prawn, *Macrobrachium rosenbergii*. *J. Food Sci.* 50: 1187-1188.
- Nip, W.K. and Moy, J.H. 1988. Microstructural Changes of ice-chilled and cooked freshwater prawn, *Macrobrachium rosenbergii*. *J. Food Sci.* 53: 319-322.
- Nishimoto, J. 1985. Study on the denaturation of Crucian (*Carassius auratus*) myofibrillar protein during frozen storage. *Bull. Jap. Soc. Sci. Fish.* 51: 1887-1892.
- Nunes, M., Batista, I. and Morão de Campos, R. 1992. Physical, chemical and sensory analysis of sardine (*Sardina pilchardus*) stored in ice. *J. Sci. Food Agric.* 59: 37-43.
- Nur-E-Borhan, S.A., Okada, S. Watabe, S. and Yamaguchi, K. 1995. Carotenoproteins from the exoskeleton of commercial black tiger prawn. *Fish. Sci.* 60: 213-215.
- Offer, G. and Knight, P. 1988. The structural basis of water-holding in meat. *In Developments in meat science 4.* (Lawrie, R. ed.), pp 63-234. Elsevier: London.
- Ogawa, M., Kurotsu, M., Ochiai, I. and Kozima, T. 1983. Mechanism of black discoloration in spiny lobster tails stored in ice. *Bull. Jpn. Soc. Sci. Fish/Nippon Suisan Gakkaishi.* 49: 1065-1075.
- Ogawa, M., Perdugao, N.B., Santiago, M.E. and Kozima, T.T. 1984. On physiological aspects of black spot appearance in shrimp. *Bull. Jap. Soc. Sci. Fish.* 50: 1763-1769.

- Ogrydziak, D.M. and Brown, W.D. 1982. Temperature effects in modified atmosphere storage of seafoods. *Food Technol.* 36(5): 86-96.
- Oh, E.S., Kim, D.S., Kim, J.H. and Kim, J.H. 2000. Purification and characterization of protease from the hepatopancreas of shrimp, *Penaeus orientalis*. *J. Food. Biochem.* 24: 251-264.
- Olalla, A., Suna, C., Sabastain, J.A. and Sillero, M.G. 1978. Purification and properties of three proteases from the larvae of the brine shrimp *Artemia salina*. *Biochim. Biophys. Acta.* 523: 181-186.
- Ordóñez, J.A., López-Gálvez, D.E., Fernández, M., Hierro, E. and de la Hoz, L. 2000. Microbial and physicochemical modifications of hake (*Merluccius merluccius*) steaks stored under carbon dioxide enriched atmospheres. *J. Sci. Food Agric.* 80: 1831-1840.
- Ordóñez, J.A., Pablo, Castro, B.F., Asensio, M.A. and Sanz, B. 1991. Selected chemical and microbiological change in refrigerated pork stored in carbon dioxide and oxygen enriched atmosphere. *J. Agric. Food Chem.* 39: 668-672.
- Osnes, K.K. and Mohr, V. 1985. On the purification and characterization of three anionic, serine-type peptide hydrolases from Antarctic krill, *Euphuusiu superba*. *Comp. Biochem. Physiol.* 82B: 607-619.
- Otwell, W.S., Iyengar, R. and McEvily, A.J. 1992. Inhibition of shrimp melanosis by 4-hexylresorcinol. *J. Aqua. Food Prod. Technol.* 1(1): 53-65.
- Otwell, W.S. and Marshall, M.R. 1986. Studies on the use of sulfites to control shrimp melanosis (blackspot): Screen alternatives to sulfiting agents to control shrimp melanosis. Florida seagrant technical paper No. 46: 1-10.
- Ozogul, F., Polat, A. and Ozogul, Y. 2004. The effects of modified atmosphere packaging and vacuum packaging on chemical, sensory and microbiological changes of sardines (*Sardina pilchardus*). *Food Chem.* 85: 49-57.
- Ozogul, F., Taylor, K.D.A., Quantick, P. and Ozogul, Y. 2002. Changes in biogenic amines in herring stored under modified atmosphere and vacuum pack. *J. Food Sci.* 67: 2497-2501.

- Pacheco-Aguolar, R., Lugo-Sanchez, M.E. and Robles-Burgueno, M.R. 2000. Post-mortem biochemical and functional characteristic of Monterey sardine muscle stored at 0°C. *J. Food Sci.* 65: 40-47.
- Parkin, K.L., Wells, M.J. and Brown, W.D. 1981. Modified atmosphere storage of rockfish fillets. *J. Food Sci.* 47: 181-184.
- Parry, R.T. 1993. Introduction. *In Principles and applications of modified atmosphere packaging of food.* (Parry, R.T. ed.), pp 1-18. Blackie: London.
- Parsons, N. and Knight, P. 1990. Origin of variable extraction of myosin from myofibrils treated with salt and pyrophosphate. *J. Sci. Food Agric.* 51: 71-90.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, M.L. 1996a. Effect of modified atmosphere packaging on shelf life of iced fresh hake slices. *J. Sci. Food Agric.* 71: 541-547.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, M.L. 1996b. Effect of carbon dioxide atmosphere on microbial growth and quality of salmon slices. *J. Sci. Food Agric.* 72: 348-352.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, M.L. 1998. Influence of sodium chloride and modified atmosphere packaging on microbiological, chemical and sensorial properties in ice storage of slices of hake (*Merluccius merluccius*). *Food Chem.* 61: 23-28.
- Patsias, A., Chouliara, I., Badeka, A., Savvaidis, I.N. and Kontominas, M.G. 2006. Shelf-life of a chilled precooked chicken product stored in air and under modified atmospheres: microbiological, chemical, sensory attributes. *Food Microbiol.* 23: 423-429.
- Pavlisko, A., Rial, A. and Coppes, Z. 1997. Characterization of trypsin purified from the pyloric ceca of the southwest Atlantic white croaker *Micropogonias furnieri* (Sciaenidae). *J. Food Biochem.* 21: 383-400.
- Pedrosa-Menabrito, A. and Regenstein, J.M. 1988. Shelf life extension of fresh fish: A review. Part. 1. Spoilage of fish. *J. Food Quality.* 11: 117-127.
- Pérez-Gilabent, M. and García Carmona, F. 2000. Characterization of catecholase and cresolase activities of eggplant polyphenol oxidase. *J. Agric. Food Chem.* 48: 695-700.

- Pizzocaro, F., Torreggiani, D. and Gilardi, G. 1993. Inhibition of apple polyphenoloxidase (PPO) by ascorbic acid, citric acid and sodium chloride. *J. Food Process. Pres.* 17: 21-30.
- Ramírez, E.C., Whitaker, J.R. and Virador, V. 2003. Polyphenol oxidase. *In Handbook of food enzymology.* (Whitaker, J.R., Voragen, A.G.J. and Wong, D.W.S. eds.), pp 509-23. Marcel Dekker: New York.
- Randell, K., Ahvenainen, R. and Hattula, T. 1995. Effect of the gas/product ratio and CO<sub>2</sub> concentration on the shelf-life of MA packed fish. *Packag. Technol. Sci.* 205-218.
- Reddy, N.R., Armstrong, D.J., Rhodehamel, E.J. and Kauter, D.A. 1992a. Shelf-life extension and safety concerns about fresh fishery products packaged under modified atmospheres: a review. *J. Food Safety.* 12: 87-118.
- Reddy, N.R., Armstrong, D.J., Rhodehamel, E.J. and Kauter, D.A. 1992b. Shelf-life extension and safety concerns about fresh fishery products packaged under modified atmospheres. A review of the effects of carbon dioxide on microbial growth and food quality. *J. Food Prot.* 38: 303-305.
- Reddy, N.R., Paradis, A., Roman, M.G., Solomon, H.M. and Rhodehamel, E.J. 1996. Toxin development by *Clostridium botulinum* in modified atmosphere packaged fresh tilapia fillets during storage. *J. Food Sci.* 61: 632-635.
- Reddy, N.R., Villanueva, M. and Kauter, D.A. 1995. Shelf life of modified atmosphere packaged fresh tilapia fillets stored under refrigeration and temperature abuse conditions. *J. Food Prot.* 58: 908-914.
- Reilly, A., Bernate, M.A. and Dangla, E. 1984. Storage stability of brackishwater prawns during processing for export. *Food Technol. Aust.* 36: 282-286.
- Reilly, A. and Dangla, E. 1986. Postharvest spoilage of shrimp (*Penaeus monodon*). *In The First Asian Fisheries Society.* (Maclean, J.L. Dizon, L.B.. and Hosillo, L.V. eds.), pp 455-458. Manila, Philippines.
- Rennerre, M. Dumont, F. and Gatellier, P. 1996. Antioxidant enzyme activities in beef in relation to oxidation of lipid and myoglobin. *Meat Sci.* 43: 111-121.



- Rhee, K.S., Krahl, L.M., Lucia, L.M. and Acuff, G.R. 1997. Antioxidation/antimicrobial effect and TBARS in aerobically refrigerated beef as related to microbial growth. *J. Food Sci.* 62: 1205-1210.
- Richardson, T. and Hyslop, D.B. 1985. Enzymes. *In Food Chemistry.* (Fennema, O.R. ed.), pp 371-476. Marcel Dekker: New York.
- Ringo, E., Bendiksen, H.R., Wesmajervi, M.S., Olsen, R.E. Jansen, P.A. and Mikkelsen, H. 2000. Lactic acid bacteria associated with digestive tract of Atlantic salmon (*Salmo salar* L.). *J. Appl. Microbiol.* 89: 317-322.
- Rolle, R.S., Guizani, N., Chen, J.S., Marshall, M.R., Yang, J.S. and Wei, C.I. 1991. Purification and characterization of phenoloxidase from Taiwanese black tiger shrimp (*Penaeus monodon*). *J. Food Biochem.* 15: 17-32.
- Rowan, A.D., Mason, P., Mach, L. and Mort, J.S. 1992. Rat procathepsin B. *J. Biol. Chem.* 267: 15993-15999.
- Roy, P., Colast, B. and Durand, P. 1996. Purification, kinetical and molecular characterizations of a serine collagenolytic protease from greenshore crab (*Carcinus maenas*) digestive gland. *Comp. Biochem. Physiol.* 115B: 87-95.
- Rudenskaya, G.N. 2003. Brachyurins, serine collagenolytic enzymes from crabs. *Russ. J. Bioorg. Chem.* 29: 101-111.
- Ruiz-Capillas, C. and Moral, A. 2002. Residual effect of CO<sub>2</sub> on hake (*Merluccius merluccius* L) as a function of the period of time in controlled atmosphere storage. *J. Sci. Food Agric.* 82: 375-379.
- Ryder, J.M., Buisson, D.H., Scott, D.N. and Fletcher, G.C. 1984. Storage of New Zealand jack mackerel (*Trachurus novaezelandiae*) in ice: chemical, microbiological and sensory assessment. *J. Food Sci.* 49: 1453-1456.
- Sapers, G.M. and Hicks, K.B. 1989. Inhibition of enzymatic browning in fruits and vegetables. *In Quality Factors of Fruits and Vegetables: Chemistry and Technology.* (Jen, J.J. ed.), pp 29-43. American Chemical Society: Washington, DC.
- Saoud, I.P., Davis, D.A. and Rouse, D.B., 2003. Suitability studies of inland well waters for *Litopenaeus vannamei* culture. *Aquaculture.* 217: 373-383.

- Sasi, M., Jeyasekaran, G., Shanmugam, S.A. and Jeya, S.R. 2003. Evaluation of the quality of seerfish (*Scomberomorus commersonii*) stored in dry ice (solid carbon dioxide). *J. Aquat. Food Prod. Technol.* 12(2): 61-72.
- Sato, H., Takino, T. and Okada, Y. 1994. A matrix metalloproteinase expressed on the surface of invasive tumour cells. *Nature.* 370: 61-65.
- Sato, K., Ohashi, C., Ohtsuki, K. and Kawabata, M. 1991. Type V collagen in trout (*Salmo gairdneri*) muscle and its solubility change during chilled storage of muscle. *J. Agric. Food Chem.* 39: 1222-1225.
- Sayavedra-Soto, L.A. and Montgomery, M.W. 1986. Inhibition of polyphenoloxidase by sulfite. *J. Food Sci.* 51: 1531-1536.
- Schnitzer, E., Pinchuk, I., Bor, A., Leikin-Frenkel, A. and Lichtenberg, D. 2007. Oxidation of liposomal cholesterol and its effect on phospholipid peroxidation. *Chem. Phys. Lipids.* 146: 43-53.
- Scullen, O.J. and Zaika, L.L. 1994. Effect of temperature, salt and pH on growth inhibition of *Listeria monocytogenes* by sodium polyphosphate. Presented at 18<sup>th</sup> annual meeting of IAMFES, San Antonio, TX, July 31-August 3.
- Segel, I.H. 1976. Biochemical calculations. pp 273-277. John Wiley and Sons: New York.
- Seymour, T.A., Morrissey, M.T., Peters, M.Y. and An, H. 1994. Purification and characterization of Pacific whiting proteases. *J. Agric. Food Chem.* 42: 2421-2427.
- Shahidi, F. 1994. Seafood proteins and preparation of protein concentrates. *In* Seafoods: chemistry, processing technology and quality. (Shahidi, F. and Botta, J.R. eds.), pp 3-9. Blackie Academic and Professional: London.
- Shamshad, S.I., Kher, U.N., Riaz, M., Zuberi, R. and Qadri, R.B. 1990. Shelf life of shrimp (*Penaeus merguensis*) stored at different temperatures. *J. Food Sci.* 55: 1201-1205.
- Shenouda, S.Y.K. 1980. Theories of protein denaturation during frozen storage of fish flesh. *Adv. Food Res.* 26: 275-311.
- Sikorski, Z.E., Kolakowska, K. and Burt, J.R. 1990. Postharvest, biochemical and microbial changes. *In* Seafood: resources, nutritional composition, and preservation. (Sikorski, Z.E. Ed.), pp 55-75. CRC Press: Boca Raton.

- Silla-Santos, S.M.H. 1996. Biogenic amines: their importance in foods. *Int. J. Food Microbiol.* 29: 213-231.
- Silliker, J.H. and Wolfe, S.K. 1980. Microbiological safety considerations in controlled-atmosphere storage of meats. *Food Technol.* 34(3): 59-63.
- Simpson, B.K. 2000. Digestive proteinases from marine animals. *In* *Seafood Enzymes: Utilization and Influence on Postharvest Seafood Quality.* (Haard, N.F. and Simpson, B.K. eds.), pp 531-540. Marcel Dekker: Newyork.
- Simpson, B.K. and Haard, H.F. 1984. Trypsin from Greenland cod (*Gadus ogac*). Isolation and comparative properties. *Comp. Biochem. Physiol.* 79B: 613-622.
- Simpson, B.K., Marshall, M.R. and Otwell, W.S. 1987. Phenoloxidase from shrimp (*Penaeus setiferus*): purification and some properties. *J. Agric. Food Chem.* 35: 918-921.
- Simpson, B.K., Marshall, M.R. and Otwell, W.S. 1988. Phenoloxidase from pink and white shrimp: Kinetic and other properties. *J. Food Biochem.* 12: 205-217.
- Sinnhuber, R.O. and Yu, T.C. 1958. 2-Thiobarbituric acid method for the measurement of rancidity in fishery products. II. The quantitative determination of malonaldehyde. *Food Technol.* 12(1): 9-12.
- Sivertsvik, M. and Birkeland, S. 2006. Effects of soluble gas stabilisation, modified atmosphere, gas to product volume ratio and storage on the microbiological and sensory characteristics of ready-to-eat shrimp (*Pandalus borealis*). *Food Sci. Technol. Int.* 12: 445-454.
- Sivertsvik, M., Jeksrud, W.K. and Rosnes, J.T. 2002. A review of modified atmosphere packaging of fish and fishery products-significance of microbial growth, activities and safety. *Int. J. Food Sci. Technol.* 37: 107-127.
- Skandamis, P.N. and Nychas, G.J.E. 2002. Preservation of fresh meat with active and modified atmosphere packaging conditions. *Int. J. Food Microbiol.* 79: 35-46.
- Slattery, S.L., Williams, D.J. and Cusak, A. 1995. A sulphite-free treatment inhibits black-spot formation in prawns. *Food Aust.* 47(11): 509-514.
- Smith, J.P., Ramaswanmy, H.S. and Simpson, B.K. 1990. Developments in food packaging technology. Part II: Storage aspects. *Trends Food Sci. Technol.* 1: 111-118.
- Sofos, J.N. 1986. Use of phosphate in low-sodium meat products. *Food Technol.* 40(9): 52-69.

- Sovik, S.L. and Rustad, T. 2006. Effect of season and fishing ground on the activity of cathepsin B and collagenase in by-products from cod species. *LWT-Food Sci. Technol.* 39: 43-53.
- Speck, M.L. 1976. *Compendium of methods for the microbiological examination of foods.* American Publish Health Association, Inc: Washington, D.C.
- Sriket, P., Benjakul, S, Visessanguan, W. and Kijroongrojana, K. 2007. Comparative studies on chemical composition and thermal properties of black tiger shrimp (*Penaeus monodon*) and white shrimp (*Penaeus vannamei*) meats. *Food Chem.* 103: 1199-1207.
- Stammen, K., Gerdes, D. and Caporaso, F. 1990. Modified atmosphere packaging of seafood. *Crit. Rev. Food Sci. Nutr.* 29: 301-331.
- Steel, R.G.D. and Torrie, J.H. 1980. *Principles and procedures of Statistics: a biometrical approach.* 2<sup>nd</sup> ed. P. 862. McGraw. Hill. New York.
- Stenstrom, I. 1985. Microbial flora of cod fillets (*Gadus morhua*) stored at 2°C in different mixture of carbon dioxide and nitrogen/oxygen. *J. Food Prot.* 48: 585-589.
- Stiles, M.E. and Holzapfel, W.H. 1997. Lactic acid bacteria of foods and their current taxonomy. *Int. J. Food Microbiol.* 36: 1-29.
- Suphamongkhon, K. 2002. Office of the Thai trade representative. Thai trade representative Thailand's policy on international trade and investment after the 1997 economic crisis for the international Boston seafood show, Massachusetts Boston, 12 March.
- Taoukis, P.S., Labuza, T.P., Lillemo, J.H. and Lin, S.W. 1990. Inhibition of shrimp melanosis (black spot) by ficin. *LWT-Food Sci. Technol.* 23: 52-54.
- Teruel, S.R.L. and Simpson, B.K. 1995. Characterization of the collagenolytic enzyme fraction from winter flounder (*Pseudopleuronectes americanus*). *Comp. Biochem. Physiol.* 112B: 131-136.
- Tipton, K.F. and Dixon, H.B.F. 1983. Effect of pH on enzymes. *In Contemporary Enzyme Kinetics and Mechanism.* (Purich, D.L. ed.), p. 97-148. Academic Press: New York.

- Toussaint, O. and Lerch, K. 1987. Catalytic oxidation of 2-aminophenols and ortho hydroxylation of aromatic amines by tyrosinase. *Biochem.* 26: 8567-8571.
- Trout, G.R. and Schmidt, G.R. 1984. Effect of phosphate type and concentration, salt level and method of preparation on binding of restructured beef rolls. *J. Food Sci.* 49: 687-694.
- Trout, G.R. and Schmidt, G. R. 1986. Effect of phosphates on the functional properties of restructures beef rolls: the role of pH, ionic strength and phosphate type. *J. Food Sci.* 51: 1416-1423.
- Turkiewicz, M., Galas, E. and Kalinowska, H. 1991. Collagenolytic serine proteinase from Antarctic krill (*Euphasia superba*). *Comp. Biochem. Physiol.* 99B: 359-371.
- Umezawa, H. 1976. Structures and activities of protease inhibitors of microbial origin. *Methods Enzymol.* 45: 678-695.
- Ueno, R., Morishita, T. and Takahashi, T. 1981. Intracellular distribution of enzymes and particle properties of lysosomes in carp muscle tissue. *Bull. Jpn. Soc. Sci. Fish.* 47: 207-214.
- Ueno, R., Liston, J. and Horiguchi, Y. 1986. Intracellular distribution of enzymes and particle properties of lysosomes in mackerel muscle tissue. *Bull. Jpn. Soc. Sci. Fish.* 52: 895-900.
- Van Wormhoudt, A., Le. Chevalier, P. and Sellos, D. 1992. Purification, biochemical characterization and N-terminal sequence of a serine-protease with chymotryptic and collagenolytic activities in a tropical shrimp, *Penaeus vannamei* (*Crustacea, Decapoda*). *Comp. Biochem. Physiol.* 103B: 675-680.
- Vega-Villasante, F., Nolasco, H. and Civera, R. 1995. The digestive enzymes of the Pacific brown shrimp (*Penaeus californiensis*) II. Properties of protease activity in the whole digestive tract. *Comp. Biochem. Physiol.* 112B: 123-129.
- Venugopal, B. and Bailey, M.E. 1978. Lysosomal proteinases in muscle tissue and leukocytes of meat animals. *Meat Sci.* 2: 227-239.
- Venugopal, V., Alur, M.D. and Lewis, N.F. 1983. Extracellular protease from *Pseudomonas marinoglutinosa*: some properties and its action of fish actomyosin. *J. Food Sci.* 48: 671-675.

- Villemure, G., Simard, R.F. and Picard, G. 1986. Bulk storage of cod fillets and gutted cod (*Gadus morhua*) under carbon dioxide atmosphere. *J. Food Sci.* 51: 317-320.
- Vinayakam A. and Nellaiappan K. 1987. Multiple forms of phenoloxidase in the cuticle of different moult stages of the mole crab, *Emerita emeritus* (L., 1767) (Decapoda, Anomura, Hippidea). *Crustaceana* 53: 200-204.
- Visessanguan, W., Menino, A.R., Kim, S.M. and An, H. 2001. Cathepsin L: A predominant heat-activated proteinase in arrowtooth flounder muscle. *J. Agric. Food Chem.* 49: 2633-2640.
- Wager, M.K. and Busta, F.F. 1985. Inhibition of *Clostridium botulinum* 52A toxicity and protease activity by sodium acid pyrophosphate in meat systems. *Appl. Environ. Microbiol.* 50: 16-20.
- Wagner, T. and Finne, G. 1986. Evaluation of alternatives to sulfiting agents as melanosis inhibitors in raw shrimp. Technical Paper of the Florida Sea Grant Programme 45NT: 2.
- Walker, J.R.L. 1977. Enzymatic browning in foods. Its chemistry and control. *Food Technol. NZ.* 12: 19-25.
- Wang, M. Y. and Brown, W. D. 1983. Effects of elevated CO<sub>2</sub> atmosphere on storage of fresh water crayfish (*Pacifastacus leniusculus*). *J. Food Sci.* 48: 158-162.
- Wang, Z., Taylor, K.D.A. and Yan, X. 1992. Studies on the protease activities in Norway lobster (*Nephrops norvegicus*) and their role in the phenolase activation process. *Food Chem.* 45: 111-116.
- Wang, Z., Taylor, K.D.A. and Yan, X. 1994. Further studies on the roles of proteases in the activation of phenolase from Norway lobster (*Nephrops norvegicus*). *Food Chem.* 51: 99-103.
- Wasson, D.H. 1992. Fish muscle proteases and the heat-induced myofibrillar degradation: A Review. *J. Aquat. Food Prod. Technol.* 1(2): 23-41.
- Wei, C.I., Fernando, S.Y. and Huang, T.S. 1991. Mutagenicity studies of kojic acid. Proceedings of the Fifteenth annual conference Tropical and Subtropical Fisheries Technological Conference of the Americas, p. 464-470. Orlando, FL.

- Whitaker, J.R. 1972. Polyphenol oxidase. *In* Principle of enzymology for the food sciences. (Whitaker, J.R. ed), pp. 571-582. Marcel Dekker: New York.
- Whitaker, J.R. 1994. Principles of Enzymology for the Food Sciences. Marcel Dekker: New York.
- Williams, A.B. 1984. Shrimps, lobsters and crabs of the Atlantic coast of the eastern United States, Maine to Florida. Smithsonian Institution Press. Washington, D.C. 550 pp.
- Williams, H.G., Davidson, G.W. Mamo, J.C. 2003. Heat-induced activation of polyphenoloxidase in western rock lobster (*Panulirus Cygnus*) hemolymph: implications for heat processing. *J. Food Sci.* 68: 1928-1932.
- Wolfe, S.K. 1980. Use of CO and CO<sub>2</sub> enriched atmospheres for meats, fish and produce. *Food Technol.* 34: 55-58.
- Xiong, Y.L. 2005. Role of myofibrillar proteins in water-binding in brine-enhanced meats. *Food Res. Int.* 38: 281-287.
- Yamagata, M. and Low, L.K. 1995. Banana shrimp, *Penaeus merguensis*, quality changes during iced and frozen storage. *J. Food Sci.* 60: 721-726.
- Yamamoto, A. 1975. *In* Enzyems in food processing. (Reed, G. ed.), pp 123. Academic Press: New York.
- Yamashita, M. and Konagaya, S. 1990a. High activities of cathepsins B, D, H and L in the white muscle of chum salmon in spawning migration. *Comp. Biochem. Physiol.* 95B: 149-152.
- Yamashita, M. and Konagaya, S. 1990b. Purification and characterization of cathepsin B from ordinary muscle of cod- *rhynchus keta*. *Comp. Biochem. Physiol.* 96B: 247-252.
- Yamashita, M. and Konagaya, S. 1991. Hydrolytic action of salmon cathepsin B and L to muscle softening. *Nippon Suisan Gakkaishi.* 57: 1917-1922.
- Yu, G-L., Wang, Y-H., Liu, S-Q. and Tian, X-L. 1996. Effects of low molecular weight chitosan (LMC-1) on shrimp preservation. *Chin. J. Oceanol. Limnol.* 14: 189-192.
- Zaika, L.L., Cooke, P. and Dobson, T. 1991. Effect of sodium polyphosphates on ultrastructure of *Listeria monocytogenes*. Presented at 91<sup>st</sup> general meeting of the American Society for Microbiology, Dallas, TX. pp. 5-9.

Zeng, Q.Z., Thorarinsdottir, K.A. and Olafsdottir, G. 2005. Quality changes of shrimp (*Pandalus borealis*) stored under different cooling conditions. J. Food Sci. 70: 459-466.