

Contents

	Page
Contents	(10)
List of Tables	(13)
List of Figures	(16)
Chapter	
1. Introduction	1
Literature Review	3
1. Sarcoplasmic protein	4
2. Stroma protein	4
3. Myofibrillar protein	4
4. Surimi and surimi production	8
5. Gelation of muscle proteins	11
6. Role of endogenous transglutaminase in setting and protein cross-linking	16
7. Improvement of gel quality of surimi	19
8. Phosphate compounds	25
Objectives	32
2. Materials and methods	33
1. Materials	33
2. Instruments	34
3. Study on the effect of phosphate compounds on setting and gel forming ability of surimi	35

Contents (Continued)

	Page
4. Study on the effect of phosphate compounds and calcium chloride or microbial transglutaminase on setting and gel forming ability of surimi	37
5. Study on the effect of phosphate on transglutaminase activity from fish muscle	38
6. Study on the effect of phosphate on fish muscle protein and gel forming ability of surimi	40
7. Statistical analysis	42
3. Results and Discussion	43
1. Effect of phosphate compounds on setting and gel forming ability of surimi	43
2. Effect of sodium pyrophosphate in combination with calcium chloride on surimi gel properties	59
3. Effect of setting time on gel forming ability of surimi gel added with sodium pyrophosphate in combination with calcium chloride	66
4. Effect of sodium pyrophosphate in combination with microbial transglutaminase on surimi gel properties	72
5. Effect of setting time on gel forming ability of surimi gel added with sodium pyrophosphate in combination with MTGase	78
6. Effect of sodium pyrophosphate and/or calcium chloride and/or microbial transglutaminase on microstructure of surimi gel	83
7. Effect of phosphate compounds on gel forming ability of surimi	86
8. Effect of phosphate compounds on fish muscle protein	94

Contents (Continued)

	Page
4. Conclusion	101
Suggestion	102
References	103
Appendix	123
Vitae	131

List of Tables

Table	Page
1. Conformational changes which may occur during the thermal denaturation of natural actomyosin	12
2. Classes, formulas, pH, solubility, and functions of several phosphates	26
3. Expressible moisture content of surimi gels from bigeye snapper added with various phosphate compounds at different concentrations and heated under various conditions	50
4. Expressible moisture content of surimi gels from threadfin bream added with various phosphate compounds at different concentrations and heated under various conditions	51
5. Whiteness of surimi gels from bigeye snapper added with various phosphate compounds at different concentrations and heated under various conditions	52
6. Whiteness of surimi gels from threadfin bream added with various phosphate compounds at different concentrations and heated under various conditions	52
7. Solubility of bigeye snapper surimi gels added with various phosphate compounds at different concentrations and heated under various conditions	54
8. Solubility of threadfin bream surimi gels added with various phosphate compounds at different concentrations and heated under various conditions	55
9. Whiteness of kamaboko gels from bigeye snapper and threadfin bream surimi added with sodium pyrophosphate and calcium chloride at various concentrations	62

List of Tables (Continued)

Table	Page
10. Solubility of kamaboko gels from bigeye snapper and threadfin bream surimi added with sodium pyrophosphate and calcium chloride at different concentrations	63
11. Expressible moisture content of bigeye snapper and threadfin bream kamaboko gel added with 0.025% PP and 50 mmol CaCl ₂ /kg and set for various times at 40°C	68
12. Whiteness of kamaboko gels from bigeye snapper and threadfin bream surimi added with 0.025% PP and 50 mmole CaCl ₂ /kg and set at 40°C for different times	69
13. Solubility of kamaboko gels from bigeye snapper and threadfin bream surimi added with 0.025%PP and 50 mmole/kg CaCl ₂ and set at 40°C for different times	70
14. Expressible moisture content of kamaboko gels from bigeye snapper and threadfin bream surimi added with PP and MTGase at various levels	74
15. Whiteness of kamaboko gel from bigeye snapper and threadfin bream surimi added with PP and MTGase at various levels	75
16. Solubility of kamaboko gel from bigeye snapper and threadfin bream surimi added with PP and MTGase at various levels	76
17. Expressible moisture content of kamaboko gels from bigeye snapper surimi (added with 0.025% PP and 0.05% MTGase) and from threadfin bream surimi (added with 0.05% PP and 0.05% MTGase) and set at 40°C for different times	80

List of Tables (Continued)

Table	Page
18. Whiteness of kamaboko gels from bigeye snapper (added with 0.025% PP and 0.05% MTGase) and for threadfin bream surimi (added with 0.05% PP and 0.05% MTGase) and set at 40°C for different times	80
19. Solubility of kamaboko gels from bigeye snapper surimi (added with 0.025%PP and 0.05% MTGase) and from threadfin bream surimi (added with 0.05% PP and 0.05% MTGase) and set at 40°C for different times	81
20. Whiteness of kamaboko gels from bigeye snapper and threadfin bream surimi added with PP or TPP at various levels in the presence of 20 mmole EGTA/kg	89
21. Whiteness of directly heated gels from bigeye snapper and threadfin bream surimi added with PP or TPP at various levels in the presence of 20 mmole EGTA/kg	89
22. Solubility of kamaboko gels from bigeye snapper and threadfin bream surimi added with PP or TPP at various levels in the presence of 20 mmole EGTA/kg	90
23. Solubility of directly heated gels from bigeye snapper and threadfin bream surimi added with PP or TPP at various levels in the presence of 20 mmole EGTA/kg	91
24. Inactivation rate constant of natural actomyosin Ca^{2+} -ATPase of bigeye snapper and threadfin bream at various temperatures	97
25. Inactivation rate constant of natural actomyosin Mg^{2+} -ATPase of bigeye snapper and threadfin bream at various temperatures	97

List of Figures

Figure	Page
1. Structure of myosin	6
2. a) Proteins of the thin filament	7
b) Arrangement of actin, troponin and tropomyosin	
3. Formation of a gel network structure	13
4. A Schematic representation of the thermal aggregation of fish myosin	16
5. Reaction catalyzed by TGase	17
6. Scheme for surimi preparation	34
7. Scheme for surimi gel preparation	35
8. Breaking force and deformation of kamaboko, directly heated and suwari gels from bigeye snapper surimi added with phosphate compounds at different levels	47
9. Breaking force and deformation of kamaboko, directly heated and suwari gels from threadfin bream surimi added with phosphate compounds at different levels	48
10. SDS-PAGE patterns of protein in kamaboko, directly heated and suwari gels from bigeye snapper surimi added with various phosphate compounds at different levels (%)	57
11. SDS-PAGE patterns of protein in kamaboko, directly heated and suwari gels from threadfin bream surimi added with various phosphate compounds at different levels (%).	58
12. Breaking force and deformation of kamaboko gels from bigeye snapper and threadfin bream surimi added with sodium pyrophosphate and calcium chloride at different levels	60

List of Figures (Continued)

Figure	Page
13. Expressible of kamaboko gel from bigeye snapper and threadfin bream surimi added with sodium pyrophosphate and calcium chloride at different levels	61
14. SDS-PAGE pattern of protein in kamaboko gel from bigeye snapper and threadfin bream surimi added with sodium pyrophosphate (PP) and calcium chloride at different levels	65
15. Breaking force and deformation of gels from bigeye snapper and threadfin bream surimi added with 0.025% PP and 50 mmole CaCl ₂ /kg and set at 40°C for different times, followed by heating at 90°C 20 min	67
16. SDS-PAGE pattern of protein in kamaboko gels from bigeye snapper and threadfin bream surimi added with 0.025% sodium pyrophosphate and 50 mmole/kg calcium chloride and set at 40°C for different times	71
17. Breaking force and deformation of kamaboko gels from bigeye snapper and threadfin bream (B) surimi added with sodium pyrophosphate and microbial transglutaminase at different level	73
18. SDS-PAGE patterns of protein in kamaboko gels from bigeye snapper and threadfin bream surimi added with sodium pyrophosphate (PP) and microbial transglutaminase at different levels	77
19. Breaking force and deformation of kamaboko gels from bigeye snapper surimi added with 0.025% sodium pyrophosphate and 0.05% microbial transglutaminase and set at 40°C for different times	79

List of Figures (Continued)

Figure	Page
20. Breaking force and deformation of kamaboko gels from threadfin bream surimi added with 0.05% PP and 0.05% MTGase and set at 40°C for different times	79
21. SDS-PAGE patterns of proteins in kamaboko gel from bigeye snapper surimi (added with 0.025% PP and 0.05% MTGase) and from threadfin bream surimi (added with 0.025% PP and 0.05% MTGase) and set at 40°C for different times	82
22. Microstructure of kamaboko gels from bigeye snapper surimi without additive, with 0.025% PP, with 0.025% PP and 50 mmole CaCl ₂ /kg, and with 0.025% PP and 0.05% MTGase	84
23. Microstructure of kamaboko gels from threadfin bream surimi without additive, with 0.025%PP, with 0.025%PP and 50 mmole CaCl ₂ /kg, and with 0.05% PP and 0.05% MTGase	85
24. Breaking force and deformation of kamaboko gels and directly heated gel from bigeye snapper surimi added with PP or TPP at different levels in the presence of 20 mmole EGTA/kg	87
25. Breaking force and deformation of kamaboko gels and directly heated gel from threadfin bream surimi added with PP or TPP at different levels in the presence of 20 mmole EGTA/kg	88
26. SDS-PAGE patterns of proteins in kamaboko gels from bigeye snapper and threadfin bream surimi added with PP or TPP at different level in the presence of 20 mmole EGTA/kg	92

List of Figures (Continued)

Figure	Page
27. SDS-PAGE patterns of proteins in directly heated gels from bigeye snapper and threadfin bream surimi added with PP or TPP at different level in the presence of 20 mmole EGTA/kg	93
28. ATPase activity of NAM from bigeye snapper and threadfin bream treated with various phosphates at different concentrations. Activities were determined after dialysis of treated NAM against 0.6 M KCl	95
29. TGase activity of fish muscle from bigeye snapper and threadfin bream in the presence of salt or salt in combination with phosphates at different levels	99
30. TGase activity of fish muscle from bigeye snapper and threadfin bream in the presence of salt or salt in combination with CaCl ₂ or PP at different levels	100