

REFERENCES

- Al-Janabi, J., Hartsuck, J., and Tang, J. 1972. Kinetics and mechanism of pepsinogen activation. *J. Biol Chem.* 247: 4628-4632.
- An, H., Seymour, T. A., Wu, J. and Morrissey, M. T. 1994. Assay systems and characterization of Pacific whiting (*Merluccius productus*) protease. *J. Food Sci.* 59: 277-281.
- AOAC. 1999. Official Method of Analysis. 16th ed. Association of Official Chemists. Washington, D. C.
- Arunchalam, K. and Haard, N.F. 1985. Isolation and characterization of pepsin from polar cod (*Boreogadus saida*). *Comp. Biochem. Physiol.* 80B: 467-473.
- Ash, M. and Ash, I. 1997. Handbook of Food Additives. Gower Publishing. Brookfield.
- Babu, I. R. and Ganesh, K. N. 2001. Enhanced triple helix stability of collagen peptides with 4R-Aminoprolyl (Amp) residues: relative roles of electrostatic and hydrogen bonding effects. *J. Am. Chem. Soc.* 123: 2079-2080.
- Bateman, J. F., Lamandé, S. R. and Ramshaw, J. A. M. 1996. Collagen superfamily. In Extracellular metrix. Molecular components and interactions. (Comper, W. D., ed). p. 22-27. Harwood Academic Publishers. UK.
- Battaglia, R. 2003. Japan Fishery Products Annual 2003. GAIN Report #FR2041. USDA Foreign Agricultural Service.
- Baudys, M., Erdene, T. G., Kostka, V., Pavlik, M. and Folmann, B. 1988. Comparison between prochymosin and pepsinogen from lamb and calf. *Comp. Biochem. Physiol.* 89B: 385-391.
- Benjakul, S., Visessanguan, W. and Tanaka, M. 2003a. Partial purification and characterization of trimethylamine-N-oxide demethylase from lizardfish kidney. *Comp. Biochem. Physiol.* 135B: 359-371.
- Benjakul, S., Visessanguan, W. and Leelapongwattana, K. 2003b. Purification and characterization of heat-stable alkaline proteinase from bigeye snapper (*Priacanthus macracanthus*) muscle. *Comp. Biochem. Physiol.* 134B: 579-591.

- Benjakul, S., Chantarasuwan, C., and Visessaneuan, W. 2003c. Effect of medium temperature setting on gelling characteristics of surimi from some tropical fish. *Food Chem.* 82: 567-574.
- Bergman, I. and Loxley, R. 1963. Two improved and simplified methods for the spectrophotometric determination of hydroxyproline. *Anal. Chem.* 35: 1961-1965.
- Bigi, A., Bracci, B., Cojazzi, G., Panzamolta, S. and Roveri, N. 1998. Drawn gelatin films with improved mechanical properties. *Biomaterials.* 19: 2335-2340.
- Bloom, O.T. 1925. Machine for testing jelly strength of glue, gelatins, and the like. US Patent #1,540,979. Assigned to Swift.
- Bohak, Z. 1973. The kinetics of the conversion of chicken pepsinogen to chicken pepsin. *Eur. J. Biochem.* 32: 547-554.
- Bornstein, P., Kang, A. H. and Piez, K. A. 1966. The limited cleavage of native collagen with chymotrypsin, trypsin and cyanogens bromide. *Biochem.* 5: 3803.
- Brewer, P., Helbig, N. and Haard, N. F. 1984. Atlantic cod pepsin – characterization and use as a rennet substitute. *Can. Inst. Food Sci. Technol. J.* 17: 038-043.
- Budavari, S. 1996. *Merck Index*, (12th ed.) Whitehouse Station, NJ: Merck.
- Burghagen, M. 1999. Collagen. In *Food Chemistry*. 2nd ed. (Belitz, H. D. and Grosch, W., eds.). p. 540-547. Springer-verlag Berlin Heidelberg. Berlin.
- Campos, L. A. and Sancho, J. 2003. The active site of pepsin is formed in the intermediate conformation dominant at mildly acidic pH. *FEBS Letters.* 538: 89-95.
- Castillo-Yanez, F. J., Pacheco-Aquilar, R., Garcia-Carreno, F. L. and Navarrete del Toro, M. A. 2004. Characterization of acidic proteolytic enzymes from Monterey sardine (*Sardinops sagax caerulea*) viscera. *Food Chem.* 85: 343-350.
- Cheow, C. S., Norizah, M. S., Kyaw, Z. Y. and Howell, N. K. 2007. Preparation and characterisation of gelatins from the skins of sin croaker (*Johnius dussumieri*) and shortfin scad (*Decapterus macrosoma*). *Food Chem.* 101: 386-391.
- Cho, S. M., Kwak, K. S., Park, D. C., Gu, Y. S., Ji, C. I., Jang, D. H., Lee, Y. B. and Kim, S. B. 2004. Processing optimization and functional properties of gelatin from shark (*Isurus oxyrinchus*) cartilage. *Food Hydrocol.* 18: 573-579.

- Chobert, J., Sitohy, M. Z., and Whitaker, J. R. 1988. Solubility and emulsifying properties of caseins modified enzymatically by *Staphylococcus aureus* V8 protease. *J. Agric. Food Chem.* 36: 220–224.
- Choi, S. S. and Regenstein, J. M. 2000. Physiochemical and sensory characteristics of fish gelatin. *J. Food Sci.* 65: 194-199.
- Chomarat, N., Robert, L., Seris, J. L. and Kern, P. 1994. Comparative efficiency of pepsin and proctase for the preparation of bovine skin gelatin. *Enzyme Microb. Technol.* 16: 756-760.
- Courts, A. 1955. The N-terminal amino acid residues of gelatin. 3. Enzymic degradation. *Biochem J.* 59: 382–386.
- Department of export promotion. 2005. Summary of USA seafood trade in 2004. Ministry of Commerce, Thailand. (<http://www.depthai.go.th>).
- Diaz-Lopez, M., Moyano-Lopez, F. J., Alarcon-Lopez, F. J., Garcia-Carreno, F. L., and Navarrete del Toro, M. A. 1998. Characterization of fish acid proteases by substrate- gel electrophoresis. *Comp. Biochem. Physiol.* 121B. 369-377.
- Drake, M. P., Davison, P. F., Rump, S. and Schmitt, F. O. 1966. Action of proteolytic enzyme on tropocollagen and insoluble collagen. *Biochem.* 5: 301-312.
- Eastoe, J. E. and Leach, A. A. 1977. Chemical constitution of gelatin. In The science and technology of gelatin. (Ward, A. G. and Courts, A, eds). p. 73-107. Academic Press. London.
- Eckhoff, K. M., Aidos, I., Hemre, G. I. and Lie, O. 1998. Collagen content in farmed Atlantic salmon (*Salmo solar*, L.) and subsequent changes in solubility during storage on ice. *Food Chem.* 62: 197-200.
- England, S. and Seifter, S. 1990. Precipitation techniques. In Methods in enzymology: Guide to protein purification. vol. 182. (Deutscher, M. P., ed.). p. 285-300. Academic Press. USA.
- Fernandez-Diaz, M. D., Montero, P. and Gomez-Guillen, M. C. 2001. Gel properties of collagen from skin of cod (*Gadus morhua*) and hake (*Merluccius merluccius*) and their modification by coenhancers magnesium sulphate, glycerol and transglutaminase. *Food Chem.* 74: 161-167.
- Fersht, A. 1984. Enzyme structure and mechanism. 2nd Ed. W.H. Freeman and Company. USA.

- Foegeding, E.A., Lanier, T. C. and Hulin, H.O. 1996. Characteristic of edible muscle tissues. In Food chemistry. 3rd ed. (Fennema, O. R., ed.). p. 902-906. Marcel Dekker. New York.
- Food Chemicals Codex. 1996. Food and Nutrition Board, National Academy of Sciences. 4th Ed. National Academy Press. Washington, DC.
- Fujimoto, Z., Fujii, Y., Kaneko, S., Kobayashi, H. and Mizuno, H. 2004. Crystal structure of aspartic proteinase from *Irpeus lacteus* in complex with inhibitor pepstatin. J. Mol. Biol. 341: 1227-1235.
- Fusek, M. and Větvička, V. 1995. Aspartic proteinases: Physiology and pathology. CRC Press. New York.
- Galea, C. A., Dalrymple, B. P. and Kuypers, R. 2000. Modification of the substrate specificity of porcine pepsin for the enzymatic production of bovine hide gelatin. Protein Sci. 9: 1947-1959.
- Gass, T. 2001. Petition for amending the National List of the USDA's National Organic Program. USDA/AMS/TM/NOP. Washington, DC.
- Gildberg, A. and Raa, J. 1983. Purification and characterization of pepsins from the Arctic fish capelin (*Mallotus villosus*). Comp. Biochem. Physiol. 75A: 337-342.
- Gildberg, A. 1987. Purification and characterization of cathepsin D from the digestive gland of the pelagic squid *Todarodes sagittatus*. J. Sci. Food Agric. 39: 85-94.
- Gildberg, A. 1988. Aspartic proteinase in fishes and aquatic invertebrates. Comp. Biochem. Physiol. 91B: 425-435.
- Gildberg, A., Olsen, R. L. and Bjarnason, J. B. 1990. Catalytic properties and chemical composition of pepsins from Atlantic cod (*Gadus morhua*). Comp. Biochem. Physiol. 96B: 323-330.
- Gilsenan, P. M. and Ross-Murphy, S. B. 2000. Rheological characterization of gelatins from mammalian and marine sources. Food Hydrocol. 14: 191-196.
- Gomez-Guillen, M. C. and Montero, P. 2001. Extraction of gelatin from mermigrim (*Lepidorhombus boscii*) skins with several organic acids. J. Food Sci. 66: 213-216.
- Gomez-Guillen, M. C., Turnay, J., Fernandez-Diaz, M. D., Ulmo, N., Lizarbe, M. A. and Montero, P. 2002. Structural and physical properties of gelatin extracted from different marine species: a comparative study. Food Hydrocol. 16: 25-34.

- Guerard, F. and Le Gal, Y. 1987. Characterization of a chymosin-like pepsin from dogfish *Scyliorhinus canicula*. Comp. Biochem. Physiol. 88B: 823-827.
- Gudmundsson, M. and Hafsteinsson, H. 1997. Gelatin from cod skins as affected by chemical treatments. J. Food Sci. 62: 37-40.
- Gudmundsson, M. 2002. Rheological properties of fish gelatins. J. Food Sci. 67: 2172-2176.
- Haard, N. F., Helbig, N. and Feltham, L. A. W. 1981. The temperature characteristics of pepsin from two stock of American salmon (*Osmerus mordax*). In Proc. Workshop of the Labrador Coastal Offshore Region. pp. 174-196. Newfoundland Institute of Cold Ocean Science.
- Haard, N. F. 1994. Protein hydrolysis in seafoods. In Seafoods: Chemistry, processing technology and quality. (Shahidi, F. and Botta, J. R., eds.). p. 10-33. Chapman & Hall. UK.
- Haard, N. F. and Simpson, B. K. 2000. Seafood enzymes: Utilization and influence on postharvest seafood quality. Marcel Dekker. New York.
- Haug, I. J., Draget, K. I. and Smidsrød, O. 2004. Physical and rheological properties of fish gelatin compared to mammalian gelatin. Food Hydrocol. 18: 203-213.
- Hwang, J. -H., Mizuta, S., Yogoyama, Y. and Yoshinaka, R. 2007. Purification and characterization of molecular species of collagen in the skin of skate (*Raja kenojei*). Food Chem. 100: 921-925.
- Hudson, C.B. 1994. Gelatine – Relating structure and chemistry to functionality, In Food Hydrocolloids: Structures, Properties, and Functions. (Nishihari, K. and Doi, E., eds). Plenum. New York.
- Igoe, R.S. 1983. Dictionary of Food Ingredients. Van Nostrand Reinhold. New York.
- Ikoma, T., Kobayashi, H., Tanaka, J., Walsh, D. and Mann, S. 2003. Physical properties of type I collagen extracted from fish scales of *Pagrus major* and *Oreochromis niloticus*. Int. J. Biol. Macro. 32: 199-204.
- Intarasirisawat, R., Benjakul, S., Visessaguan, W., Prodpran, T., Tanaka, M. and Howell, N. K. 2006. Autolysis study of bigeye snapper (*Priacanthus macracanthus*) skin and its effect on gelatin. Food Hydrocol. In press.

- Jamilah, B. and Harvinder, K.G. 2002. Properties of gelatin from skins of fish - black tilapia (*Oreochromis mossambicus*) and red tilapia (*Oreochromis nilotica*). Food Chem. 77: 81-84.
- Johnson, T. and Zabik, M. 1981. Egg albumin proteins interactions in an angel food cake system. J. Food Sci. 46: 1231.
- Johnston-Banks, F. A. 1990. Gelatin. In Food Gels. (Harris, P., ed.). p. 233-289. Elsevier Applied Science. London.
- Jongjareonrak, A., Benjakul, S., Visessanguan, W., Nagai, T. and Tanaka, M. 2005a. Isolation and characterization of acid and pepsin-solubilised collagen from brownstrip red snapper (*Lutjanus vitta*) skin. Food Chem. 93: 475-484.
- Jongjareonrak, A., Benjakul, S., Visessanguan, W. and Tanaka, M. 2005b. Isolation and characterization of collagen from bigeye snapper (*Priacanthus macracanthus*) skin. J. Sci. Food Agric. 85: 1203-1210.
- Jongjareonrak, A., Benjakul, S., Visessanguan, W. and Tanaka, M. 2006. Skin gelatin from bigeye snapper and brownstripe red snapper: Chemical compositions and effect of microbial transglutaminase on gel properties. Food Hydrocol. 20: 1216-1222.
- Kim, J. S. and Park, J. W. 2004. Characterization of acid-soluble collagen from Pacific whiting surimi processing by products. J. Food Sci. 69: C637-C642.
- Kinsella, J. E. 1977. Functional properties of protein: possible relationships between structure and function in foams. J. Food Chem. 7: 273-288.
- Kittiphattanabawon, P. 2004. Extraction and characterization of collagen and gelatin from bigeye snapper (*Priacanthus tayenus*) skin and bone. Master of Science Thesis. Prince of Songkhla University. 142 p.
- Kittiphattanabawon, P., Benjakul, S., Visessanguan, W., Nagai, T. and Tanaka, M. 2005. Characterization of acid soluble collagen from the skin and bone of bigeye snapper (*Priacanthus tayenus*). Food Chem. 89: 363-372.
- Klomklao, S., Benjakul, S. and Visessanguan, W. 2004. Comparative studies on proteolytic activity of splenic extract from three tuna species commonly used in Thailand. J. Food Biochem. 28: 355-372.

- Komsa-Penkova, R., Koyonava, R., Kostov, G. and Tenchov, B. 1999. Discrete reduction of type I collagen thermal stability upon oxidation. *Biophys. Chem.* 83: 185-195.
- Kołodziejska, I., Kaczorowski, K., Piotrowska, B. and Sadowska, M. 2004. Modification of properties of gelatin from skins of Baltic cod (*Gadus morhua*) with transglutaminase. *Food Chem.* 86: 203-209.
- Kubota, M. and Ohnuma, A. 1970. Studies on bonito pepsin-II. Enzymic properties of bonito. *Bull. Jap. Soc. Sci. Fish.* 36: 1152-1156.
- Leamml, U. K. 1970. Cleavage of structural proteins during assembly of head of bacteriophage T4. *Nature*. 277: 680-685.
- Lin, L. -H. and Chen, K. -M. 2006. Preparation and surface activity of gelatin derivative surfactants. *Colloids Surf., A: Physicochem. Eng. Aspects*. 272: 8-14.
- Lin, Y. K. and Liu, D. C. 2006. Effects of pepsin digestion at different temperatures and times on properties of telopeptide-poor collagen from bird feet. *Food Chem.* 94: 621-625.
- Liu, H., Li, D. and Guo, S. 2006. Studies on collagen from the skin of channel catfish (*Ictalurus punctatus*). *Food Chem.* In press.
- Leuenberger, B.H. 1991. Investigation of the viscosity and gelation properties of different mammalian and fish gelatins. *Food Hydrocol.* 5: 353-361.
- Love, R. M., Yamaguchi, Y., Creach, Y. And Lavety, J. 1976. The connective tissues and collagens of cod during starvation. *Comp. Biochem. Physiol.* 55B: 487-492.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. And Randall, R. J. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193: 256-275.
- Makinodan, Y., Yokoyama, Y., Kinoshita, M. and Toyohara, H. 1987. Characterizaton of an alkaline proteinase of fish muscle. *Comp. Biochem Physiol.* 87B: 1041-1046.
- Mason, R. W. and Barett, A. J. 1984. Purification and properties of human liver cathepsin L. *Biochem. Soc. Trans.* 12: 11-16.
- Mechanic, G. L., Katz, E. P., Henmi, M., Noyes, C. and Yamauchi, M. 1987. Locus of a histidine-based, stable trifunctional, helix to helix collagen cross-link: Stereospecific collagen structure of type I skin fibrils. *Biochemistry*. 26: 3500-3509.
- McCormick, R. 1987. Exploiting the novel properties of pectin and gelatin gels. *Prepared Foods*. 5: 204-205.

- McCormick, R. J. 1994. Structure and Properties of Tissues. In Muscle Foods; Meat Poultry and Seafood Technology. (Kinsman, D.M., Kotula, A. W. and Breidenstein, B. C., eds.). p. 54-58. Chapman & Hall. New York.
- McWilliams, M. 2001. Foods-Experimental Perspectives. 4th ed. Prentice Hall. Englewood Cliffs.
- Michell, J. R. 1986. Foaming and emulsifying properties of proteins. In Development in food protein. 4th ed. (Hudson, B. J. F., ed.). p. 291-338. Elsevier Applied Science Publishers. London.
- Miller, E. J. 1972. Structural studies on cartilage collagen employing limited cleavage and solubilization with pepsin. Biochem. 11: 4903-4909.
- Mizuta, S., Hwang, J-H. and Yoshinaka, R. 2002. Molecular species of collagen from wing muscle of skate (*Raja Kenojei*). Food Chem. 76: 53-58.
- Mizuta, S., Hwang, J-H. and Yoshinaka, R. 2003. Molecular species of collagen in pectoral fin cartilage of skate (*Raja Kenojei*). Food Chem. 80: 1-7.
- Montero, P., Bonderias, J., Turnay, J. and Leyzarbe, M. A. 1990. Characterization of hake (*Merluccius L.*) and trout (*Salmo irideus Gibb*) collagen. J. Agric. Food Chem. 38: 604-609.
- Montero, P. and Borderias, A. J. 1991. Emulsifying capacity of collagenous material from muscle and skin of hake (*Merluccius merluccius L.*) and trout (*Salmoirideus Gibb*) : Effect of pH and NaCl concentration. Food Chem. 41: 251-267.
- Montero, P., Jimenez-Colmenero, F. and Borderias, J. 1991. Effect of pH and the presence of NaCl on some hydration properties of collagenous material from trout (*Salmo irideus Gibb*) muscle and skin. J. Sci. Food Agric. 54: 137-146.
- Montero, P., Alvarez, C., Marti, M. A. and Borderias, A. J. 1995. Plaice skin collagen extraction and functional properties. J. Food Sci. 60: 1-3.
- Montero, P., Gomez-Guillen, M. C. and Borderias, A. J. 1999. Functional characterization of muscle and skin collagenous material from hake (*Merluccius merluccius L.*). Food Chem. 65: 55-59.
- Montero, P. and Gomez-Guillen, M.C. 2000. Extraction conditions for Megrime (*Lepidorhombus boscii*) skin collagen affect functional properties of resulting gelatin. J. Food Sci. 65: 434-438.

- Morrissey, M. T., Park, J. W. and Huang, L. 2000. Surimi Processing Waste: Its Control and Utiliaztion. In Surimi and Surimi Seafood. (Park, J. W., ed). p.127-166. Marcel Dekker, Inc. New York.
- Moriarty, D. J. W. 1973. The physiology of digestion of blue-green algae in the cichlid fish, *Tilapia nilotica*. J. Zool. Lond. 171: 25-39.
- Muyonga, J. H., Cole, C. G. B. and Duodu, K. G. 2004a. Characterization of acid soluble collagen from skins of young and adult Nile perch (*Lates niloticus*). Food Chem. 85 : 81-89.
- Muyonga, J. H., Cole, C. G. B. and Duodu, K. G. 2004b. Extraction and physico-chemical characterisation of Nile perch (*Lates niloticus*) skin and bone gelatin. Food Hydrocol. 18: 581-592.
- Nagai, T. and Suzuki, N. 2000. Isolation of collagen from fish waste material – skin, bone and fins. Food Chem. 68: 277-281.
- Nagai, T., Worawattanamateekul, W., Suzuki, N. Nakamura, T., Ito, T., Fujiki, K., Nakao, M. and Yano, T. 2000. Isolation and characterization of collagen from rhizostomous jelly fish (*Rhopilema asamushi*). Food Chem. 70: 205-208.
- Nagai, T., Yamashita, E., Taniguchi, K., Kanamori, N. and Suzuki, N. 2001. Isolation and characterization of collagen from the outer skin waste material of cuttlefish (*Sepia lycidas*). Food Chem. 72: 425-429.
- Nagai, T., Araki, Y. and Suzuki, N. 2002. Collagen of the skin of ocellate puffer fish (*Takifugu rubripes*). Food Chem. 78: 173-177.
- Nagai, T. and Suzuki, N. 2002. Preparation and partial characterization of collagen from paper nautilus (*Argonauta argo*, Linnaeus) outer skin. Food Chem. 76: 149-153.
- Nishimoto, M., Mizuta, S. and Yoshinaka, R. 2004. Characterization of molecular species of collagen in muscles of Japanese amberjack, *Seriola quinqueradiata*. Food Chem. 84 : 127-132.
- Noda, M and Murakami, K. 1981. Studies on proteinases from the digestive organs of sardine. Purification and characterization of two acid proteinases from the stomach. Biochim. Biophys. Acta. 658: 27-34.

- Nomura, Y., Sakai, H., Ishii, Y. and Shirai, K. 1996. Preparation and some properties of type I collagen from fish scales. *Biosci. Biotechnol. Biochem.* 60: 2092-2094.
- Norland, R. E. 1990. Fish gelatin. In *Advance in Fisheries Technology and Biotechnology for Increased Profitability*. (Voight, M. N. and Botta, J. K., eds.). p. 325-333. Techcomic Publishing. Lancaster.
- Norris, E. R. and Mathies, J. C. 1953. Preparation, properties, and crystallization of tuna pepsin. *J. Biol. Chem.* 204: 673-680.
- Ockerman, H. W. and Hansen, C. L. 1988. Glue and gelatin. In *Animal by-product processing*. p. 133-157. Ellis Horwood. New York.
- Ogawa, M., Moody, M. W., Portier, R. J., Bell, J., Schexnayder, M. A. and Losso, J. N. 2003. Biochemical properties of black drum and sheepshead seabream skin collagen. *J. Agric. Food Chem.* 51: 8088-8092.
- Ogawa, M., Portier, R. J., Moody, M. W., Bell, J., Schexnayder, M. A. and Losso, J. N. 2004. Biochemical properties of bone and scale collagens isolated from the subtropical fish black drum (*Pogonias cromis*) and sheepshead seabream (*Archosargus probatocephalus*). *Food Chem.* 88: 495-501
- Okuzaki, H. 2001. General theory of gel preparation. In *Gel Handbook*. vol. I. (Osada, Y. and Kajiwara, K., eds.). p. 112-116. Academic Press. New York.
- OMRI. 2002. Gelatin processing. In *National Organic Standards Board Technical Advisory Panel Review*. Complied by Organic Materials Review Institute for the USDA National Organic Program. 1 March 2002. 25p.
- Owen, T. G. and Wiggs, A. J. 1971. Thermal compensation in the stomach of brook trout (*Salvelinus fontinalis* Mitchell). *Comp. Biochem. Physiol.* 40B: 465-473.
- Pearce, K. N., and Kinsella, J. 1978. Emulsifying properties of protein: evaluation of a turbidimetric technique. *J. Agric. Food Chem.* 26: 719-723.
- PB Leiner. 2004. Gelatin Application (Online). Available : <http://www.gelatin.com> [2004, September 1]
- Poppe, J. 1997. Gelatin, In *Thickening and Gelling Agents for Food*. 2nd ed. (Imeson, A., ed.). p. 144-168. Blackie Academic and Professional. London.

- Raa, J. 1990. Biotechnology in aquaculture and the fish processing industry. A success story in Norway. In *Advances in fisheries technology and biotechnology for increased profitability.* (Voigt, M. N. and Botta, J. R., eds.). p. 509-524. Technomic. Lancaster.
- Rajagopalan, T. G., Moore, S. and Stein, W. H. 1966. Pepsin from pepsinogen; preparation and properties. *J. Biol. Chem.* 241: 4940-4950.
- Regenstein, J. M. and Regenstein, C. E. 1991. Current issues in kosher foods. *Trends Food Sci. Technol.* 2: 50-54.
- Rigby, B. J. 1968. Amino-acid composition and thermal stability of the skin collagen of the Antarctic ice-fish. *Nature.* 219: 166-167.
- Robinson, H.W. and Hodgen, C.G. 1940. The biuret reaction in the determination of serum protein. I. A study of the condition necessary for the production of the stable color which bears a quantitative relationship to the protein concentration. *J. Biol. Chem.* 135: 707-725.
- Rochdi, A., Foucat, L. and Renou, J. 2000. NMR and DSC studies during thermal denaturation of collagen. *Food Chem.* 69: 295-299.
- Sadowska, M., Kolodziejska, I. and Niecikowska, C. 2003. Isolation of collagen from the skins of Baltic cod (*Gadus morhua*). *Food Chem.* 81: 257-262.
- Sanchez-Chaing L., Cisternas E. and Ponce O. 1987. Partial purification of pepsins from adult and juvenile salmon fish *Oncorhynchus keta*. Effect of NaCl on proteolytic activities. *Comp. Biochem. Physiol.* 87B: 793-797.
- Segtnan, V. H., Kvaal, K., Rukke, E. O., Schüller, R. B. and Isaksson, T. 2003. Rapid assessment of physico-chemical properties of gelatine using near infrared spectroscopy. *Food Hydrocol.* 17: 585-592.
- Slade, L. and Levine, H. 1987. Polymer-chemical properties of gelatin in foods. In *Advances in meat research.* vol. 4. (Pearson, A. M., Dutsom, T. R. and Bailey, A. J., eds). p. 251-266. Van Nostrand Reinhold Company. New York.
- Scopes, R. K. 1978. Techniques for protein purification. In *Techniques in the life sciences: Techniques in protein and enzyme biochemistry-Part I.* vol. B1/I. (Kornberg, H. L., J. C. Metcalfe, J. C., Northcote, D. H., Pogson, C. I. and Tipton, K. F., eds). p. B101: 7-18. Elsevier/North-Holland Scientific Publishers. Netherlands.

- Sikorski, Z. E., KolaKoska, A. and Pan, B. S. 1990. The nutritive composition of the major groups of marine food organisms. In *Seafood : resources, nutritional composition and preservation.* (Sikorski, Z. E., ed.). p. 29-54. CRC Press. Florida.
- Silva, E. A., Neitzel, I. and Silva, L. H. M. 2001. Gelatin drying process. *Braz. J. Chem. Eng.* 18: 467-478.
- Simon, A., Vandajon, L., Levesque, G. and Bourseau, P. 2002. Concentration and desalination of fish gelatin by ultrafiltration and continuous diafiltration processes. *Desalination.* 144: 313-318.
- Squires, E. J., Haard, N. F. and Feltham, L. A. W. 1986. Gastric proteases of the Greenland cod *Gadus ogac.* I. Isolation and kinetic properties. *Biochem. Cell. Biol.* 64: 205-214.
- Steel, R. G. D. and Torrie, J. H. 1980. *Principles and procedures of statistics: A biometrical approach.* McGraw-Hill, New York.
- Tanji, M., Kageyama, T. and Takahashi, K. 1988. Tuna pepsinogens and pepsins. Purification, characterization and amino-terminal sequences. *Eur. J. Biochem.* 177: 251-259.
- Tello-Solis, S. R. and Romero-Garcia, B. R. 2001. Thermal denaturation of porcine pepsin: a study by circular dichroism. *Int. J. Biol. Macromol.* 8: 129-133.
- The Customs Department. 2005. Import statistics of gelatin 2004. Ministry of finance, Thailand. (<http://www.customs.go.th>).
- Twining, S. S., Alexander, P. A., Huibregtse, K. and Glick, D. M. 1983. A pepsinogen from rainbow trout. *Comp. Biochem. Physiol.* 75B: 109-112.
- Umezawa, H. 1976. Structures and activities of protease inhibitors of microbial origin. *Methods Enzymol.* 45: 678-695.
- US FDA. 1997. FDA Center of Biologics Evaluation and Research, Transmissible Spongiform Encephalopathies Advisory Committees. Transcript of Meeting April 23.
- Veis, A. 1964. *The Macromolecular Chemistry of Gelatin.* Academic Press. New York.
- Vojdani, F. 1996. Solubility. In *Methods of Testing Protein Functionality.* 1st ed. (Hall, G. M., ed.). p. 11-60. St. Edmundsbury Press. Great Britain.
- Ward, A. G. and Courts, A. 1977. *The science and technology of gelatin.* Academic Press. London.

- Whitaker, J. R. 1994. Principles of enzymology for the food sciences. 2nd Ed. Marcel Dekker. New York.
- Wilk, S. and Orlowski, M. 1983. Evidence pituitary cation-sensitive neutral endopeptidase a multicatalytic protease complex. *J. Neurochem.* 40: 842-849.
- Wong, D. W. S. 1989. Mechanism and Theory in Food Chemistry. Van Nostrand Reinhold. New York.
- Xiong, Y. L. 1997. Structure-function relationships of muscle proteins. In *Food Proteins and their applications*. (Damodaran, S. and Paraf, A., eds.). p. 341392. Marcel Dekker, Inc. New York.
- Xu, Y., Bhate, M. and Brodsky, B. 2002. Characterization of the nucleation step and folding of a collagen triple-helix peptide. *Biochem.* 41: 8143-8151.
- Yata, M., Yoshida, C., Fujisawa, S., Mizuta, S. and Yoshinaka, R. 2001. Identification and characterization of molecular species of collagen in fish skin. *J. Food Sci.* 66: 247-251.
- Zayas, J. F. 1997. *Functionality of Proteins in Food*. Springer. Berlin.
- Zhou, P. and Regenstein, J. M. 2003. Extraction of gelatin from Pollock skin. Aquatic food products. 2003 IFT Annual Meeting. Chicago.
- Zhou, P., Mulvaney, S. J. and Regenstein, J. M. 2006. Properties of Alaska pollock skin gelatin: a comparison with tilapia and pork skin gelatins. *J. Food Sci.* 71: C313-C321.