

References

- Ajandouz, E. H., Tchiakpe, L. S., Ore, F. D., Benajiba, A. and Puigserver, A. 2001. Effects of pH on caramelization and Maillard reaction kinetics in fructose-lysine model systems. *J. Food Sci.* 66: 926-931.
- Al-Abdessalaam, T. Z. 1995. Marine species of the Sultanate of Oman. Oman: Marine Science and Fisheries Center, Ministry of Agriculture and Fisheries.
- Amano, K., 1962. The influence on nutritive value of fish with special reference to fermented fish products of South East Asia. In *Fish in Nutrition*, Proceeding of the FAO Congress. (Heen, E. and Kreuzer, R., eds.). p. 180-197. Fishing News Books. Washington DC
- An, H., Seymour, T. A. Wu, J. and Morrissey, M. T. 1994. Assay systems and characterization of Pacific whiting (*Merluccius productus*) protease. *J. Food Sci.* 59: 277-281.
- An, H. and Visessanguan, W. 2000. Recovery of enzymes from seafood processing wastes. In *Seafood Enzymes, Utilization and Influence on Postharvest Seafood Quality*. (Haard, N.F. and Simpson B. K., eds.) p. 641-664. Marcel Dekker. New York.
- Ando, M., Yoshinoto, Y., Inabu, K., Nakagawa, T. and Makinodan, Y., 1995. Post-mortem changes of three-dimensional structure of collagen fibrillar network in fish muscle pericellular connective tissues corresponding to post-mortem tenderization. *Fish. Sci.* 61: 327-330.
- AOAC., 2000. Official Method of Analysis. 16th ed. Association of Official Analytical Chemists, Washington, DC, USA.

- Aquerreta, Y., Astiasaran, I. and Bello, J. 2001. Use of exogenous enzymes to elaborate the Roman fish sauce “garum”. *J. Sci. Food Agric.* 82: 107-112.
- Aranishi, F., Ogata, H., Hara, K., Osatomi, K. and Ishihara, T. 1997. Purification and characterization of cathepsin L from hepatopancreas of carp (*Cyprinus carpio*). *Comp. Biochem. Physiol.* 118B: 531-537.
- Arason S. 1994. Production of fish silage. In *Fisheries processing, biotechnological applications* (Martin A. M., ed.). p. 245–72. Chapman & Hall. London.
- Areekijseree, M., Engkagul, A., Kovitvadhi, U., Thongpan, A., Mingmuang, M., Pakkong, P. and Rungruangsak-Torriksen, K. 2004. Temperature and pH characteristics of amylase and proteinase of adult fresh water pearl mussel, *Hyriopsis (Hyriopsis) bialatus* Simpson 1900. *Aquaculture* 234: 575-587.
- Argento-Ceru, M. P., and Autuori, F. 1985. In *Structure and function of amine oxidases* (Mondovi B., ed.). p. 89-104. CRC Press, Boca Raton.
- Arnold, S. H., Price, R. J. and Brown, W. D. 1980. Histamine formation by bacteria isolated from skipjack tuna (*Katsuwonus pelamis*). *Bull. Jpn. Soc. Sci. Fish.* 46: 991-995.
- Arunchalam, K. and Haard, N. F. 1985. Isolation and characterization of pepsin isoenzymes from polar cod (*Boreogadus saida*). *Comp. Biochem. Physiol.* 80B: 467-473.
- Asgeirsson, B. and Bjarnason, J. B. 1991. Structural and kinetic properties of chymotrypsin from Atlantic cod (*Gadus morhua*) comparison with bovine trypsin. *Comp. Biochem. Physiol.* 99B: 327-335.

- Asgeirsson, B., Fox, J.W. and Bjarnason, J.B. 1989. Purification and characterization of trypsin from the poikilotherm (*Gadus Morhua*). *Eur. J. Biochem.* 180: 85-94.
- Ashie, I. N. A., Smith, J. P. and Simpson, B. K. 1996. Spoilage and shelf-life extension of fresh fish and shellfish. *Crit. Rev. Food Sci.* 36: 87-121.
- Ashoor, S. H. and Zent, J. B. 1984. Maillard browning of common amino acids and sugars. *J. Food Sci.* 49: 1206-1207.
- Baisier, W. M. and Labuza, T. P. 1992. Maillard browning kinetics in liquid model system. *J. Agric. Food Chem.* 4: 707-713.
- Balakrishnan, M., Thirupathi, V. and Devadas, C. T. 2001. Role of Food Additives in Food Processing-An overview. Beverage and Food World. Connecticut, USA.
- Bardócz, S. 1995. Polyamines in food and their consequences for food quality and human health. *J. Food. Sci. Technol.* 6: 341-346.
- Bates, L., Ames, J. M., Mac Dougall, D. B. and Taylor, P. C. 1998. Laboratory reaction cell to model Maillard colour development in a starch-glucose-lysine system. *J. Food Sci.* 6: 991-996.
- Beddows, C. G. 1985. Fermented fish products. In *Microbiology of fermented food* vol.2. (Wood, B. J. B., ed.). p. 2-23. Elsevier Applied Science Publishers. London.
- Beddows, C.G. and Ardestir, A.G. 1979a. The production of fish protein soluble for use in fish sauce manufacture. The use of added enzymes. *J. Food Technol.* 14: 603-613.

- Beddows, C.G. and Ardeshir, A.G. 1979b. The production of fish protein soluble for use in fish sauce manufacture. The use of acid at ambient temperature. *J. Food Technol.* 14: 613-623.
- Beddows, C. G., Ardeshir, A. G. and Daud, W. J. 1979. Biochemical changes during the manufacture of budu. *J. Sci. Food Agric.* 30: 1097-1103.
- Beddows, C. G., Ardeshir, A. G. and Daud, W. J. 1980. Development and origin of the volatile fatty acids in budu. *J. Sci. Food Agric.* 31: 86-92.
- Ben-Gigirey, B., Vieites Baptista de Sousa, J. M., Villa, T. G. and Barros-Velaquez, J. 1998. Changes in biogenic amines and microbiological analysis in albacore (*Thunnus alalunga*) muscle during frozen storage. *J. Food Prot.* 61: 608-615.
- Benjakul, S., Lerttikkul, W. And Bauer, F. 2005a. Antioxidant activity of Maillard reaction products from porcine plasma protein-sugar model system. *Food Chem.* 93: 189-196.
- Benjakul, S., Seymour, T. A., Morrissey, M. T. and An, H. 1997. Physicochemical changes in Pacific whiting muscle proteins during ice storage. *J. Food Sci.* 62: 729-733.
- Benjakul, S., Visessanguan, W., Phongkanpai, V. and Tanaka, M. 2005b. Antioxidative activity of caramelisation products and their preventive effect on lipid oxidation in fish mince. *Food Chem.* 90: 231-239.
- Benjakul, S., Visessangun, W., Riebroy, S., Ishizaki, S. and Tanaka, M. 2002. Gel-forming properties of surimi produced from bigeye snapper, (*Priacanthus tayenus*) and (*P. marlatus*) stored in ice. *J. Sci. Food Agric.* 82: 1442-1451.

- Bersamin, S.V. and Napugan, R. S. J. 1961. Preliminary studies on the comparative chemical composition of the different commercial brands of Patis in the Philippines. *J. Fish.* 151-157.
- Brillantes, S., Paknoi, S. and Totakien, A. 2002. Histamine formation in fish sauce production. *J. Food Sci.* 67: 2090-2094.
- Calligaris, S., Manzocco, L., Anese, M. and Nicoli, M. C. 2004. Effect of heat-treatment on the antioxidant and proxidant activity of milk. *Intern. Dairy J.* 14: 421-427.
- Capillas, C. R. and Moral, A. 2001. Correlation between biochemical and sensory quality indices in hake stored in ice. *34:* 441-447.
- Castillo-Yañez, F. J., Pacheco-Aguilar, R. P., Garcia-Carreño, F. L. and Navarrete-Deltaro, M. A. 2004. Characterization of acid proteolytic enzymes from Monterey sardine (*Sardinops sagaxcaerulea*) viscera. *Food Chem.* 85: 343-350.
- Chaiyanan, S., Sarayanit, S., Chamaoot, M. and Kerornmala, C. 1988. Fish sauce fermentation by using microbial inoculation and recycling system. In *Food science and technology in industrial development* vol. I. p. 320-325. Proceeding of the food conference (24-26 October 1988). Bangkok, Thailand.
- Chaveesuk, R., Smith, J. P. and Simpson, B. K. 1993. Production of fish sauce and acceleration of sauce fermentation using proteolytic enzymes. *J. Aquatic Food Prod. Technol.* 2(3): 59-77.

- Chayavan, S., Rao, R. M., Liuzzo, J. A. and Khan, A. 1983. Chemical characterization and sensory evaluation of dietary sodium potassium fish sauce. *J. Agric. Food Chem.* 31: 859-863.
- Chin, K. D. L., Koehler, P. E. 1983. Identification and estimation of histamine, tryptamine, phenylamine, and tyramine in soy sauce by thin layer chromatography and dansyl constituents. *J. Food Sci.* 48: 1826-1828.
- Cho, S. M, Gu, Y. S. and Kim, S. B. 2005 Extracting optimization and physical properties of yellowfin tuna (*Thunnus albacares*) skin gelatin compared to mammalian gelatins. *Food Hydrocolloids* 19: 221–229.
- Choi, Y. J., Im, Y. S., Kim, S. M. and Cho, Y. J. 1999. Enzymatic method for measuring ATP-related compound in fish sauces. *J. Kor. Fish. Soc.* 32: 385-390.
- Choi, S. S. and Regenstein, J. M. 2000. Physicochemical and sensory characteristics of fish gelatin. *J. Food Sci.* 65: 194–199.
- Chong, A. S. C., Hashim, R., Chow-Yang, L. and Ali, A. B. 2002. Partial characterization and activities of proteases from the digestive tract of discus fish (*Sympphysodon aequifasciata*). *Aquaculture*. 203: 321-333.
- Chu, C. H. and Bjeldanes, L. F. 1981. Effect of diamines, polyamines and tuna fish extract on the binding of histamine to mucin *in vitro*. *J. Food. Sci.* 47: 7980-7988.
- Chytiri, S., Chouliara, I., Savvaidia, I. N. and Kontominas, M. G. 2004. Microbiological, chemical and sensory assessment of ice whole and filleted aquacultured rainbow trout. *J. Food Microbiol.* 21: 157-165.

- Clarks, J., Macdonald, N. L. and Stark, J. R. 1985. Metabolism in marine flatfish. II. Measurement of elastase activity in the digestive tract of dover sole (*Solea solea L.*). Comp. Biochem. Physiol. 81B: 677-684.
- Clarks, J., Macdonald, N. L. and Stark, J. R. 1985. Metabolism in marine flatfish. III. Measurement of elastase activity in the digestive tract of dover sole (*Soles solea L.*). Comp. Biochem. Physiol. 81B: 695-700.
- Cohen, T., Gertler, A. and Birk, A. Y. 1981. Pancreatic proteolytic enzymes from carp (*Cyprinus carpio*)-I. Purification and physical properties of trypsin, chymotrypsin, elastase and carboxypeptidase B. Comp. Biochem. Physiol. 69B: 639-646.
- Collette, B. B. 1986. Fishes of the North-Eastern Atlantic and the Mediterranean V. II (Whitehead, P. J. P., Bauchot, M. L., Hureau, J. C. Nielsen, J. and Tortonese, E. eds.). p. 981-997. Bungay: The United Nations Educational, Scientific and Cultural Organization.
- Connell, J. J. 1975. Control of fish Quality, 1st Ed. Fishing News Books Limited, London.
- Conway, E. J. and Byrne, A. 1936. An absorption apparatus for the micro-determination of certain volatile substances I. The micro-determination of ammonia. J. Biochem. 27: 419-429.
- Dapkevicius, M. L. N. E., Nout, M. J. R., Rombouts, F. M., Houben, J. H. and Wymenga, W. 2000. Biogenic amine formation and degradation by potential fish silage starter microorganisms. Inter. J. Food Microbiol. 57: 107-114.

- Dawood, A. A., Karkalas, J., Roy, R. N. and Williams, C. S. 1988. The occurrence of non-volatile amines in chilled storage rainbow trout (*Salmo irideus*). Food Chem. 27: 33-45.
- Debevere, J. and Baskou, G. 1996. Effect of Modified atmosphere packaging on the TVB/TMA producing microflora of cod fillets. Inter. J. Food Microbiol. 31: 221-229.
- Delaage, M., Abita, J. P., Lazdunski, M. 1968. Physico-chemical properties of bovine chymotrypsinogen B. Eur. J. Biochem. 5: 285-293.
- Del Rosario, S. R., Maldo, S. M. 1984. Biochemistry of patis formation. Activity of cathepsins in patis hydrolysates. Phil. Agric. 67: 167-175.
- De Poix, A., Rouet-Mayer, A. and Philippon, J. 1980. Combined action of chlorides and ascorbic acid on inhibition of enzymic browning in apple puree. Lebensmittel-Wissenschaft und -Technologie. 13: 105-110.
- Dimes, L. E., Garcia-Carreno, F. L. and Haard, N. F. 1994. Estimation of protein digestibility: III. Studies on the digestive enzymes from the pyloric caeca of rainbow trout and salmon. Comp. Biochem. Physiol. 109A: 349-360.
- Dohmoto, N. Wang, K. C. Mori, T. Kimura, I., Koriyama, T. and Abe, H. 2001. Development of a new type fish sauce using the soy sauce fermentation method. Nippon Suisan Gakkaishi. 67: 1103-1109.
- Eitenmiller, R. R., Orr, J. H., Wallis, W. W. 1982. Histamine formation in fish. In Chemistry and Biochemistry of Marine Food Products, Microbial and Biochemical Condition. (Martin, R. E., Flick , G. J., Hebard, C. E. and Ward, D. R. eds.), p. 39-50. Avi, Westport, CT. USA.

- El-beltagy, A. E., El-Adawy, T. A., Rahma, E. H. and El-Bedawey, A. A. 2004. Purification and characterization of an acidic protease from the viscera of boltifish (*Tilapia nilotica*). *Food Chem.* 86: 33-39.
- Eshel, A., Lindner, P., Smirnoff, P., Newton, S. and Harpaz, S. 1993. Comparative study of proteolytic enzymes in the digestive tracts of the European sea bass and hybrid striped bass reared in freshwater. *Comp. Biochem. Physiol.* 106A: 627-634.
- Faid, M., Zouiten, A., Elmarrakchi, A. and Achkari-Begdouri, A. 1997. Biotransformation of fish waste into stable feed ingredient. *Food Chem.* 60: 13–18.
- Fang, L. S. and Chiou, S. F. 1989. Effect of salinity on the activities of digestive proteases from the tilapia fish (*Oreochromis niloticus*) in different culture environments. *Comp. Biochem. Physiol.* 93A: 439-443.
- FAO. 1997. Review of the state of world fishery resources: Marine fisheries C. Species topics 1. Global resources of tuna and tuna-like species. FDA Fisheries Circular NO.920 FIRM/C920.
- Fernandez-Salguero, J. and Mackie, I. M. 1987. Comparative rates of spoilage of fillets whole fish during storage of haddock (*Melanogrammus aeglefinus*) and herring (*Clupea harengus*) as determination by the formation of non-volatile and volatile amines. *Intern. J. Food Sci. Technol.* 22: 385-390.
- Finne, G., 1992. Non-protein nitrogen compounds in fish and shellfish. In *Advances in seafood biochemistry, Composition and Quality*. (Flick, J. G. Martin, R. E. Eds.). p. 393-401. Technomic Publishing. Pennsylvania.

- Frank, H. A. 1985. VI. 1. Use of normographs to estimate histamine formation in tuna. In Histamine formation in Marine Products, Production by Bacteria, Measurement and Prediction of pH Formation, FAO Fisheries Technical Paper No. 252. p. 18-20. Food and Agriculture Organization of the United Nations. Rome.
- Frank, H. A., Baranowski, J. D., Chongiriatana, M., Brust, P. A. and Premaratne, R. J. 1985. Identification and decarboxylase activitys of bacteria isolated from decomposed mahimahi (*Coryphaena hippurus*) after incubation at 0 and 32 ° C. Intern. J. Food Microbiol. 2: 331-340.
- Frøkjaer S. 1994. Use of hydrolysates for protein supplementation. Food Technol 48 (4): 86–88.
- Fujii, T., Kurihara, K. and Okuzumi, M. 1994. Viability and histidine decarboxylase activity of halophilic histamine-forming bacteria during frozen storage. J. Food Prot. 57: 611-613.
- Fukami, K., Funatsu, Y., Kawasaki, K. and Watabe, S. 2004. Improvement of fish sauce odor by treatment with bacteria isolated from the fish sauce mush (Moromi) made from Frigate mackerel. J. Food Sci. 69: 45-49.
- Fukami, K., Ishiyama, S., Yaguramaki, H., Masuzawa, T., Nabeta, Y., Endo, K. and Shimoda, M. 2002. Identification of distinctive volatile compounds in fish sauce. J. Agric. Food Chem. 50: 5,412-5,416.
- Funatsu, Y., Sunago, R., Konagawa, S., Imai, T., Kawasaki, K. and Takeshima, F. 2000. A comparison of extractive components of a fish sauce prepared from frigate mackerel using soy sauce Koji with those of Japanese-made fish sauces and soy sauce. Nippon Suisan Gakkaishi. 66: 1036-1045.

- Garcia-Carreno, F. L. and Haard, N. F. 1993. Characterization of proteinase classes in langostilla (*Pleuroncodes planipes*) and crayfish (*Pacifastacus astacus*) extracts. J. Food Biochem. 17: 97-113.
- Geiger, E., Courtney, G. and Schnakenberg, G. 1994. The content and formation of histamine in fish muscle. Arch. Biochem. 3:311-319.
- Gildberg, A. 1988. Aspartic protease in fishes and aquatic invertebrates. Comp. Biochem. Physiol. 91B:425-435.
- Gilberg, A. 1992. Recovery of proteinase and protein hydrolysate from fish viscera. Biores. Technol. 39: 271-276.
- Gildberg, A., 2001. Utilization of male Arctic capelin and Atlantic cod intestines for fish sauce production evaluation of fermentation conditions. Biores. Technol. 76: 119-123.
- Gildberg A, Batista I, Strøm E. 1989. Preparation and characterization of peptones obtained by a two-step enzymatic hydrolysis of whole fish. Biotechnol Appl. Biochem. 11:413–23.
- Gilberg, A., Hermes, J. E. and Orejana, F.M. 1984. Acceleration of autolysis during fish sauce fermentation by adding acid and reducing the salt content. J. Sci. Food Agric. 35: 1363-1369.
- Gildberg, A., Olsen, R. L. and Bjarnason, J.B. 1990. Catalytic properties and chemical composition of pepsin from Atlantic cod (*Gadus morhua*). Comp. Biochem. Physiol. 69B: 323-330.

- Gildberg, A. and Raa, J. 1983. Purification and characterization of pepsins from Arctic fish capelin (*Mallotus villosus*). Comp. Biochem. Physiol. 75A: 337-342.
- Gildberg, A. and Thongthai, C. 2001. The effect of reduced salt content and addition of halophilic lactic bacteria on quality and composition of fish sauce made from sprat. J. Aquat. Food Product Technol. 10(1): 77-88.
- Gilberg, A. and Xian-Quan, S. 1994. Recovery of tryptic enzymes from fish sauce. Proc. Biochem. 29: 151-155.
- Giraffa, G., 2002. Enterococci from foods. FEMS Microbiology Reviews, 26: 163-171.
- Gram, L., Huss, H. H., 1996. Microbial spoilage of fish and fish products. Intern. J. Food Microbiol. 33: 121-137.
- Gudmundsson, M. 2002. Rheological properties of fish gelatins. J. Food Sci. 67: 2172-2176.
- Guérard, F., Dufossé, L., Broise, D. De La. And Binet, A. 2001. Enzymatic hydrolysis of proteins from yellowfin tuna (*Thunnus albacares*) wastes using Alcalase. J. Mol. Catal. B: Enzymol. 11: 1051-1059.
- Guérard, F., Guimas, L. and Binet, A. 2002. Production of tuna waste hydrolysates by a commercial neutral protease preparation. J. Mol. Catal. B: Enzymol. 19: 489-495.
- Guérard, F. and LeGal, Y. 1987. Characterization of a chymosin-like pepsin from the dogfish (*Selachiorhinus canicula*). Comp. Biochem. Physiol. 88B: 823-827.

- Guizani, N., Al-Busaidy, M. A., Al-Belushi, I. M., Mothershaw, A. and Rahman, M. S. 2005. The effect of storage temperature on histamine production and the freshness of yellowfin tuna (*Thunnus albacares*). *Food. Res. Intern.* 38: 215-222.
- Guizani, N., Rolle, R. S., Marshall, M. R. and Wei, C. I. 1991. Isolation, purification and characterization of a trypsin from the pyloric caeca of a trypsin from the pyloric caeca of mullet (*Mugil cephalus*). *Comp. Biochem. Physiol.* 98B: 517-521.
- Haard, N. F. 1986. Characterization with casein and milk substrate and influence of sepharose immobilization on salt activation, temperature characteristics and milk clotting reaction. *J. Food Sci.* 51: 313-316.
- Haard, N. F., 1994. Protein hydrolysis in seafoods. In *Seafood chemistry. Processing technology and quality*. (Shahidi, F. and Botta, J. R. Eds.). p. 10-33. Chapman & Hall, New York.
- Haard, N. F. and Simpson, B. K. 1994. Proteases from aquatic organisms and their uses in the seafood industry. In *Fisheries processing: Biotechnology applications*. Chapman and Hall Publishing, London.
- Hálász, A., Barath, A., Simon-Sakardi, L. and Holzafel, W. 1994. Biogenic amines and their micro-organisms in food. *Trends Food Sci. Technol.* 5: 42-49.
- Hansen, L. T., Gill, T. And Hessa, H. H. 1995. Effect of salt and storage temperature on chemical microbiological and sensory changes in cold smoked salmon. *Food Res. Intern.* 28: 123-130.
- Hayashi, Y. 1970. Rna in midgut of tussock moth, *orgyia leucostigma*, infected with cytoplasmic-polyhedrosis virus. *Can. J. Microbiol.* 16: 1101-1107.

- Hebard, C. E., Flick, G. J., Martin, R. E., 1982. Occurrence and significance of trimethylamine oxide and its derivatives in fish and shellfish. In Chemistry and biochemistry of marine food products. (Martin, R. E., Flick, G. J., Hebard, C.E., Ward, D. R. eds.). p. 149-304. AVI Publishing, Westport, Connecticut, USA.
- Hendel, C. E., Bailey, G. F., Taylor, D. H. 1950. Measurement of non enzymatic browning of dehydrated vegetables during storage. Food Technol. 3(9): 44-48.
- Hernández-Herrero, M. M., Raig-Sagués, A. X., Rodríguez-Jerez, J. J. and Mora-ventura, M. T. 1999. Halotolerant during the ripening of salted anchovies (*Engraulis encrasicholus*). J. Food. Prot. 62: 509-514.
- Heu, M. S., Kim, H. R. and Pyeum, J. H. 1995. Comparison of trypsin and chymotrypsin from the viscera of anchovy (*Engraulis japonica*). Comp. Biochem. Physiol. 112B: 557-568.
- Hidalgo, M. C., Urea, E. and Sanz, A. 1999. Comparative study of digestive enzymes in fish with different nutritional habits. Proteolytic and amylase activities. Aquaculture. 170:267-283.
- Hjelmeland, K. and Raa, J. 1982. Characteristics of two trypsin type isozymes isolated from the Arctic fish capelin (*Mallotus villosus*). Comp. Biochem. Physiol. 71B: 557-562.
- Holtmann, L., Rustad, T. 2004. Iced storage of Atlantic salmon (*Salmo salar*)-effects on endogenous enzymes and their impact on muscle proteins and texture. Food Chem. 87: 31-41.

- Hoyle, N .T. and Merritt, J. H. 1994. Quality of fish protein hydrolysates from herring (*Clupea harengus*). *J. Food Sci.* 59: 76-79.
- Hultmann, L. and Rustad, T. (2004). Iced storage of Atlantic salmon (*Salmo salar*) effect on endogenous enzymes and their impact on muscle proteins and texture. *Food Chem.* 87: 31-41.
- Huss, H. H., 1995. Quality and quality changes in fresh fish. FAO fisheries technical paper no. 348.
- Ienistea, C. 1971. Bacterial production and destruction of histamine in foods, and food poisoning caused by histamine. *Die Nahrung.* 15: 109-113.
- Ijomah, P., Clifford, M. N., Walker, R., Wright, J., Hardy, R. and Murray, C. K. 1992. Further volunteer studies on scombrotoxicosis. In *Pelagic Fish, The Resource and its Exploitation* (Burt, J. R., Hardy, R., Whittle, K. J. Eds.). p. 194-199., Fishing News Books, Oxford.
- Ijong, F. G. and Ohta, Y. 1996. Physicochemical and microbiological changes associated with bakasang processing, a traditional Indonesian fermented fish sauce. *J. Sci Food Agric.* 71: 69-74.
- Ishizuka, H., Horinouchi, S. and Beppu, T. 1993. Putrescine oxidase of *Micrococcus rubens*: primary structure and *Escherichia coli*. *Gen. Microbiol.* 139: 425-432.
- Javeed, A. and Mahendrakar, N. S. 1996. Autolysis and rancidity development in tropical freshwater fish viscera during fermentation. *Biores. Technol.* 58: 247-251.

- Jay, M. J. 1996. Intrinsic and extrinsic parameters of foods that affect microbial growth. In Modern Food Microbiology. 5th ed. Chapman and Hall. New York.
- Jeyasekaran, G., Ganesan, P., Maheswari, K., Shakila, R. J. and Sukumar, D. 2004. Effect of delayed icing on the microbiological quality of tropical fish: barracudas (*Sphyraena barracuda*). J. Food Sci. 69, 197-200.
- Jiang, S. T., Lee, B. L., Tsao, C. Y. and Lee, J. J. 1997. Mackerel cathepsin B and L effects on thermal degradation of surimi. J. Food Sci. 62: 310– 315.
- Jiang, S. T., Wang, Y. T. and Chen, C. S. 1991. Purification and characterization of a proteinase identified as cathepsin D from tilapia muscle (*Tilapia nilotica* x *Tilapia aurea*). J. Agric. Food Chem.39: 237–241.
- Jonas, E., Ragyanszki, M., Olah, J. and Boross, L. 1983. Proteolytic digestive enzymes of carnivorous (*Silurus glanis L.*) herbivorous (*Hopophthalmichthys molitrix Vol.*) and omnivorous (*Cyprinus carpio*) fishes. Aquaculture. 30: 145-154.
- Kahyaoglu, T. and Kaya, K. 2005. Modelling of moisture, color and texture changes in sesame seeds during the conventional roasting. J. Food Engineer. In press.
- Kalisz, M. H. 1988. Microbial proteases. In Advances in Biochemical, Engineering Biotechnology, vol. 35. (Fiechter, A. ED). p1-65.
- Karacam, H., Kutlu, S. and Kose, S. 2002. Effect of salt concentrations and temperature on the quality and shelf-life of brined anchovies. Intern. J. Food Sci. Technol. 37: 19-28.

- Kawashima, K. and Yamanaka, H. 1996. Free amino acids responsible for the browning of cooked scallop adductor muscle. Fish. Sci. 62: 293-296.
- Kim, J. H., Ahn, H. J., Yook, H. S., Kim, K. S., Rhee, M. S., Ryu, G. H. and Byun, M. W. 2004. Colour, flavour, and sensory characteristics of gamma-irradiated salted and fermented anchovy sauce. Rad. Phys. Chem. 69: 179-187.
- Kim, S. H., Ben-gigirey, B., Baeeos-Velazquez, J., Price, R. J. and An, H. 2000. Histamine and biogenic amine production by *Morganii* isolated from temperature-abuse albacore. J. Food Prot. 63: 244-251.
- Kim, I. S., Choi, Y. J., Heu, M.S., Cho, Y. J., Im, Y. S., Gu, Y. S., Yeo, S. G. and Park, J. W. 1999. Peptide properties of rapid salted and fermented anchovy sauce using various proteases.I. Hydrolysis of anchovy sauce and actomyosin by various proteases. J. Kor. Fish. Soc. 32: 481: 487.
- Kim, S. H., Field, K. G., Morrissey, M. T., Price, R. J., Wei, C. and An, H. 2001. Source and identification of histamine producing bacteria from fresh and temperature abused albacore. J. Food. Prot. 64: 1035-1044.
- Kim, S., Jeon, Y., Byeun, H., Kim, Y. and Lee, C. 1997. Enzymatic recovery of cod frame proteins with crude proteinase from tuna pyloric caeca. Fish. Sci. 63: 421-427.
- Kinoshita, M., Toyohara, H. and Shimizu, Y. 1990. Diverse distribution of four distinct types of modori (gel degradation) inducing proteinases among fish species. Nippon Suisan Gukkaishi. 56: 1485-1492.

- Kirschke, H. and Barrett, A.J. 1987. Chemistry of lysosomal proteases. In Lysosomes, Their role in protein breakdown. (Glaumannt, H. and Ballard, F.J. eds.). p. 193. New York: Academic Press.
- Klausen, N. K. and Huss, H. H. 1987. Growth and histamine production by *Morganella morganii* under various temperature conditions. Intern. J. Food Microbiol. 5: 147-156.
- Klomklao, S., Benjakul, S. and Visessanguan, W. 2004. Comparative studies on proteolytic activity of splenic extract from three tuna species commonly used in Thailand. J. Food Biochem. 28: 355-372.
- Koutsoumanis, K., Lampropoulou, K., and Nychas, G.E. 1999. Biogenic amines and sensory changes associated with the microbial flora of Mediterranean gilt-head sea bream (*Sparus aurata*) stored aerobically at 0, 8, and 15°C. J. Food Protect. 62; 398-402.
- Kristjansson, M. 1991. Purification and characterization of trypsin from the pyloric caeca of rainbow trout (*Oncorhynchus mykiss*). J. Agric. Food. Chem. 39: 1738-1742.
- Kristinsson, H. G. and Rasco, B. A. 2000. Fish protein hydrolysates: production, biochemical, and functional properties. Crit. Rev. Food Sci. Nutr. 40: 43–81.
- Kurbanoglu E. B. and Algur O. F. 2002. The influence of ram horn hydrolyzate on the crop yield of the mushroom Agaricus bisporus. Sci. Hort. 94: 351–357.
- Labuza, T. P. and Baisier, W. M. 1992. Kinetics of nonenzymatic browning. In Physical chemistry of foods (Schwartzberg, H. G. and Hartel R. W., eds.). p. 595-649. Marcel Dekker, New York.

- Lahl, W. J. and Braun, S. D. 1994. Enzymatic production of protein hydrolysates for food use. *Food Technol.* 58(10): 68.
- Lakshmanan, R., Shakila, R. J. and Jeyasekaran, G. 2002. Changes in the halophilic amine forming bacteria flora curing salt-drying of sardines (*Sardinella gibbosa*). *Food Res. Intern.* 35: 541-546.
- Lannelongue, M., Finne, G., Hanna, M.D., Nickelson, F. and Vanderzant, C. 1982. Microbiological and chemical changes during storage of swordfish (*Xiphia sgladius*) steaks in retail packages containing CO₂ enriched atmospheres. *J. Food Prot.* 45: 1197-1203.
- Lazano, R., Drudis, R. and Ibarz, A. 1994. Enzymatic browning in apple pulps. *J. Food Technol.* 59: 564-567.
- Lee, B. J. and Hendricks, D. G. 1997. Antioxidant effects of L-carnisine on liposomes and beef homogenates. *J. Food Sci.* 62: 931-934.
- Lee, Y. S., Homma, S. and Aida, K. 1997. Characterization of melanoidin in soy sauce and fish sauce by electrofocusing and high performance gel permeation chromatography. *Nippon Shokuhin Gakkaishi.* 34: 313-319.
- Lee, Y. Z., Simpson, B. K. and Haard, N. F. 1982. Supplementation of squid fermentation with proteolytic enzymes. *J. Food Biochem.* 6: 127-134.
- Lehane, L. 2000. Ciguatera update. *Med. J. Aust.* 172: 176-179.
- Lericci, C. R., Barbanti, D., Manzano, M. and Cherubin, S. 1990. Early indicators of chemical changes in foods due to enzymic or non enzymic browning reactions. 1: study on heat treated model systems. *Lebensmittel-Wissenschaft und Technologie.* 23: 289-294.

- Lertsiri, S., Phontree, K., Thepsingha, W. and Bhumiratana. 2003. Evidence of enzymatic browning due to laccase-like enzyme during mash fermentation in Thai soybean paste. *Food Chem.* 80: 171-176.
- Leuschner, R. G. K. and Hammes, W. P. 1998. Tyramine degradation by micrococci during ripening of fermented sausage. *Meat Sci.* 49: 289-296.
- Leuschner, R. G., Heidel, M. And Hammes, W. P. 1998. Histamine and tyramine degradation by food fermenting microorganisms. *Intern. J. Food Microbiol.* 39: 1-10.
- Lopetcharat, K. 1999. Fish sauce: The alternative solution for Pacific whiting and surimi by products. M. Sc. Thesis. Corvallis, OR, Oregon State University.
- Lopetcharat, K., Choi Y.J., Park J.W. and Daeschel, M.A. 2001. Fish sauce products and manufacturing: a review. *Food Rev. Intern.* 17: 65-88.
- Lopetcharat K. and Park, J. W. 2002. Characteristics of fish sauce made from Pacific whiting and surimi by products during fermentation stage. *J. Food Sci.* 67: 511-516.
- Love, R. M. 1980. Histidine. In *The Chemical Biology of Fishes, Advances 1968-1977*, Vol. 2, Academic Press, London, New York, Toronto, Sydney, San Francisco, p. 427.
- Lowry, O. H., Rosebrough, N. J., Fan, A. L. and Randall, R. J. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193: 256-275.
- Mahmoud, M.I. 1994. Physiochemical and functional properties of protein hydrolysates in nutritional products. *Food Technol.* 58(10): 89–95Makan, M. 2001. Kinetic

- of color changes of kiwifruit during hot air and microwave drying. *J. Food Engineer.* 48: 169-175.
- Martinez, A., Olsen, R. L. and Serra, J. L. 1988. Purification and characterization of two trypsin-like enzymes from the digestive tract of anchovy (*Engraulis encrasicholus*). *Comp. Biochem. Physiol.* 91B: 677-684.
- Martinez, A. and Serra, J. L. 1989. Proteolytic activities in the digestive tract of anchovy (*Engraulis eucrasi cholus*). *Comp. Biochem. Physiol.* 93B: 61-66.
- Martins, S. I. F. S., Marcelis, A. T. M. and Van Boekel, M. A. J. S. 2003. Kinetic modelling of the Amadori N-(1-Deoxy-D- fructose-1-yl) glycine degradation pathways. Part I- reaction mechanism. *Carbohydr. Res.* 338: 1651-1663.
- Martuscelli, M., Crudele, M. A., Gardini, F. and Suzzi, G. 2000. Biogenic amine formation and oxidation by *Staphylococcus xylosus*. *Lett. Appl. Microbiol.* 31: 228-232.
- Mason, R. W., Toylor, M. A. and Etherington, D. J. 1984. The purification and properties of cathepsin L from rabbit liver. *J. Biochem.* 17: 209-217.
- Mauron, J. 1981. The Maillard reaction in food: Criti. Rev. from Nutri. Food Nutri. Sci. 5: 5-35.
- Mazorra-Manzano, M. A., Pacheco-Aguilar, R., Diaz-Rojas, E. I. and Lugo-Sanchez, M. E. 2000. Postmortem changes in black skipjack muscle during storage in ice. *J. Food Sci.* 65: 774-779.
- Mihalyi, E. 1978. Application of proteolytic enzymes to protein structure studies. CRC Press, Boca Raton, Florida.

- Morales, F. J. and Jimenez-Perez, S. 2001. Free radical scavenging capacity of Maillard reaction products as related to color and fluorescence. Food Chem. 72: 119-125.
- Morales, F. J., Romero, C. and Jimenez-Perez, S. 1996. Fluorescence associated with Maillard reaction in milk and milk-resembling systems. Food Chem. 67: 423-428.
- Morales, F. J and Van Boekel, M. A. J. S. 1997. A study on advanced Maillard reaction in heated casein-sugar solution: Fluorescence accumulation. Intern. Dairy J. 7: 675-683.
- Morioka, K., Fujii, S., Iton, Y., Chengchu., L. and Obatake, A. 1999. Recovery of amino acid from protein in the head and viscera of frigate mackerel by autolysis. Fish. Sci. 65: 588-591.
- Morrissey, M. T., WU, J. W., Lin, D. D. and An, H. 1993. Protease inhibitor effects on torsion measurements and autolysis of Pacific Whiting surimi. J. Food Sci. 58: 1050-1054.
- Moss, J. R. and Otten, L/ 1989. A relation ship between color development and moisture content during roasting of peanut. J. Can. Inst. Food Sci. Technol. 22: 34-39.
- Munilla-Moran, R. and Rey, F. B. 1996. Digestive enzymes in marine species. Proteinase activities in gut from redfish (*Sebastes mentella*), seabream (*Sparus aurata*) and turbot (*Scophthalmus maximus*). Comp. Biochem. Physiol. 113B: 395-402.

- Namwong, S., Tanasupawat, S., Smitinont, T., Visessanguan, W., Kudo, T. and Itoh, I. 2005. Isolation of *Lentibacillus salicampi* strains and *Lentibacillus* sp. nov. from fish sauce in Thailand. Intern. J. Syst. Evol. Microbiol. 55: 315-320.
- Nicoli, M. C., Anese, M. and Parpinel, M. 1999. Influence of processing on the antioxidant properties of fruit and vegetable. Trends Food Sci. Technol. 10: 94-100.
- Niwa, E. 1992. Chemistry of surimi gelation. In Surimi technology. (Lanier, T.C. and Lee, C.N. eds.). p. 389-427. Marcel Dekker. New York.
- Noda, M. and Murakami, K. 1981. Studies on proteases from the digestive organs of sardine. II. Purification and characterization of two acid proteases from the stomach. Biochem. Biophys. Acta. 65B: 27-34.
- Noda, M., Van, T. V., Kusakabe, I. and Murakami, K. 1982. Substrate specificity and salt inhibition of five proteinases isolated from the pyloric caeca and stomach of sardine. Agric. Biol. Chem. 46: 198-203.
- Oetterer, M., Perujo, S. D., Gallo, C. R., Arruda, L. F. D., Borghesi, R. and Cruz, A. M. P. D. 2003. Monitoring the sardine (*Sardinella brasiliensis*) fermentation process to obtain anchovies. J. Food. Sci. Technol. 60: 511-517.
- Okazaki, T., Shigeta, Y., Aoyama, and Namba, K. 2003. Autolysis of unsalted fish protein under pressurization. Fish. Sci. 69: 1257-1262.
- Okuzumi, M., Okuda, S. and Awano, M. 1981. Isolation of psychrophilic and halophilic histamine-forming bacteria from *Scomber japonicus*. Bull. Jpn. Soc. Sci. Fish. 47: 1591-1598.

- Omar, N. M., 2002. Indole as an index of spoilage of mackerel (*Restrelliger kanagurta*) during storage at ambient temperature and in ice. Fish. Sci. 39: 142-145.
- Ooshiro, Z. 1971. Studies on proteinase in the pyloric caeca of fish-II. Some properties of proteinase purified from the pyloric caeca of mackerel. Bull. Jap. Soc. Sci. Fish. 26: 685-690.
- Orejana, F. M., Liston, J. 1981. Agents of proteolysis and its inhibition in patis (fish sauce fermentation). J. Food Sci. 47: 198-203, 209.
- Özogul, F., Gökbüyük, C., Özogul, Y. and Özyurt, G. 2005. Biogenic amine production and nucleotide rates in gutted wild sea bass (*Dicentrarchus labrax*) stored in ice wrapped in aluminium foil and wrapped in cling film at 4 °C. Food Chem. (In press).
- Pastoriza, L., Sampedro, G., Herrena, J. J. and Cabo, M. 1996. Effect of modified atmosphere packaging on shelf-life of iced fresh hake slice. J. Sci. Food. Agric. 71: 541-547.
- Poosaran, N. 1986. Fish sauce I: acid hydrolysis at ambient temperature. Songklanakarin J. Sci. Technol. 8: 43-46.
- Prasertsan, P., Wuttijumnong, P., Sophanodora, P. and Choorit, W. 1988. Seafood processing industries within Songkhla-Hatyai region: The survey of basic data emphasis on wastes. Songklanakarin. J. Sci. Technol. 10: 447-451.
- Quiñones, B. 2000. Partial purification and characterization of trypsin from the pyloric caeca and intestine of the queen snapper (*Etelis oculatus*). Master's Thesis. University of Puerto Rico-Mayagüez Campus.

- Raksakulthai, N., and Haard, N. F. 1992. Correlation between the concentration of peptides and amino acids and the flavour of fish sauce. ASEAN Food J. 7: 86-90.
- Raksakulthai, N., Lee, Y. Z. and Haard, N. F. 1986. Effect of enzyme supplements on the production of fish sauce from male capelin (*Mallotus villosus*). Can. Inst. J. Food Sci. Technol. 19: 28-33.
- Rawles, D. D., Flick, G. J. and Martin, R. E. 1996. Biogenic amine in fish and shellfish. Adv. Food Nutr. Res. 39: 329-364.
- Riaz, M., Fatima, R. and Qadri, R. B. 1986. Biochemical changes during the preparation of fish sauces. Trop Sci. 26: 213-222.
- Ritchie, A. H. and Mackie, I. M. 1997. The formation of diamine and polyamines during storage of mackerel (*Scomber scombrus*). In Advances in fish sciences and technology. (Connell, J. J. ed.). p. 489-494. Fishing News Books. Surrey.
- Rodtong, S., Nawong, S. and Yongsawatdikul, J. 2005. Histamine accumulation and histamine-forming bacteria in Indian anchovy (*Stolephorus indicus*). J. Food Microbiol. 22: 475-482.
- Roscoe, H. G., Kupfer, D. 1972. A new histaminase assay. Anal. Biochem. 47: 418-425.
- Sabapathy, U. and Teo, L. H. 1993. A quantitative study of some digestive enzymes in the rabbitfish (*Siganus canaliculatus*) and the sea bass (*Lates calcarifer*). J. Fish Biol. 42: 595-602.

- Saila, S. B. and Norton, V. J. 1974. Tuna: status, trends, and alternative management arrangement. Washington: Resource for the future, Inc.
- Saisithi, P., 1994. Traditional fermented fish: fish sauce production. In *Fisheries Processing, Biotechnology Application* (Martin, A. M. ed). p. 111-131. Chapman & Hall, London.
- Saisithi, P., Kasemsarn, B. O., Liston, J. and Dolla, A. M., 1966. Microbiology and chemistry of fermented fish. *J. Food Sci.* 31: 105-110.
- Sanceda, N.G., Kurata, T. and Aakawa, N. 1990. Overall quality and sensory acceptance of a lysine-fortified fish sauce. *J. Food Sci.* 55: 983-988.
- Sanceda, N. G., Kurata, T. and Arakawa, N. 1996. Accelerated fermentation process for the manufacture of fish sauce using histidine. *J. Food Sci.* 61: 220-225.
- Sanceda, N., Suzuki, E. and Kurata, T (2003). Quality and sensory acceptance of fish sauce partially substituting sodium chloride or natural salt with potassium chloride during the fermentation process. *Intern. J. Food. Sci. Technol.* 38: 435-443.
- Sanceda, N., Suzuki, E. and Kurata, T. 2000. Development of normal and branched chain volatile fatty acids during the fermentation process in the manufacture of fish sauce. *J. Sci. Food Agric.* 81: 1013 – 1018.
- Sanceda, N. G., Suzuki, R., Ohashi, M. and Kurata, T., 1999. Histamine behavior during the fermentation process in the manufacture of fish sauce. *J. Agric. Food Chem.* 47: 3596-3600.
- Sanchez-Chiang, L., Cisternas, E. and Ponce, O. 1987. Partial purification of pepsins of adult and Juvenile salmon fish (*Oncorhynchus keta*). Effect of NaCl on proteolytic activities. *Comp. Biochem. Physiol.* 87B: 793-797.

- Sands, A. and Crisan, E. V. 1974. Microflora of fermented korean seafoods. *J. Food. Sci.* 39: 1002-1974.
- Sapin-Jaloustre, H. and Sapin-Jaloustre, J. 1957. A little-known food poisoning: histamine poisoning from tuna. *Concours Med Paris.* 79: 2705-2708.
- Schayer, R. W. 1960. Relationship of induced histamine synthesis to shock from stress and from endotoxin. *Am. J. Physiol.* 198. 1187-1192.
- Schwelberger, H. G., and Bodner, E. (1997). Purification and characterization of diamine oxidase from porcine kidney and intestine. *Biochem. Biophys. Acta.* 152-164.
- Senöz, B., Isykly, N., & Coksöyler, N. (2000). Biogenic amines in Turkish sausages (sucuk). *J. Food Sci.* 65: 764–767.
- Seymour, T. A., Morrissey, M. T., Peters, M. Y. and An, H. 1994. Purification and characterization of Pacific whiting proteases. *J. Agric. Food Chem.* 42: 2634-2638.
- Shahidi, F. 1994. Proteins from seafood processing discards. In *Seafood proteins.* (Sikorski, Z. E., Pan, B. S. and Shahidi, F. eds.). p. 171-193. Chapman and Hall, New York.
- Shahidi, F., X. Q. Han, and J. Synowiecki. 1995. Production and Characteristics of protein hydrolysates from capelin (*Mallotus villosus*). *Food Chem.* 53: 285-293.
- Shakila, R. J., Vijayalakshmi, K. and Jeyasekaran, G. 2003. Changes in histamine and volatile amine in six commercially important species of fish of the Toothukkudi coast of Tamil Nadu, India stored at ambient temperature. *Food Chem.* 83: 347-352.

- Shemy, M. E. and Levin, R. E. 1997. Characterization of affinity-purified trypsin from hybrid tilapia (*Tilapia niloticus/surea*). *J. Food biochem.* 21: 163-175.
- Sherekar, S. V., Doke, S. N., Gore, M. S. and Ninjoor, V. 1986. Proteinases of tropical fish and their role in autolysis. *Indian J. Exp. Biol.* 24: 440-444.
- Shewan, J.M. 1962. The bacteriology of fresh and spoiling fish and some related chemical changes. In Recent advances in food science. Hawthorn, J. and Muil Leitch, J. (eds.). Butterworths, London. 1: 167-193.
- Shih, I-L., Chen, L-G., Yu, T-S, Chang, W-T. and Wang, S-L. 2003. Microbial reclamation of fish processing wastes for the production of fish sauce. *Enz. Microbiol. Technol.* 33: 154-162.
- Siebert, G. and Bottke, J. 1963. Enzyme in der leibeshole von fischen. *Arch. Fur Fishchereiwiss.* 14: 57-68.
- Sikorski, Z.E. Gildberg, A. and Ruiter, A. 1995. Fish Products: Fish and Fishery Products. Ruiter, A. (edite) CAB. International, Wallingford, United Kingdom 315-346.
- Sikorski, Z.E., Kolokowski, A. and Burt, J. R. 1990. Postharvest biochemical and microbiological changes. In Seafood resources nutritional composition and preservation. (Sikorski, Z. E. ed). p. 55. CRC Press, Inc., USA.
- Silla Santos, M. H. 1996. Biogenic amines: their importance in foods. *Intern. J. Food Microbiol.* 29: 213-231.
- Simpson, B. K. 2000. Digestive proteinases from marine animals. In Seafood Enzymes: Utilization and Influence on Postharvest Seafood Quality. (Haard, N. F. and Simpson, B. K. eds.). p. 191-124. Marcel Dekker, New York.

- Simpson, B. K. and Haard, H. F. 1984. Trypsin from Greenland cod (*Gadus ogac*). Isolation and comparative properties. Comp. Biochem. Physiol. 79B: 613-622.
- Simpson, B. K., Simpson, M. V. and Haard, N. F. 1990. Properties of trypsin from the pyloric caeca of Atlantic cod (*Gadus morhua*). J. Food Sci. 55: 959-961.
- Sipos, T., Rappoport, S., Avnir, D., Ottolenghi, M. and Braum, S. 1992. Properties of trypsin and acid phosphatase immobilized in sol-gel glass matrix. Biotechnol. Appl. Biochem. 15: 227.
- Skrede, G. 1985. Colour quality of black current syrups during storage evaluated by Hunter L, a, b value. J. Food Sci. 50: 514-517, 525.
- Snyder, S. H. and Hendley, E. D. 1968. A simple and sensitive fluorescence assay for monoamine oxidase and diamine oxidase. J. Pharmacol. Exp. Ther. 163: 386-392.
- Sovik, S. L. and Rustad, T. 2006. Effect of season and fishing ground on the activity of cathepsin B and collagenase in by products from cod species. Food Sci. Technol. 39(1) : 43-53.
- Soyiri, I.N., Tano-Debrah, K. and Amoa-Awuaah, A.W. 2003. Physico-Chemical and Quality characteristics of fish sauce produced from tuna processing waste. 2nd International Workshop on Food-based approaches for a healthy nutrition in West Africa: the role of food technologists and nutritionists, 23 -28 November 2003 in Ouagadougou Burkina Faso.
- Squires, E. J., Haard, N. F. and Feltham, L. A. 1986. Gastric proteases of the Greenland cod (*Gadus ogac*) I. Isolation and kinetic properties. Biochem. Cell Biol. 64: 205-214.

- Srikanth, L. N., Khuntia, B. K., Reddy, G. V. S. and Srinivasa, B. R. 1993. Influence of storage temperature on the quality of salted mackerel (*Rastrelliger japonicus*) and pink perch (*Nemipterus japonicus*). J. Sci. Food Agric. 63: 319-322.
- Steel, R. G. D. and Torrie, J. H. 1980. Principles and procedures of statistics; A biometrical approach. 2nd ed. p. 633. McGraw-Hill. New York.
- Stom, T. and Eggum, B. O., 1981. Nutritional value viscera silage. J. Sci. Food Agric. 32: 115 – 120.
- Stratton, J. E., Hutkins, R. W. and Taylor, S. L. 1991. Biogenic amines in cheese and other fermented food. A review. J. Food Prot. 54: 460-470.
- Straub, B. W., Kicherer, M., Schilcher, S. M. and Hammes, W. P. 1995. The formation of biogenic amine by fermentation organisms. Z. Lebensm Unters. Forsch. 201: 79-82.
- Straub, B. W., Tichaczek, P. S., Kicherer, M. and Hammes, W. P. 1994. Formation of tyramine by *Lactobacillus curvatus* LTH 972. Z. Lebensm Unters. Forsch. 1: 9-12.
- Su, H., Lin, T. S. and Lanier, T. C. 1981. Investigation into potential sources of heat stable alkaline protease in mechanically separated Atlantic croaker (*Micropogon undulates*). J. Food Sci. 46: 1654.
- Tanasupawat, S., Ezaki, T., Suzuki, K., Okada, S., Komagata, K. and Kozaki, M. 1992. Characterization and identification of *Lactobacillus pentosus* and *Lactobacillus plantarum* strains from fermented foods in Thailand. J. Gen. Appl. Microbiol. 38: 121-134.
- Tanasupawat, S. and K. Komagata. 1995. Lactic acid bacteria in fermented foods in Thailand. World J. Microbiol. Biotechnol. 11: 253-256.

- Tannenbaum, S.R., R.P. Bates and L. Brodfeld, 1970. Solubilization of fish protein concentrate. II.Utilization of the alkaline-process product. Food.Tecnol. Champaign. 24: 607-609.
- Taylor, S. L. 1986. Histamine food poisoning: Toxicology and Clinical aspects. Crit. Rev. Toxicol. 17(2): 91-128.
- Taylor, S. L. and Sumner, S. S. 1986. Determination of histamine, putrescine, and cadaverine. In (Kramer, D. E., Liston, J. Eds.). p. 235-245. Seafood Quality Determination, Elsevier, Amsterdam.
- Tengjaroenkul, B., Smith, B. J., Caceci, T. and Smith, S. A. 2000. Distribution of intestinal enzyme activities along the intestinal tract of cultured Nile tilapia. (*Oreochromis Miloticus*). Aquaculture. 182: 317-327.
- Thai Industrial Standard. 1983. Local fish sauce standard. Department of Industry, Bangkok, Thailand.
- Thongthai, C. and Siriwongpairat, M. 1990. The sequential quantitation of microorganisms in traditionally fermented fish sauce (nam pla). In Post harvest Technology, Preservation and Quality of Fish in Southeast Asia. (Reilly, P. J. A., Parry, R. W. H. and Barile, L. E. Eds.) p. 61-65. International Foundation for Science. Stockholm.
- Torrisen, K.R. 1984. Characterization of proteases in the digestive tract of Atlantic salmon (*Salmo salar*) in comparison with rainbow trout (*Salmo gairdneri*). Comp. Biochem. Physiol. 77B: 669-674.
- Tungkawachara, S., Park, J. W. and Choi, Y. J. 2003. Biochemical properties and consumer acceptance of Pacific whiting fish sauce. J. Food. Sci. 68: 855-860.

- Van Boekel, M. A. J. S. 1998. Effect of heating on Maillard reactions in milk. *Food Chem.* 62: 403-414.
- Van Veen, J. 1965. The tectonic and stratigraphic history of the Cardoño area, Contabrian Mountains Northwest spain. *Leid geol Meded* 35: 45-104.
- Vázquez, J. A., González, M. P. and Murago, M. A. 2004. Peptones from autohydrolysed fish viscera for nisin and pediocin production. *J. Biotechnol.* 112: 299-311.
- Vecchi, S.D. and Coppes, Z. 1998. Marine fish digestive proteases relevant to food industry and the South-west Atlantic region-a review. *J. Food Biochem.* 20: 193-214.
- Vecht-Lifshitz, S. E., Almas, K. A. and Zomer, E. 1990. Microbial growth on peptones from fish industrial wastes. *Lett. Appl. Microbiol.* 10: 183–186.
- Veciana-Nogues, M. T. M., Vidal-Carou, M. C. and Marine-Font, A. and Vidal-Carou, M. C. 1997. Biogenic amines as higenic quality indicators of tuna relationship with microbial counts ATP-related compounds, volatile amines, and organoleptic changes. *J. Agric. Food Chem.* 45: 2036-2041.
- Vega-Villasante, Nolasco, H. and Civera, R. 1995. The digestive enzymes in marine species. Proteinase activities in gut from redfish (*sebastes mentella*), seabream (*Sparus aurata*) and turbot (*Scophthalmus maximus*). *Comp. Biochem. Physiol.* 113B: 395-402.
- Wade, M. J. 1991. Genetic variance for rate of population increase in natural populations of flour beetles, *Tribolium spp.* *Evolution.* 45: 1574-1584.

- Wang, J. H., Ma, W. C., Su, J. C., Chen, C. S. and Jiang, S. T. 1993. Comparison of the properties of m-calpain from tilapia and grass shrimp muscle. *J. Agric. Food Chem.* 41: 1379-1384.
- Whitaker, J.R. 1994. Classification and nomenclature of enzymes. In *Principles of Enzymology for the Food Sciences*, 2nd ed. p. 367-385. Marcel Dekker, New York.
- Wilaiwan, P. 1990. Halophilic bacteria producing lipase in fish sauce. M. Sc. Thesis. Bangkok, Thailand, Chulalongkorn University.
- Yamanaka, H. 1984. Changes in histamine contents in red meat fish during storage at different temperatures. *Bull. Jpn. Soc. Sci. Fish.* 50: 695-701.
- Yamashita, M., Sakaue, M., Iwata, N., Sugino, H. and Murooka, Y. 1993. Purification and characterization of monoamine oxidase from *Klebsiella aerogenes*. *J. Ferm. Bioeng.* 76: 289-295.
- Yatsunami, K. and Takenaka, T. 2000. Characterizhation of brine proteases as agents of hydrolysis during the ripening of fermentes sardine with rice-bran. *Fish sci.* 66: 569-573.
- Yongsawatdigul, J., Choi, Y. J. and Udomporn, S. 2004. Biogenic amines formation in fish sauce prepared from fresh and temperature abused Indian anchovy (*Stolephorus indicus*). *J. Food Sci.* 69: 312-319.
- Yoshinaka, R., Sato, M., Tsuchiya, N. and Ikeda, S. 1983. Production of fish sauce from sardines by utilization of its visceral enzymes. *Bull. Jpn. Soc. Sci. Fish.* 49: 463-469.