

CONTENTS

Contents	Page
List of tables	xiii
List of figures	xv
Chapter	
1. Introduction	1
Literature review	3
1. Chicken	3
2. Chemical composition of chicken meat	4
3. Quality characteristic of muscle foods	5
4. Factor affecting quality characteristics of muscle	5
5. Effect of pH on meat quality	8
6. Effect of thermal processing on physical and chemical properties of muscle	10
Objectives	13
2. Materials and Methods	14
1. Study on quality characteristics of Thai indigenous chicken muscles as influenced by age and rearing system	14
1.1 Raw materials	14
1.2 Physical analyses	14
1.2.1 Determination of cooking loss	14
1.2.2 Determination of colour	14
1.2.3 Shear force analysis	15
1.2.4 Thermal denaturation	15
1.3 Microstructure of muscle	15
1.4 Chemical analyses	16
1.4.1 Proximate analysis	16
1.4.2 Determination of total collagen content	16
1.4.3 Determination of soluble collagen content	16

CONTENTS (Continued)

	Page
2. Effect of heating under Tom Yum soup condition on chemical composition physical properties of Thai indigenous chicken, spent hen and broiler	17
2.1 Material and preparation	17
2.1.1 Raw materials	17
2.1.2 Tom Yum soup preparation	17
2.1.3 Cooking under Tom Yum soup condition	17
2.1.4 Thermal processing under Tom Yum soup condition	17
2.2 Physical analysis	18
2.2.1 Determination of cooking loss	18
2.2.2 Determination of color	18
2.2.3 Shear force analysis	18
2.2.4 Thermal determination	18
2.3 Microstructure of muscle	18
2.4 Chemical analyses	18
2.4.1 pH	18
2.4.2 Proximate composition of raw muscle	18
2.4.3 Total collagen and soluble collagen content	18
2.5 Sensory evaluation	19
3. Study on quality changes of thermal processed Tom Yum spent hen muscles during storage	19
3.1 Material and preparation	19
3.2 Physical analysis	19
3.2.1 Determination of weight loss	19
3.2.2 Determination of color	19
3.2.3 Shear force analysis	19
3.2.4 Fiber diameter	19
3.3 Chemical analyses	20
3.3.1 pH	20

CONTENTS (Continued)

	Page
3.3.2 Total collagen and soluble collagen content	20
3.4 Sensory evaluation	20
4. Statistical analysis	20
3. Results and Discussion	21
1. Effect of age and rearing systems on quality characteristics and microstructure of Thai indigenous chicken muscles	21
1.1 Proximate compositions	21
1.2 Total collagen and soluble collagen	23
1.3 Color	23
1.4 Shear force	26
1.5 Cooking loss	26
1.6 Thermal denaturation	28
1.7 Microstructure of muscle	29
2. Chemical and physical properties of chicken meat from different breeds and muscle type	37
2.1 Chemical composition	37
2.2 Physical properties	39
2.3 Microstructure of muscle	42
2.4 Effect of heating under Tom Yum soup condition	45
2.5 Effect of heating under different solutions	46
2.6 Effect of thermal processing temperatures on properties of Tom Yum spent hen	51
3. Changes in chemical, physical properties and microstructure of chicken muscles in Tom Yum soup condition during storage	54
3.1 Changes in collagen content	54
3.2 Changes in pH and weight loss of muscle	54
3.3 Changes in fiber diameter	57
3.4 Changes in shear force value	61

CONTENTS (Continued)

	Page
3.5 Changes in color	62
3.6 Changes in sensory	62
4. Conclusions	65
References	67
Appendix	
Additional results	76
Additional figures	82
Vitae	85

LIST OF TABLES

Table		Page
1	Chemical composition of meat from Thai indigenous chicken, broiler and spent hen	4
2	Total collagen and percentage of heat-soluble collagen in cooked, unprocessed and cooked aseptically processed broiler and spent hen chicken breast	12
3	Effect of age and rearing system on proximate compositions of Thai indigenous chicken <i>pectoralis major</i> and <i>biceps femoris</i> muscles	22
4	Effect of age and rearing system on total collagen and soluble collagen of Thai indigenous chicken <i>pectoralis major</i> and <i>biceps femoris</i> muscles	24
5	Effect of age and rearing system on colour of Thai indigenous chicken <i>pectoralis major</i> and <i>biceps femoris</i> muscles	25
6	Effect of age and rearing system on shear force values of Thai indigenous chicken <i>pectoralis major</i> and <i>biceps femoris</i> muscles	27
7	Effect of age and rearing system on cooking loss of Thai indigenous chicken <i>pectoralis major</i> and <i>biceps femoris</i> muscles	28
8	Effect of age and rearing system on transformation temperature and denaturation enthalpy of <i>pectoralis major</i> and <i>biceps femoris</i> muscles	30
9	Effect of age and rearing system on microstructure of Thai indigenous chicken <i>pectoralis major</i> and <i>biceps femoris</i> muscles	32
10	Chemical properties of raw chicken muscles from Thai indigenous chicken, spent hen and broiler	38
11	Physical properties of raw chicken muscles from Thai indigenous chicken, spent hen and broiler	40
12	Transformation temperature of chicken muscle from Thai indigenous chicken, spent hen and broiler	41
13	Physical and chemical properties of Thai indigenous chicken, spent hen and broiler muscles after heating in Tom Yum soup condition heating in difference soup solutions	48

LIST OF TABLES (Continued)

Table		Page
14	Physical and chemical properties of spent hen <i>pectoralis major</i> muscle after heating in difference soup solutions	49
15	Colour of thermal processed spent hen muscles in Tom Yum soup	51
16	Shear force value of thermal processed spent hen muscles in Tom Yum soup	52
17	Total collagen and heat soluble collagen of thermal processed spent hen muscles in Tom Yum soup	52
18	Sensory evaluation of thermal processed spent hen muscles in Tom Yum soup	53
19	Changes in sensory evaluation of thermal processed spent hen <i>pectoralis major</i> and <i>biceps femoris</i> muscles in Tom Yum soup during storage at room temperature (25-30°C)	64
20	F-value of analysis of variance for physical properties of Thai indigenous chicken muscles as influence by age and rearing system	77
21	F-value of analysis of variance for chemical properties of Thai indigenous chicken muscles as influence by age and rearing system	77
22	Changes in total collagen and soluble collagen of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	78
23	Changes in pH of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	78
24	Changes in weight loss of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	79
25	Changes in fiber diameter of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	79
26	Changes in shear force values of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	80
27	Changes in L*, a* and b* values of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	80-81

LIST OF FIGURES

Figure		Page
1	SEM micrographs of transverse sections at differing ages (16, 18 and 20 weeks) of indigenous chicken <i>pectoralis major</i> muscle reared under intensive (I) and extensive (E) systems	33
2	SEM micrographs of transverse sections at different age (16, 18 and 20 weeks) of indigenous chicken <i>biceps femoris</i> muscle reared under intensive (I) and extensive (E) systems	34
3	SEM micrographs of longitudinal sections at differing ages (16, 18 and 20 weeks) of indigenous chicken <i>pectoralis major</i> muscle reared under intensive (I) and extensive (E) systems	35
4	SEM micrographs of longitudinal sections at differing ages (16, 18 and 20 weeks) of indigenous chicken <i>biceps femoris</i> muscle reared under intensive (I) and extensive (E) systems	36
5	SEM micrographs of transverse sections of chicken muscles; <i>pectoralis major</i> muscle of Thai indigenous chicken (A), spent hen (B) and broiler (C); <i>biceps femoris</i> muscle of Thai indigenous chicken (D), spent hen (E) and broiler (F)	43
6	SEM micrographs of longitudinal sections of chicken muscles; <i>pectoralis major</i> muscle of Thai indigenous chicken (A), spent hen (B) and broiler (C); <i>biceps femoris</i> muscle of Thai indigenous chicken (D), spent hen (E) and broiler (F)	44
7	SEM micrograph of transverse sections of <i>pectoralis major</i> muscle after heating under difference soup solutions	50
8	Changes in total collagen and soluble collagen of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	55
9	Changes in pH of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	56
10	Changes in weight loss of thermal processed spent hen muscles in Tom Yum soup during storage at room temperature (25-30°C)	56

LIST OF FIGURES (Continued)

Figure		Page
11	Changes in fiber diameter of thermal processed spent hen <i>pectoralis major</i> and <i>biceps femoris</i> muscles in Tom Yum soup during storage at room temperature (25-30°C)	58
12	SEM micrograph of transverse sections of thermal processed spent hen <i>pectoralis major</i> muscle in Tom Yum soup during storage at room temperature (25-30°C)	59
13	SEM micrograph of transverse sections of thermal processed spent hen <i>biceps femoris</i> muscle in Tom Yum soup during storage at room temperature (25-30°C)	60
14	Changes in shear force values of thermal processed spent hen <i>pectoralis major</i> and <i>biceps femoris</i> muscles in Tom Yum soup during storage at room temperature (25-30°C)	61
15	Changes in L*, a* and b*-values of thermal processed spent hen <i>pectoralis major</i> and <i>biceps femoris</i> muscles in Tom Yum soup during storage at room temperature (25-30°C)	63
16	Thermal processed spent hen <i>pectoralis major</i> (A) and <i>biceps femoris</i> (B) muscles in Tom Yum soup after storage for 6 months at room temperature (25-30°C)	64
17	Intensive production system where chickens were kept in house and provided concentrated feeds: (a) outside of chicken's house and (b) inside of chicken's house which divided into 12 blocks (23 chickens/block)	83
18	Extensive production system where allowed to scavenge on natural food around the homestead (18 meter squares) during the day and supplemented with concentrated feeds in the evening when they came back to roost and sheltered at night; (a) outside of chicken's house and; (b) group of chicken scavenged on nature food in their homestead (12 blocks x 23 chickens)	84