

CONTENTS

| | Page |
|---|------|
| Contents..... | viii |
| Lists of Tables..... | xiv |
| List of Figures..... | xv |
| Chapter | |
| 1. Introduction and Literature Review | |
| Introduction..... | 1 |
| Literature Review..... | 3 |
| Cephalopod..... | 3 |
| Tissue structure of cephalopod..... | 5 |
| Chemical compositions of cephalopod..... | 5 |
| Protein..... | 7 |
| Lipid..... | 8 |
| Freezing and frozen storage..... | 10 |
| Lipid oxidation and its effect on protein denaturation and discoloration | |
| of muscle food during processing and storage..... | 14 |
| Mechanism of lipid oxidation..... | 15 |
| The important factors affecting lipid oxidation..... | 17 |
| Effect of lipid oxidation on protein denaturation..... | 19 |
| Physicochemical changes of protein..... | 20 |
| Changes in functional properties of protein..... | 22 |
| Changes in color..... | 23 |
| Lipid oxidation of microsome and liposome systems..... | 26 |
| Objectives of study..... | 29 |
| 2. Chemical composition and thermal property of cuttlefish (<i>Sepia pharaonis</i>) | |
| muscle | |
| Abstract..... | 30 |

| | |
|----------------------------|----|
| Introduction..... | 30 |
| Materials and Methods..... | 31 |
| Chemicals..... | 31 |

CONTENTS (Continued)

| | Page | Sample |
|--|--------|--------|
| preparation..... | 32 | |
| Proximate analysis and determination of collagen content..... | 32 | |
| Determination of lipid composition and fatty acid profile..... | 32 | |
| Fractionation of muscle protein..... | 33 | |
| Thermal transition measurement..... | 33 | |
| Determination of trace mineral content..... | 34 | |
| Statistical analysis..... | 34 | |
| Results and Discussion..... | 34 | |
| The proximate analysis and collagen content..... | 34 | |
| Lipid composition and fatty acid profile of cuttlefish..... | 36 | |
| Composition of muscle proteins..... | 37 | |
| Trace mineral content in cuttlefish..... | 41 | |
| Thermal transition of cuttlefish muscle..... | 42 | |
| Conclusion..... | 43 | |

3. The effect of metal ions on lipid oxidation, color and chemical properties of cuttlefish

(*Sepia pharaonis*) subjected to multiple freeze-thaw cycles

| | |
|--|----|
| Abstract..... | 44 |
| Introduction..... | 45 |
| Materials and Methods..... | 46 |
| Chemicals..... | 46 |
| Materials..... | 46 |
| Sample preparation..... | 47 |
| Preparation of natural actomyosin..... | 47 |

| | |
|--|----|
| Thiobarbituric acid reactive substances (TBARS)..... | 47 |
| Color..... | 48 |
| Surface hydrophobicity..... | 48 |
| Total sulfhydryl content..... | 48 |
| Disulfide bond content..... | 49 |
| Sodium dodecyl sulfate-polyacrylamide gel electrophoresis..... | 49 |

CONTENTS (Continued)

| | Page |
|--|------|
| ATPase activity..... | 49 |
| Protein solubility..... | 50 |
| Statistical analysis..... | 50 |
| Results and Discussion..... | 50 |
| Lipid oxidation..... | 50 |
| Discoloration..... | 51 |
| Surface hydrophobicity..... | 52 |
| Sulfhydryl and disulfide bond content..... | 55 |
| SDS-PAGE..... | 56 |
| ATPase activity | 57 |
| Solubility..... | 58 |
| Conclusion..... | 61 |

4. Yellow discoloration of cuttlefish (*Sepia pharaonis*) liposome system as influenced by lipid oxidation

| | |
|--|----|
| Abstract..... | 62 |
| Introduction..... | 62 |
| Materials and Methods..... | 63 |
| Chemicals..... | 63 |
| Cuttlefish collection and preparation..... | 64 |
| Preparation of cuttlefish liposome..... | 64 |

| | |
|---|----|
| Lipid oxidation of cuttlefish liposome systems..... | 64 |
| Determination of lipid oxidation..... | 65 |
| Determination of free amine groups..... | 65 |
| Measurement of phospholipid pyrrolization..... | 65 |
| Color measurement..... | 66 |
| Statistical analysis..... | 66 |
| Results and Discussion..... | 66 |
| Effect of temperature and time on the changes of cuttlefish liposome system..... | 66 |

CONTENTS (Continued)

| | Page |
|--|------|
| Effect of FeCl ₃ and ascorbic acid on the changes of cuttlefish liposome system..... | 69 |
| Effect of NaCl on the changes of cuttlefish liposome system..... | 70 |
| Conclusion..... | 72 |
| 5. Lipid oxidation in microsomal fraction of squid muscle (<i>Loligo peali</i>) | |
| Abstract..... | 73 |
| Introduction..... | 73 |
| Materials and Methods..... | 74 |
| Chemicals..... | 74 |
| Materials..... | 75 |
| Preparation of squid microsomal fraction..... | 75 |
| Lipid oxidation in microsomal fraction of squid muscle..... | 76 |
| Preparation of liposomes from squid microsomal lipids..... | 76 |
| Measurement of lipid oxidation..... | 77 |
| Color measurement..... | 77 |
| Statistics..... | 77 |
| Results and Discussion..... | 78 |

| | |
|---|-----|
| Enzymatic and nonenzymatic lipid oxidation in microsomal fraction of squid muscle..... | 78 |
| The effect of nonenzymatic lipid oxidation on discoloration of squid muscle microsomes and microsomal lipids..... | 82 |
| Conclusion..... | 85 |
| 6. Development of yellow pigmentation in squid (<i>Loligo pealei</i>) as a result of lipid oxidation | |
| Abstract..... | 86 |
| Introduction..... | 86 |
| Materials and Methods..... | 88 |
| CONTENTS (Continued) | |
| Page | |
| Materials..... | 88 |
| Preparation of squid microsomal fraction, liposomes and salt-soluble proteins..... | 88 |
| Lipid oxidation of liposomes or the microsomal fraction of squid muscle | 89 |
| Measurement of chemical and physical alterations in the microsomes, liposomes and salt-soluble myofibrillar proteins..... | 90 |
| Statistics..... | 92 |
| Results and Discussion..... | 92 |
| Conclusion..... | 102 |
| 7. The effect of antioxidants on the quality changes of cuttlefish (<i>Sepia pharaonis</i>) muscle during frozen storage | |
| Abstract..... | 103 |
| Introduction..... | 103 |
| Materials and Methods..... | 105 |

| | |
|--|-----|
| Chemicals..... | 105 |
| Cuttlefish collection and preparation..... | 105 |
| Treatment of cuttlefish with various antioxidants..... | 106 |
| Thawing of frozen cuttlefish..... | 106 |
| Preparation of natural actomyosin..... | 106 |
| Surface hydrophobicity..... | 107 |
| Total sulfhydryl content..... | 107 |
| Disulfide bond content..... | 107 |
| Sodium dodecyl sulfate-polyacrylamide gel electrophoresis..... | 108 |
| Thiobarbituric acid reactive substances (TBARS)..... | 108 |
| Protein solubility..... | 108 |
| Thaw drip..... | 109 |
| Color..... | 109 |
| Statistical analysis..... | 109 |

CONTENTS (Continued)

| | Page |
|--|------|
| Results and Discussion..... | 110 |
| Changes in lipid oxidation..... | 110 |
| Changes in color..... | 111 |
| Changes in physicochemical properties of natural actomyosin..... | 112 |
| Changes in solubility..... | 116 |
| Changes in thaw drip..... | 117 |
| Conclusion..... | 119 |

8. Summary and Future works

| | |
|-------------------|-----|
| Summary..... | 120 |
| Future works..... | 121 |
| References..... | 122 |
| Vitae..... | 138 |

LIST OF TABLES

| Table | Page |
|--|------|
| 1. Proximate composition of mantle muscle of squid and cuttlefish..... | 7 |
| 2. Lipid composition of squid mantle..... | 9 |
| 3. Chemical compositions of cuttlefish muscle..... | 35 |
| 4. Lipid composition in cuttlefish muscle..... | 36 |
| 5. Fatty acid profile of cuttlefish lipid..... | 38 |
| 6. Protein and non-protein nitrogen components in cuttlefish muscle..... | 39 |
| 7. Trace mineral content of cuttlefish muscle..... | 41 |
| 8. Peak transition temperature (T_{\max}) and denaturation enthalpies (ΔH) of cuttlefish muscle protein..... | 43 |
| 9. Changes in TBARS of cuttlefish paste with different metal ions at concentration 5 ppm and 25 ppm during multiple freeze-thaw cycles..... | 53 |
| 10. Effect of sodium chloride on thiobarbituric acid reactive substances (TBARS) formation, b*-value and pyrrole compounds of cuttlefish liposome systems..... | 72 |
| 11. Interaction between <i>trans</i> -2-hexenal and salt-soluble myofibrillar proteins..... | 96 |

LIST OF FIGURES

| Figure | Page |
|---|-------------|
| 1. Cuttlefish (Pharaoh Cuttlefish: <i>Sepia pharaonis</i>)..... | 4 |
| 2. Squid (Long finned squid: <i>Loligo pealei</i>)..... | 4 |
| 3. The structure of cephalopod..... | 6 |
| 4. Factors affecting protein denaturation and aggregation during frozen storage... .. | 12 |
| 5. The overall mechanism of lipid oxidation and its effect on the changes of quality in food during processing and storage..... | 15 |
| 6. Production of reactive carbonyls during phosphatidylethanolamine oxidation and the later formation of pyrrolized phospholipids by amino-carbonyl reactions..... | 25 |
| 7. A proposed pathway of the enzymatic lipid oxidation in microsomal fraction of flounder muscle..... | 27 |
| 8. Formation of lipid membranes, micelles, emulsions and liposomes from amphipathic lipids..... | 28 |
| 9. SDS-PAGE pattern of cuttlefish whole muscle and fractions..... | 41 |
| 10. Changes in L*, a* and *b-values of cuttlefish paste containing different metal ions (25 ppm) and subjected to multiple freeze-thaw cycles..... | 54 |
| 11. Changes in surface hydrophobicity (S_0 ANS) of cuttlefish proteins exposed to different metal ions (25 ppm) and subjected to multiple freeze-thaw cycles..... | 55 |
| 12. Changes in sulphydryl content and disulfide bond content of cuttlefish proteins exposed to different metal ions (25ppm) and subjected to multiple freeze-thaw cycles..... | 56 |
| 13. SDS-PAGE pattern of cuttlefish mince with subjected to different metal-ions (25 ppm) and subjected three freeze-thaw cycles..... | 57 |
| 14. Changes in ATPase activity of cuttlefish proteins exposed to different metal | |

| | |
|---|----|
| ions (25 ppm) and subjected to multiple freeze-thaw cycles..... | 59 |
|---|----|

LIST OF FIGURES (Continued)

| Figure | Page |
|---|-------------|
| 15. Changes in Ca ²⁺ -sensitivity of cuttlefish proteins exposed to different metal ions (25 ppm) and subjected to multiple freeze-thaw cycles..... | 60 |
| 16. Changes in protein solubility of cuttlefish paste exposed to different metal ions (25 ppm) and subjected to multiple freeze-thaw cycles..... | 60 |
| 17. Formation of thiobarbituric acid reactive substances (TBARS) in liposomes made from cuttlefish lipids (5 mg lipid/mL) in the presence of 50 μM FeCl ₃ and 200 μM ascorbic acid during incubation at different temperatures for 24 h..... | 67 |
| 18. Changes in L*, a* and b*-values in liposomes made from cuttlefish lipids (5 mg lipid/mL) in the presence of 50 μM FeCl ₃ and 200 μM ascorbic acid during incubation at different temperatures for 24 h..... | 68 |
| 19. Thiobarbituric acid reactive substances (TBARS) formation and yellow pigment formation (b*-value), the loss of amine groups and the formation of pyrrole compounds in liposomes made from cuttlefish lipids (5 mg lipid/ml) in the presence of 50 μM FeCl ₃ and 200 μM ascorbic acid during incubation at 37 °C for different times..... | 69 |
| 20. Effect of FeCl ₃ and ascorbic acid concentrations on the formation of Thiobarbituric acid reactive substances (TBARS), yellow pigments formation (b*-value), loss of amine groups and pyrrolization in liposomes made from cuttlefish lipids (5 mg lipid/mL) during incubation at different temperatures for 24 h..... | 71 |
| 21. Formation of thiobarbituric acid reactive substances (TBARS) in squid muscle microsomes (2 mg microsomal protein/mL) in the presence or absence of 200 μM ADP, 50 μM FeCl ₃ , 200 μM NADH, 200 μM | |

| | |
|--|----|
| NADPH during incubation at 4 °C and 37 °C..... | 79 |
| 22. Formation of thiobarbituric acid reactive substances (TBARS) in squid muscle microsomes (2 mg microsomal protein/mL) in the presence of 200 µM ADP, 50 µM FeCl ₃ and 200 µM ascorbic acid during incubation at 4, 25 and 37 °C..... | 80 |

LIST OF FIGURES (Continued)

| Figure | Page |
|---|------|
| 23. Effect of iron concentration on thiobarbituric acid reactive substances (TBARS) formation in squid muscle microsomes in the presence of 200 µM ADP and 200 µM ascorbic acid during incubation at 37 °C..... | 81 |
| 24. Effect of ascorbic acid concentration on thiobarbituric acid reactive substances (TBARS) formation in squid muscle microsomes in the presence of 200 µM ADP and 50 µM FeCl ₃ during incubation at 37 °C..... | 81 |
| 25. Formation of thiobarbituric acid reactive substances (TBARS) in squid muscle microsomes (approximately 0.9 mg microsomal lipid/mL; microsomal system) or lipids isolated from muscle microsomes (0.2 mg lipid/mL; liposome system) in the presence of 200 µM ADP, 50 µM FeCl ₃ and 200 µM ascorbic acid during incubation at 37 °C..... | 83 |
| 26. Changes in L*, a* and b*-values in squid muscle microsomes (2 mg microsomal protein/mL; microsomal system) or lipids isolated from muscle microsomes (0.2 mg lipid/mL; liposome system) in the presence of 200 µM ADP, 50 µM FeCl ₃ and 200 µM ascorbic acid during incubation at 37 °C..... | 84 |
| 27. Relationship between development of lipid oxidation (measured as TBARS) with yellow pigment formation (b*-value), loss of amines and formation of pyrrole compounds in squid muscle microsomes. Oxidation was promoted with 200 µM ascorbic acid and 50 µM FeCl ₃ in the presence of 5 mg of microsomal protein/mL of 0.12 M KCl/25 mM potassium phosphate | |

| | |
|---|----|
| buffer, pH 7.2, at 37 °C..... | 94 |
| 28. Changes in sulfhydryls and solubility of 10 parts of squid muscle salt-soluble proteins (5 mg/mL 0.6 M KCl/25 mM potassium phosphate buffer, pH 7.2) after 8 h of exposure at 25 °C to 1 part of 0.12 M KCl/25 mM potassium phosphate buffer, pH 7.2, 1 part of 200 µM ascorbic acid and 50 µM FeCl ₃ , 1 part of squid muscle microsomes (5 mg of microsomal protein/mL of 0.12 M KCl/25 mM potassium phosphate buffer, pH 7.2) oxidized by | |

LIST OF FIGURES (Continued)

| Figure | Page |
|--|------|
| 200 µM ascorbic acid and 50 µM FeCl ₃ , and 2 parts of squid muscle microsomes (5 mg of microsomal protein/mL of 0.12 M KCl/25 mM potassium phosphate buffer, pH 7.2) oxidized by 200 µM ascorbic acid and 50 µM FeCl ₃ | 96 |
| 29. Formation of thiobarbituric acid reactive substances (TBARS) and yellow pigments (b* value), loss of amines and formation of pyrrole compounds in liposomes (5 mg of lipid/mL of 0.12 M KCl/25 mM potassium phosphate buffer at pH 7.2) prepared from squid microsomal lipids and egg yolk lecithin during oxidation by 200 µM ascorbic acid and 50 µM FeCl ₃ at 37 °C..... | 97 |
| 30. Changes in b*-value in egg yolk lecithin liposomes (5 mg of lipid/mL of 0.12 M KCl/25 mM potassium phosphate buffer at pH 7.2) upon exposure to various concentrations of <i>trans</i> -2-hexenal or <i>trans</i> -2-nonenal at 37 °C for 15 h..... | 98 |
| 31. Changes in b*-value, amines and pyrrole compounds in egg yolk lecithin liposomes (5 mg of lipid/mL of 0.12 M KCl/25 mM potassium phosphate buffer at pH 7.2) exposed to different aldehydic lipid oxidation products (5.0 mM) at 37 °C for 15 h..... | 99 |

| | |
|--|-----|
| 32. Proposed model for the impact of physical location of aldehydic lipid oxidation products on their reactivity with phospholipid amines..... | 101 |
| 33. Thiobarbituric acid reactive substances (TBARS) in cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C for 16 weeks..... | 111 |
| 34. L*, a* and b*-values of cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C for 16 weeks. | 113 |

LIST OF FIGURES (Continued)

| Figure | Page |
|---|-------------|
| 35. Surface hydrophobicity (S _o ANS) of natural actomyosin extracted from cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C for 16 weeks | 115 |
| 36. Sulfhydryl content and disulfide bond content of natural actomyosin extracted from cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C for 16 weeks..... | 115 |
| 37. SDS-PAGE pattern of cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C for 16 weeks. | 116 |
| 38. Protein solubility of cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C for 16 weeks. | 118 |
| 39. Thaw drip of cuttlefish treated with 5% NaCl and 0.3% H ₂ O ₂ , followed by soaking in different antioxidant solutions during frozen storage at -18 °C | |

| | |
|-------------------|-----|
| for 16 weeks..... | 119 |
|-------------------|-----|