

เอกสารอ้างอิง

- จันทร์เพ็ญ อิสลาม. 2544. การคัดเลือกแบคทีเรียที่ผลิตสารลดแรงตึงผิวชีวภาพและสภาวะที่เหมาะสมต่อการผลิต. วิทยานิพนธ์วิทยาศาสตร์มหาบัณฑิต. มหาวิทยาลัยสงขลานครินทร์.
- ชัยวัฒน์ เจนวนิชย์. 2527. บทนำ : แนวคิดพื้นฐาน. ใน เคมีโพลิเมอร์พื้นฐาน. หน้า 1-80. โอดีเยนส์โตร์. กรุงเทพฯ.
- ดวงพร คันธิชิต. 2537. อนุกรรมวิธานของแบคทีเรียและปฏิกิริยา. โอดีเยนส์โตร์. กรุงเทพฯ.
- มั่นศิน ตันตูลเวศ์. 2542. วิศวกรรมการประปา เล่ม 1. สำนักพิมพ์แห่งจุฬาลงกรณ์มหาวิทยาลัย.
- ศิริพร หมายหล้า. 2544. การคัดเลือกแบคทีเรียนอ่อนหุ่นภูมิสูงที่ผลิตพอลิเมอร์ กำจัดเนื้อชนิดและคุณสมบัติของพอลิเมอร์. วิทยานิพนธ์วิทยาศาสตร์มหาบัณฑิต. มหาวิทยาลัยสงขลานครินทร์.
- สุวรรณा เหลืองชลธาร. 2542. เคมีอนินทรีย์ทางเภสัชศาสตร์. สำนักพิมพ์แห่งจุฬาลงกรณ์มหาวิทยาลัย.
- Abdel-El-Haleem, D. 2003. *Acinetobacter* : environmental and biotechnological applications. African J. Biotechnol. 2(4) : 71-74.
- Arino, S., Marchal, R. and Vandecasteele, J.P. 1996. Identification and production of a rhamnolipidic biosurfactant by a *Pseudomonas* species. Appl. Microbiol. Biotechnol. 45 : 162-168.
- Arjen, J., Olsthoorn, J. and Duine, J.A. 1996. Production characterization and reconstitution of recombinant quinoprotein glucose dehydrogenase (soluble type; EC1.1.99.17) apoenzyme of *Acinetobacter calcoaceticus*. Arch. Biochem. Biophys. 336(1) : 42-48.
- Bach, H., Berdichevsky, Y. and Gutnick, D. 2003. An exocellular protein from the oil-dregration microbe *Acinetobacter venetianus* RAG-1 enhances the emulsifying activity of the polymerric bioemulsifier emulsan. Appl. Environ. Microbiol. 69(5) : 2608-2615.

- Benson, F., Tenoux, I., Hourdou, L.M. and Michel, G. 1992. Synthesis of β -hydroxy fatty acids and β -amino fatty acids by the strains of *Bacillus subtilis* producing iturinic antibiotics. *Biochem. Biophys. Acta.* 1123 : 51-58.
- Bognolo, G. 1999. Biosurfactants as emulsifying agents for hydrocarbons. *Colloid. Surf. A.* 152 : 41-52.
- Borel, C., Welti, D.H., Fernandez, I. and Colmenares, M. 1993. Didranin, an antimicrobial and 15-lipoxygenase inhibitor from the moss *Didranum scoporium*. *J. Natural. Product.* 56 : 1071-1077.
- Cameotra, S.S. and Makkar, R.S. 1998. Synthesis of biosurfactants in extreme conditions. *Appl. Microbiol. Biotechnol.* 50 : 520-529.
- Campbell, D. and White, J.R. 1989. Molecular weight determination. In. polymer characterization : Physical techniques. Chapman and Hall. London.
- Carreno-Gomez, B. and Duncan, R. 1997. Evaluation of the biological properties of soluble chitosan and chitosan microspheres. *Int. J. Pharm.* 148 : 231-240.
- Carrillo, P.G., Merdaraz, C., Pitta-Alvarez, S.I. and Giulietti, A.M. 1996. Isolation and selection of biosurfactant-producing bacteria. *World J. Microbiol. Biotechnol.* 13 : 137-139.
- Chaplin, M.F. and Kennedy, J.F. 1986. Carbohydrate Analysis : A Practical Approach. Washington DC : IRL Press.
- Chen, S.J., Cheng, C.Y. and Chen, T.L. 1998. Production of an alkaline lipase by *Acinetobacter radioresistens*. *J. Ferment. Bioeng.* 86(3). 308-312.
- Choi, H.J. and Kunioka, M. 1995. Preparation conditions and swelling equilibria of hydrogel prepared by γ -irradiation from microbial poly (γ -glutamic acid). *Radiat. Phys. Chem.* 46(2) : 175-179.
- Choi, H.J., Yang, R. and Kunioka, M. 1995. Synthesis and characterization of pH-sensitive and biodegradable hydrogels prepared by γ -irradiation using microbial poly (γ -glutamic acid) and poly (ϵ -lysine). *J. applied. Polymer. science.* 58 : 807-814.

- Collins, E.A., Bares, J. and Billmeyer, F.W. 1973. Experiments in Polymer Science. New York : John Wiley & Sons, Inc.
- Damodaran, S. 1996. Amino acid, peptides and proteins. In Food chemistry. p. 321-430. O.R. Fennema (Ed). 3th ed. New York. Marcel Dekker, Inc.
- Dawson, R.M.C., Elliott, D.C., Elliott, W.F. and Jones, K.M. 1986. Method for the detection of biochemical compounds on paper and thin-layer chromatograms with some notes on separation. In Data for Biochemical Research. 3rd ed. p. 453-502. The Bath Press. Avon.
- Deleu, M., Razafindralambo, H., Popineau, Y., Jaoques, P., Thonart, P. and Paquot, M. 1999. Interfacial and emulsifying properties of lipopeptides from *Bacillus subtilis*. A : Physiochem. Engineer. Aspects. 152 : 3-10.
- Dermlim, W. 1999. Screening for polymer-producing bacteria from seafood activated sludge and bioflocculant characterization. Master of Science Thesis in biotechnology. Prince of Songkla University.
- Dermlim, W., Prasertsan, P. and Doelle, H. 1999. Screening and characterization of bioflocculant produced by isolated *Klebsiella* sp. Appl. Microbiol. Biotechnol. 52 : 698-703.
- Desai, J.D. and Banat, I.M. 1997. Microbial production of surfactant and their commercial potential. Microbiol. Mol. Biol. Rev. 61(1) : 47-64.
- Deziel, E., Lepine, F., Milot, S. and Villemur, R. 2000. Mass spectrometry monitoring of rhamnolipids from a growing culture of *Pseudomonas aeruginosa* strain 57RP. Biochim. Biophys. Acta. 1485 : 145-152.
- Dlamini, A.M. and Peiris, P.S. 1997. Production of exopolysaccharide by *Pseudomonas* sp. ATCC 31461 (*Pseudomonas elodea*) using whey as fermentation substrate. Appl. Microbiol. Biotechnol. 47 : 52-57.
- Dodgson, K.S. and Price, R.G. 1962. A note on the determination of the ester sulphate content of sulphated polysaccharides. Biochem. J. 84 : 106-110.
- Drouin, C.M. and Cooper, D.G. 1992. Biosurfactant and aqueous two phase fermentation. Biotechnol. Bioeng. 40 : 86-90.

- Facklam, R. and Elliott, J.A. 1995. Identification, classification and clinical relevance of catalase-nagative, gram-positivite, excluding the *Streptococci* and *Enterococci*. *Clin. Microbiol. Rev.* 8 : 479-495.
- Fiechter, A. 1992. Biosurfactants : moving towards industrial application. *TIBTECH.* 10 : 208-217.
- Fujita, M., Ike, M., Tachibana, S., Kitada, G., Kim, S.M. and Inoue. Z. 2000. Characterizatic of a bioflocculant produced by *Citrobacter* sp. TKF04 from acetic and propionic acids. *J. Biosci. Bioeng.* 89(1) : 40-46.
- Goto, A. and Kunioka, M. 1992. Biosynthesis and Hydrolysis of poly-(γ -glutamic acid) from *Bacillus subtilis* IFO3335. *Biosci. Biotech. Biochem.* 56(7) : 1031-1035.
- Gutnick, D.L. and Bach, H. 2000. Engineering bacterial biopolymers for the biosorption of heavy metals ; new products and novel formulations. *Appl. Microbiol. Biotechnol.* 54 : 451-460.
- Han, S.J., Back, J.H., Yoon, M.Y., Shin, P.K, Cheong, C.S., Sung, M.H., Hong, S.P. Chung, I.Y. and Han, Y.S. 2003. Expression and characterization of a novel enantioselective lipase from *Acinetobacter* species SY-01. *Biochimie.* 85(5) : 501-510.
- He, N., Li, Y., Chen, J. and Lun, S.Y. 2002. Idetification of a novel bioflocculant from a newly isolated *Corynebacterium glutamicum*. *Biochem. Eng. J.* 11 : 137-148.
- He, N., Li, Y. and Chen, J. 2004. Production of a novel polygalacturonic acid biosurfactant REA-11 by *Corynebacterium glutamicum*. *Biores. Technol.* 94(1) : 99-105.
- <http://www.genialab.de/inventory/alginate.htm>
- Huy, N.Q., Jin, S., Amada, K., Haruki, M., Huu, N.B., Hang, D.T., Camna, D.T., Imanaka, T., Morikawa, M. and Kanaya, S. 1999. Characterization of petroleum-degrading bacteria from oil-contaminated sites in Vietnam. *J. Biosci. Bioeng.* 88 : 100-102.

- Ito, Y., Tanaka, T., Ohmachi, T. and Asada, Y. 1996. Glutamic acid independent production of poly-(γ -glutamic acid) from *Bacillus subtilis* TAM-4. Biosci. Biotech. Biochem. 60(8) : 1239-1242.
- Kim, H.S., Yoon, B.D., Lee, C.H., Suh, H.H., Oh, H.M., Katsuragi, T and Tani, Y. 1997. Production and properties of a lipopeptide biosurfactant from *Bacillus subtilis* C9. J. Ferment. Bioeng. 84(1) : 41-46.
- Kim, H.S., Yoon, B.D., Choung, D.H., Oh, H.M., Katsuragi, T. and Tani, Y. 1999. Characterization of a biosurfactant, mannosylerythritol lipid produced from *Candida* sp. SY16. Appl. Microbiol. Biotechnol. 52 : 713-721.
- Kim, H.S., Jeon, J.W., Lee, H.W., Park, Y.I., Seo, W.T., Oh, H.M., Katsuragi, T., Tani, Y. and Yoon, B.D. 2002. Extracellular production of a glycolipid biosurfactant, mannosylerythritol lipid, from *Candida antarctica*. Biotechnol. Lett. 24 : 225-229.
- Koma, D., Hasumi, F., Yamamoto, E., Ohta, T., Chung, S.Y. and Kubo, M. 2001. Biodegradation of long-chain n-paraffins from waste oil of car engine by *Acinetobacter* sp. J. Biosci. Bioeng. 91(1) : 94-96.
- Kosaric, N. 1993. Biosurfactants production properties application. Marcel Dekker, Inc. New York.
- Kosaric, N. 1996. Biosurfactants. In Biotechnology. Vol 6. 2nd ed. by Rehm, H.J. and Reed, G. Weinheim ([Germany](#)) : Wiley-VCH.
- Krieg, N.R. and Holt, J.G. 1984. Bergey's Manual of Systematic Bacteriology. Vol.1. Williams and Wilkins. Baltimore.
- Kubota, H., Matsunobu, T., Uotani, K., Takebe, H., Satoh, A., Tanaka, T. and Taniguchi, M. 1993. Production of poly-(γ -glutamic acid) from *Bacillus subtilis* F-2-01. Biosci. Biotech. Biochem. 56(7) : 1212-1213.
- Kunioka, M. and Choi, H.J. 1995. Properties of biodegradable hydrogels prepared by γ -irradiation of microbial poly (ϵ -lysine) aqueous solution. J. Appl. Polymer. Scien. 58 : 801-806.
- Kunioka, M. 1997. Biosynthesis and chemical reaction of poly (amino acid)s from microorganisms. Appl. Microbiol. Biotechnol. 47 : 469-475.

- Kurane, R. and Matsuyama, H. 1994. Production of a bioflocculant by mixed culture. *Biosci. Biotech. Biochem.* 58(9) : 1589-1594.
- Kurane, R. and Nohata, Y. 1994. A new water-absorbing polysaccharide from *Alcaligenes latus*. *Biosci. Biotech. Biochem.* 58(2) : 235-238.
- Kurane, R., Toeda, K., Takeda, K. and Suzuki, T. 1986. Culture conditions for production of microbial flocculant by *Rhodococcus erythropolis*. *Agric. Biol. Chem.* 50(9) : 2309-2313.
- Kurane, R., Hatakeyama, S. and Tsugeno, H. 1991. Correlation between flocculation production and morphological changes in *Rhodococcus erythropolis* S-1. *J. Ferment. Bioeng.* 72 : 498-500.
- Kurane, R., Hatamochi, K., Kakuno, T., Kiyohara, M., Hirano, M. and Taniguchi, Y. 1994a. Production of a bioflocculant by *Rhodococcus erythropolis* S-1 grown on alcohols. *Biosci. Biotech.Biochem.* 58(2) : 428-429.
- Kurane, R., Hatamochi, K., Kakuno, T., Kiyohara, M., Kawaguchi, K., Mizuno, Y., Hirano, M. and Taniguchi, Y. 1994b. Purification and charcterization of lipid bioflocculant produced by *Rhodococcus erythropolis*. *Biosci. Biotech.Biochem.* 58(11) : 1977-1982.
- Lang, S. 2002. Biological amphiphiles (microbial biosurfactants). *Curr. Opin. Colloid. Interface. Science.* 7 : 12-20.
- Lang, S and Wullbrandt, D. 1999. Rhamnose lipids-biosynthesis, microbial production and application potential. *Appl. Microbiol. Biotechnol.* 51 : 22-32.
- Lee, S.H., Lee, S.O., Jang, K.L. and Lee, T.H. 1995. Microbial flocculant from *Arouadendron* sp. TS-49. *Biotech. Lett.* 17(1) : 95-100.
- Lin, S.C., Lin, K.G., Lo, C.C. and Lin, Y.M. 1998. Enhanced biosurfactant production by a *Bacillus licheniformis* mutant. *Enzym. Micro. Technol.* 23 : 267-273.
- Lowry, O.H., Rosenberg, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with folin phenol reagent. *J. Biol. Chem.* 193 : 265-275.
- MacFaddin, J.F. 1980. Biochemical Tests for Identification of Medical Bacteria. 2nd ed. Baltimore : Williams & Wilkins.

- Makkar, R.S. and Cameotra, S.S. 1998. Production of biosurfactant at mesophilic and thermophilic conditions by a strain of *Bacillus subtilis*. *J. Ind. Microbiol. Biotechnol.* 20 : 48-52.
- Magaritis, A and Pace, G.W. 1985. Microbial polysaccharides *In* Comprehensive Biotechnology. : Vol 3, The Practice of Biotechnology : Current Commodity Products. (Blanch, H.W., Drew, S. and Wang, D.I.C. eds.). p : 1006-1014. Oxford : Pergamon Press, Ltd.
- McIlroy**, R.J. 1967. Introduction to carbohydrate chemistry. London. Butterworths & Co. Publishers Limited.
- Morikawa, M., Daido. H., Takao, T., Murate, S., Shimonishi, Y. and Imanaka, T. 1993. A new lipopeptide biosurfactant from *Arthrobacter* sp. strain MIS38. *J. Bacteriol.* 175(20) : 6459-6466.
- Morikawa, M., Hirata, Y. and Imanaka, T. 2000. A study on the structure-function relationship of lipopeptide biosurfactants. *Biochim. Biophys. Acta.* 1488 : 211-218.
- Morin, A. 1998. Screening of polysaccharide-producing microorganisms, factors influencing the production, and recovery of microbial polysaccharides. *In* Polysaccharides : Structure diversity and functional versatility. (Dumitriu, S. ed) p. 275-296. New York : Marcel Dekker.
- Nakata, K. and Kurane, R. 1999. Production of an extracellular polysaccharide bioflocculant by *Klebsiella pneumoniae*. *Biosci. Biotechnol. Biochem.* 63(12) : 2064-2068.
- Nam, J.S., Kwon, G.S., Lee, S.O., Hwang, J.S., Lee, J.D., Yoon, B.D. and Lee, T.H. 1996. Bioflocculant produced by *Aspergillus* sp. JS-42. *Biosci. Biotech.Biochem.* 60 (2) : 325-327.
- Nampoothiri, K.M., Singhania, R.R., Sabarinath, C. and Pandey, A. 2003. Fermentation production of gellan using *Sphingomonas paucimobilis*. *Process Biochem.* 38 : 1513-1519.

- Navon-venezia, S., Zosim, Z., Gottlieh, A., legmann, R., Carmeli, S., Ron, E.Z. and Rosenberg. E. 1995. Alasan, a new bioemulsifier from *Acinetobacter radioresistens*. Appl. Environ. Microbiol. 61 : 3240-3244.
- Odian, G. 1993. Introduction In. Principle of polymerization. 3rd ed : A wiley-interscience publication. United States of America.
- Ostroha, J., Pong, M., Lowman, A. and Dan. N. 2004. Controlling the collapse/swelling transition in charged hydrogel. Biomaterials. 25(18) : 4345-4353.
- Ogawa, Y., Yamaguchi, F., Yuasa, K. and Tahara, Y. 1997. Efficient production of γ -polyglutamic acid by *Bacillus subtilis* (natto) in jar fermenters. Biosci. Biotech.Biochem. 61(10) : 1684-1687.
- Passeri, A., Lang, S. and Wagner, F. and Wray, V. 1991. Marine Biosurfactants, II. Production and characterization of an anionic trehalose tetraester from the marine bacterium *Arthrobacter* sp. EK1. Z. Naturforsch. 46c : 204-209.
- Prommachan, O. 2002. [Production and application of biosurfactant from *Bacillus MUV4*.](#) Master of Science Thesis in biotechnology. Prince of Songkla University.
- Ron, E.Z. and Rosenberg, E. 2001. Natural roles of biosurfactants. Environ. Microbiol. 3(4) : 229-236.
- Rosenberg, E. and Ron, E.Z. 1997. Bioemulsans : microbial polymeric emulsifiers. Curr. Opin. Biotechnol. 9 : 313-316.
- Rosenberg, E. and Ron, E.Z. 1999. High- and low-molecular-mass microbiol surfactants. Appl. Microbiol. Biotechnol. 52 : 154-162.
- Saito, K., Sato, S., Shimoji, H., Iefuji, H. and Tadenuma, M. 1990. Flocculation mechanism of *Hansenula anomala* J224. Agric. Biol. Chem. 54 : 1425-1432.
- Salehizadeh, H. and Shojaosadati, S.A. 2001. Extracellular biopolymeric flocculants recent trends and biotechnological importance. Biotech. Advance. 19 : 371-385.
- Salehizadeh, H., Vossoughi, M. and Alemzadeh, I. 2000. Some investigations on bioflocculant producing bacteria. Biochem. Eng. J. 5 : 39-44.

- Shih, I.L. and Van, Y.T. 2001. The production of poly-(γ -glutamic acid) from microorganisms and its various applications. *Bioresour. Techno.* 79 : 207-225.
- Shih, I.L., Van, Y.T., Yeh, L.C., Lin, H.C. and Chang, Y.N. 2001. Production of a biopolymer flocculant from *Bacillus licheniformis* and its flocculation properties. *Bioresour. Techno.* 78 : 267-272.
- Shima, S., Fukuhara, Y. and Sakai, H. 1982. Inactivation of bacteriophages by ϵ -poly-L-lysine produced by *Streptomyces*. *Agric. Biol. Chem.* 46(7) : 1917-1919.
- Shima, S., Matsuoka, H., Iwamoto, T. and Sakai, H. 1984. Antimicrobial action of ϵ -poly-L-Lysine. *J. Antibiotic.* 37(11) : 1449-1455.
- Shimada, A., Nakata, H. and Nakamura, I. 1997. Acidic exopolysaccharide produced by *Enterobacter* sp. *J. Ferment. Bioeng.* 84(2) : 113-118.
- Shimofuruva, H., koide, A., Shirota, K., Tsuji, T., Nakamura, M. and Suzuki, J. 1996. The production of flocculanting substance(s) by *Streptomyces griseus*. *Biosci. Biotech. Biochem.* 61 : 498-500.
- Singh, P. and Cameotra, S.S. 2004. Potential applications of microbial surfactants in biomedical science. *Trends. Biotechnol.* 22(3) : 142-151.
- Sutherland, I.W. 1998. Novel and established applications of microbial polysaccharides. *TIBTECH.* 16 : 41-46.
- Shu, C.H. and Lung, M.Y. 2004. Effect of pH on the production and molecular weight distribution of exopolysaccharide by *Antrodia camphorata* in batch cultures. *Process. Biochem.* 39(8) : 931-937.
- Sousa, M., Teixeira, J. and Mota, M. 1992. Difference in flocculation mechanism of *Kluyveromyces marxianus* and *Saccharomyces cerevisiae*. *Biotechnol. Lett.* 14 : 213-218.
- Tahzibi, A., Kamal, F and Assadi, M.M. 2004. Improved production of rhamnolipids by a *Pseudomonas aeruginosa* mutant. *Iranian. Biomedical. J.* 8(1) : 25-31.
- Takeda, M., Kurane, R., Koizumi, J. and Nakamura, I. 1991. A protein bioflocculant produced by *Rhodococcus erythropolis*. *Agric. Biol. Chem.* 55(10) : 2663-2664.

- Takeda, M., Koizumi, J.J., Matsuoka, H. and Hikuma, M. 1992. Factors affecting the activity of a protein biosurfactant produced by *Norcadia amarea*. *J. Ferment. Bioeng.* 74(6) : 408-409.
- Tallon, R., Bressollier, P. and Urdaci, M.C. 2003. Isolation and characterization of two exopolysaccharides produced by *Lactobacillus plantarum* EP56. *Research. Microbiol.* 154 : 705-712.
- Tong, Z., Zhe, L. and Huai-lan, Z. 1999. Microbial flocculant and its application in environmental protection. *J. Environ. Sci.* 11 : 1-12.
- Toren, A., Novon-Venezia, S., Ron, E.Z. and Rosenberg, E. 2001. Emulsifying activity of purified alasan proteins from *Acinetobacter radioresistens*. *Appl. Environ. Microbiol.* 67 : 1102-1106.
- Tuleva, B.K., Ivanov, G.R. and Christova, N.E. 2002. Biosurfactant production by a new *Pseudomonas putida* strain. *Z. Naturforsch.* 57C : 356-360.
- Vashuk, E.V., Vorobieva, E.V., Basalyga, I.I. and Krutko, N.P. 2001. Water-absorbing properties of hydrogels based on polymeric complexes. *Mat. Res. Innovat.* 4 : 350-352.
- Weiner, R.M. 1997. Biopolymers from marine prokaryotes. *TIBTECH.* 15 : 390-394.
- Wu, J. and Ju, L.K. 1998. Extracellular particles of polymeric material formed in n-hexadecane fermentation by *Pseudomonas aeruginosa*. *J. Biotechnol.* 59 : 193-202.
- Yakimov, M.M., Timmis, K.N., Wray, V. and Fredrickson, H.L. 1995. Characterization of a new lipopeptide surfactant produced by thermotolerant and halotolerant subsurface *Bacillus licheniformis* BAS50. *Appl. Environ. Microbiol.* 61(5) : 1706-1713.
- Yokoi, H., Natsuda, O., Hirose, J., Hayashi, S. and Takasaki, Y. 1995. Characteristics of a biopolymer flocculant produced by *Bacillus* sp. PY-90. *J. Ferment. Bioeng.* 79(4) : 378-380.

- Yokoi, H., Arima, T., Hirose, J., Hayashi, S and Takasaki, Y. 1996. Flocculation properties of poly (γ -glutamic acid) produced by *Bacillus subtilis*. *J. Ferment. Bioeng.* 82(1) : 84-87.
- Yokoi, H., Yoshida, T., Mori, S., Hirose, J., Hayashi, S. and Takasaki, Y. 1997. Biopolymer flocculant produced by an *Enterobacter* sp. *Biotechnol. Lett.* 19(6) : 569-573.
- Yoon, S.H., Do, J.H., Lee, S.Y. and Chang, H.N. 2000. Production of poly- γ -glutamic acid by fed-batch culture of *Bacillus licheniformis*. *Biotechnol. Lett.* 22 : 585-588.
- Youssef, N.H., Duncan, K.E., Nagle, D.P., Savage, K.N., Knapp, R.M. and McInerney, M.J. 2004. Composition of methods to detect biosurfactant production by diverse microorganism. *J. Microbial. Method.* 56(3) : 339-347.