

เอกสารอ้างอิง

กองทันตสาธารณสุข กรมอนามัย กระทรวงสาธารณสุข. 2545. การสำรวจสภาวะทันตสุขภาพแห่งชาติ ครั้งที่ 5 ปีพ.ศ.2543-2544. กรุงเทพมหานคร.

Al-Khateeb, S. ; ten Cate, J.M. ; Angmar-Månsson, B. ; de Josselin de Jong, E. ; Sundström, G. ; Exterkate, R.A.M. and Oliveby, A. 1997a. "Quantification of formation and remineralization of artificial enamel lesions with a new portable fluorescence device", Adv Den Res. 11 (1997a), 502-506.

Al-Khateeb, S. ; Oliveby, A. ; de Josselin de Jong, E. ; and Angmar-Månsson, B. 1997b. "Laser fluorescence quantification of remineralisation in situ of incipient enamel lesions: Influence of fluoride supplements", Caries Res. 31 (1997b), 132-140.

Alwas-Danowska, H.M. ; Plasschaert, A.J. ; Suliborski, S. and Verdonschot, E.H. 2002. "Reliability and validity issues of laser fluorescence measurements in occlusal caries diagnosis", J Dent. 30 (2002), 129-234.

Allison, P.J. and Schwartz, S. 2003. "Interproximal contact points and proximal caries in posterior primary teeth", Pediatr Dent. 25 (2003), 334-340.

Ando, M. ; Hall, A.F. ; Eckert, G.J. ; Schemehorn, B.R. ; Analoui, M. and Stookey, G.K. 1997. "Relative ability of laser fluorescence techniques to quantitate early mineral loss in vitro", Caries Res. 31 (1997), 125-131.

Ando, M. ; de Josselin de Jong, E. ; Fontana, M. ; Stookey, G.K. and Zero, D.T. 2000. "Evaluation of the effect of dehydration on demineralized enamel with quantitative light-induced fluorescence (QLF)", Caries Res. 34 (2000), 326. (Abstract)

- Ando, M. ; Van der Veen, M.H. ; Schemehorn, B.R. and Stookey, G.K. 2001. "Comparative study to quantify demineralized enamel in deciduous and permanent teeth using laser- and light-induced fluorescence techniques", Caries Res. 35 (2001), 464-470.
- Angmar-Månsson, B. ; Al-Khateeb, S. and Tranæus, S. 1996. "Monitoring the caries process: Optical methods for clinical diagnosis and quantification of enamel caries", Eur J Oral Sci. 104 (1996), 480-485.
- Angmar-Månsson, B. and ten Bosch, J.J. 1987. "Optical methods for the detection and quantification of caries", Adv Dent Res. 1 (1987), 14-20.
- Angmar-Månsson, B. and ten Bosch, J.J. 1993. "Advances in methods for diagnosing coronal caries - a review", Adv Dent Res. 7 (1993), 70-79.
- Anttonen, V. ; Seppä, L. and Hausen, H. 2003. "Clinical study of the use of the laser fluorescence device DIAGNOdent for detection of occlusal caries in children", Caries Res. 37 (2003), 17-23.
- Araujo, F.B. ; Araujo, D.R. ; Santos, C.K. and Souza, M.A.L. 1996. "Diagnosis of approximal caries in primary teeth: Radiographic versus clinical examination using tooth separation", Am J Dent. 9 (1996), 54-56.
- Attrill, D.C. and Ashley, P.F. 2001. "Occlusal caries detection in primary teeth: a comparison of DIAGNOdent with conventional methods", Br Dent J. 190 (2001), 440-443.

- Bader, J.D. ; Shugars, D.A. and Bonito, A.J. 2001. "Systematic reviews of selected dental caries diagnostic and management methods", J Dent Educ. 65 (2001), 960-968.
- Bamzahim, M. ; Shi, X-Q. and Angmar-Månsson, B. 2002. "Occlusal caries detection and quantification by DIAGNOdent and electronic caries monitor: in vitro comparison", Acta Odontol Scand. 60 (2002), 360-364.
- Banerjee, A. and Boyde, A. 1998. "Autofluorescence and mineral content of carious dentine: Scanning optical and backscattered electron microscopic studies", Caries Res. 32 (1998), 219-226.
- Baseren, N.M. and Gokalp, S. 2003. "Validity of a laser fluorescence system (DIAGNOdent) for detection of occlusal caries in third molars: an in vitro study", J Oral Rehabil. 30 (2003), 1190-1194.
- Berman, D.S. and Slack, G.L. 1973. "Caries progression and activity in approximal tooth surfaces", Br Dent J. 134 (1973), 51-57.
- Bjelkhagen, H. ; Sundström, F. ; Angmar-Månsson, B. and Rydén, H. 1982. "Early detection of enamel caries by the luminescence excited by visible laser light", Swed Dent J. 6 (1982), 1-7.
- Chadwick, B.L. and Dummer, P.H.M. 1998. "Factors affecting the diagnostic quality of bitewing radiographs: a review", Br Dent J. 184 (1998), 80-84.
- Côrtes, D.F. ; Ellwood, R.P. and Ekstrand, K.R. 2003. "An in vitro comparison of a combined FOTI/visual examination of occlusal caries with other caries diagnostic

methods and the effect of stain on their diagnostic performance”, Caries Res. 37 (2003), 8-16.

Costa, A.M. ; Yamaguti, P.M. ; De Paula, L.M. and Bezerra, A.C. 2002. “In vitro study of laser diode 655 nm diagnosis of occlusal caries”, ASDC J Dent Child. 69 (2002), 249-253.

Dean, J.A. ; Barton, D.H. ; Vahedi, I. and Hatcher, E.A. 1997. “Progression of interproximal caries in the primary dentition”, J Clin Pediatr Dent. 22 (1997), 59-62.

Downer, M.C. 1975. “Concurrent validity of an epidemiological diagnostic system for caries with the histological appearance of extracted teeth as validating criterion”, Caries Res. 9 (1975), 231-246.

Eggertsson, H. ; Analoui, M. ; Van der Veen, M.H. ; González-cabezas, C. ; Eckert, G.J. and Stookey, G.K. 1999. “Detection of early interproximal caries in vitro using laser fluorescence, dye-enhanced laser fluorescence and direct visual examination”, Caries Res. 33 (1999), 227-233.

El-Housseiny, A.A. and Jamjoum, H. 2001. “Evaluation of visual, explorer, and a laser device for detection of early occlusal caries”, J Clin Pediatr Dent. 26 (2001), 41-48.

Emami, Z. ; Al-Khateeb, S. ; de Josselin de Jong, E. ; Sundström, F. ; Trollsås, K. and Angmar-Månsson, B. 1996. “Mineral loss in incipient caries lesions quantified with laser fluorescence and longitudinal microradiography: a methodologic study”, Acta Odontol Scand. 54 (1996), 8-13.

- Espelid, I. and Tviet, A.B. 1986. "Clinical and radiographic assessment of approximal carious lesions", Acta Odontol Scand. 44 (1986), 31-37.
- Featherstone, J.D.B. 2000. "Caries detection and prevention with laser energy", Dent Clin North Am. 44 (2000), 955-969.
- Featherstone, J.D.B. and Mellberg, J.R. 1981. "Relative rates of progress of artificial carious lesions in bovine,ovine and human enamel", Caries Res. 15 (1981), 109-114.
- Feldens, C.A. ; Tovo, M.F. ; Kramer, P.F. ; Feldens, E.G. ; Ferreira, S.H. and Finkler, M. 2003. "An in vitro study of the correlation between clinical and radiographic examinations of proximal carious lesions in primary molars", J Clin Pediatr Dent. 27 (2003), 143-148.
- Forgie, A.H. ; Pine, C.M. and Pitts, N.B. 1999. "Laser fluorescence detection of approximal caries : Initial in vitro results", J Dent Res. 78 (1999), 1044. (Abstract)
- Francescut, P. and Lussi, A. 2003. "Correlation between fissure discoloration, Diagnodent measurements, and caries depth: an in vitro study", Pediatr Dent. 25 (2003), 559-564.
- Friedman, J. and Marcus, M.I. 1970. "Transillumination of the oral cavity with use of fibre optics", J Am Dent Assoc. 80 (1970), 801-809.
- Hafström-Björkman, U. ; Sundström, F. ; de Josselin de Jong, E. ; Oliveby, A. and Angmar-Månsson, B. 1992. "Comparison of laser fluorescence and longitudinal microradiography for quantitative assessment of in vitro enamel caries", Caries Res. 26 (1992), 241-247.

- Hall, A.F. ; DeSchepper, E. ; Ando, M. and Stookey, G.K. 1997. "In vitro studies of laser fluorescence for detection and quantification of mineral loss from dental caries", Adv Dent Res. 11 (1997), 507-514.
- Heinrich-Weltzien, R. ; Kühnisch, J. ; Oehme, T. ; Ziehe, A. ; Stösser, L. and Garcia-Godoy, F. 2003. "Comparison of different DIAGNOdent cut-off limits for in vivo detection of occlusal caries", Oper Dent. 28 (2003), 672-680.
- Hibst, R. and Paulus, R. 2000. "Molecular basis of red excited caries fluorescence", Caries Res. 34 (2000), 323. (Abstract)
- Hintze, H. ; Winzel, A. and Larsen, M.J. 1995. "Stereomicroscopy, film radiography, microradiography and naked-eye inspection of tooth sections as validation for occlusal caries diagnosis", Caries Res. 29 (1995), 359-363.
- Hintze, H. ; Winzel, A. ; Danielsen, B. and Nyvad, B. 1998. "Reliability of visual examination, fibre-optic transillumination, and bite-wing radiography, and reproducibility of direct visual examination following tooth separation for the identification of cavitated carious lesions in contacting approximal surfaces", Caries Res. 32 (1998), 204-209.
- Horowitz, H.S. 2000. "Decision-making for national programs of community fluoride use", Community Dent Oral Epidemiol. 28 (2000), 321-329.
- de Josselin de Jong, E. ; Sundström, F. ; Emami, Z. and Angmar-Månsson, B. 1992. "Quantification of mineral loss in initial caries lesions on natural enamel surfaces with laser fluorescence", Caries Res. 26 (1992), 216. (Abstract)

- Longbottom, C. and Pitts, N.B. 1992. "A comparison of five caries diagnostic methods employing an in vitro simulation and histological validation", Caries Res. 26 (1992), 216. (Abstract)
- Longbottom, C. ; Wagner, M. ; Pitts, N.B. and Lussi, A. 1999. "Simulated in vivo comparison of bitewing radiography and DIAGNOdent for approximal caries detection", Caries Res. 33 (1999), 298. (Abstract)
- Lussi, A. ; Imwinkelried, S. ; Pitts, N.B. ; Longbottom, C. and Reich, E. 1999. "Performance and reproducibility of a laser fluorescence system for detection of occlusal caries in vitro", Caries Res. 33 (1999), 261-266.
- Lussi, A. ; Megert, B. ; Longbottom, C. ; Reich, E. and Francescut, P. 2001. "Clinical performance of a laser fluorescence device for detection of occlusal caries lesions", Eur J Oral Sci. 109 (2001), 14-19.
- Lussi, A. and Francescut, P. 2003. "Performance of conventional and new methods for the detection of occlusal caries in deciduous teeth", Caries Res. 37 (2003), 2-7.
- McComb, D. and Tam, L.E. 2001a. "Diagnosis of occlusal caries: Part I conventional methods", J Can Dent Assoc. 67 (2001a), 454-457.
- McComb, D. and Tam, L.E. 2001b. "Diagnosis of occlusal caries: Part II recent diagnostic technologies", J Can Dent Assoc. 67 (2001b), 459-463.
- Mendes, F.M. ; Hissadomi, M. and Imparato, J.C.P. 2004. "Effects of drying time and the presence of plaque on the in vitro performance of laser fluorescence in occlusal caries of primary teeth", Caries Res. 38 (2004), 104-108.

- Mileman, P.A. ; van der Weele, L.T. 1990. "Accuracy in radiographic diagnosis: Dutch practitioners and dental caries", J Dent. 18 (1990), 130-136.
- Mitropoulos, C.M. 1985a. "The use of fibre-optic transillumination in the diagnosis of posterior approximal caries in clinical trials", Caries Res. 19 (1985a), 379-384.
- Mitropoulos, C.M. 1985b. "A comparison of fibre-optic transillumination with bitewing radiographs", Br Dent J. 159 (1985b), 21-23.
- Mortimer, K.V. 1970. "The relationship of deciduous enamel structure to dental disease", Caries Res. 4 (1970), 206-223.
- Murray, J.J. and Majid, Z.A. 1978. "The prevalence and progression of approximal caries in the deciduous dentition in British children", Br Dent J. 145 (1978), 161-164.
- Nyvad, B. ; Machiulskiene, V. and Baelum, V. 1999. "Reliability of a new caries diagnostic system differentiating between active and inactive caries lesions", Caries Res. 33 (1999), 252-260.
- Peers, A. ; Hill, F.J. ; Mitropoulos, C.M. and Holloway, P.J. 1993. "Validity and reproducibility of clinical examination, fibre-optic transillumination, and bite-wing radiology for the diagnosis of small approximal carious lesions: An in vitro study", Caries Res. 27 (1993), 307-311.
- Peltola, J. and Wolf, J. 1981. "Fibre optics transillumination in caries diagnosis", Proc Finn Dent Soc. 77 (1981), 240-244.

Petersson, L.G. ; Arthursson, L. ; Östberg, C. ; Jönsson, G. and Gleerup, A. 1991.

“Caries-inhibiting effects of different modes of duraphat varnish reapplication: a 3-year radiographic study”, Caries Res. 25 (1991), 70-73.

Peyron, M. ; Matsson, L. and Birkhed, D. 1992. “Progression of approximal caries in primary molars and the effect of duraphat treatment”, Scand J Dent Res. 100 (1992), 314-318.

Pinelli, C. ; Campos Serra, M. and de Castro Monteiro Loffredo, L. 2002. “Validity and reproducibility of a laser fluorescence system for detecting the activity of white-spot lesions on free smooth surfaces in vivo”, Caries Res. 36 (2002), 19-24.

Pitts, N.B. 1984. “Systems for grading approximal carious lesions and overlaps diagnosed from bitewing radiographs: proposals for future standardization”, Community Dent Oral Epidemiol. 12 (1984), 114-122.

Pitts, N.B. 1996. “The use of bitewing radiographs in the management of dental caries: scientific and practical considerations”, Dentomaxillofac Radiol. 25 (1996), 5-16.

Pitts, N.B. and Rimmer, P.A. 1992. “An in vivo comparison of radiographic and directly assessed clinical caries status of posterior approximal surfaces in primary and permanent teeth”, Caries Res. 26 (1992), 146-152.

Purdell-Lewis, D.J. and Pot, T. 1974. “A comparison of radiographic and fibre-optic diagnoses of approximal caries lesions”, J Dent. 2 (1974), 143-148.

Ricketts, D.N.J. ; Whaites, E.J. ; Kidd, E.A.M. ; Brown, J.E. and Wilson, R.F. 1997. “An evaluation of the diagnostic yield from bitewing radiographs of small approximal

and occlusal carious lesions in a low prevalence sample in vitro using different film types and speeds”, Br Dent J. 182 (1997), 51-58.

Russel, M. and Pitts, N.B. 1993. “Radiovisiographic diagnosis of dental caries: Initial comparison of basic mode videoprints with bitewing radiography”, Caries Res. 27 (1993), 65-70.

Shi, X-Q. ; Welander, U. and Angmar-Månsson, B. 2000. “Occlusal caries detection with KaVo DIAGNOdent and radiography: an in vitro comparison”, Caries Res. 34 (2000), 151-158.

Shi, X-Q. ; Tranaeus, S. and Angmar-Månsson, B. 2001a. “Comparison of QLF and DIAGNOdent for quantification of smooth surface caries”, Caries Res. 35 (2001a), 21-26.

Shi, X-Q. ; Tranaeus, S. and Angmar-Månsson, B. 2001b. “Validation of DIAGNOdent for quantification of smooth-surface caries: an in vitro study”, Acta Odontol Scand. 59 (2001b), 74-78.

Shellis, R.P. 1984. “Relationship between human enamel structure and the formation of caries-like lesions”, Archs Oral Biol. 29 (1984), 975-981.

Sidi, A.D. and Naylor, M.N. 1988. “A comparison of bitewing radiography and interdental transillumination as adjuncts to the clinical identification of approximal caries in posterior teeth”, Br Dent J. 164 (1988), 15-18.

Stephen, K.W. ; Russel, J.I. ; Creanor, S.L. and Burchell, C.K. 1987. “Comparison of fibre optic transillumination with clinical and radiographic caries diagnosis”, Community Dent Oral Epidemiol. 15 (1987), 90-94.

- Stookey, G.K. and González-Cabezas, C. 2001. "Emerging methods of caries diagnosis", J Dent Educ. 65 (2001), 1001-1006.
- Vaarkamp, J. ; ten Bosch, J.J. ; Verdonshot, E.H. and Bronkhorst, E.M. 2000. "The real performance of bitewing radiography and fiber-optic transillumination in approximal caries diagnosis", J Dent Res. 79 (2000), 1747-1751.
- Vanderas, A.P. ; Manetas, C. ; Koulatzidou, M. and Papagiannoulis, L. 2003. "Progression of proximal caries in the mixed dentition: a 4-year prospective study", Pediatr Dent. 25 (2003), 229-234.
- Wagner, M. ; Longbottom, C. and Pitts, N.B. 1999. "An in vitro comparison of a laser device with bitewing radiography for approximal caries detection", Caries Res. 33 (1999), 298. (Abstract)
- Waggoner, W.F. and Ashton, J.J. 1989. "Predictability of cavitation based upon radiographic appearance: comparison of two film types", Quintessence Int. 20 (1989), 55-60.
- Welsh, G.A. ; Hall, A.F. ; Hannah, A.J. and Foye, R.H. 2000. "Variation in DIAGNOdent measurements of stained artificial caries lesions", Caries Res. 34 (2000), 324. (Abstract)
- Wenzel, A. 2004. "Dental caries", In Oral radiology: Principles and interpretation, p.297-313. White, S.C. and Pharoah, M.J. 5th ed. St.Louis, Missouri : Mosby Inc.
- Wilson, P.R. and Beynon, A.D. 1989. "Mineralization differences between human deciduous and permanent enamel measured by quantitative microradiography", Archs Oral Biol. 34 (1989), 85-88.