

เอกสารอ้างอิง

- ชื่อทิพย์ อาทิตย์มาศ. 2531. พวรรณไม่น้ำของไทย. ภาควิชาชีววิทยา คณะวิทยาศาสตร์ มหาวิทยาลัยสงขลานครินทร์. 104-109.
- นิสิตา บำรุงวงศ์. ยาภัคฆามะเริง. 2542. ภาควิชาเภสัชวิทยา คณะวิทยาศาสตร์ มหาวิทยาลัยสงขลานครินทร์. 2-6.
- พยองค์ วนิเกียรติ, ยุพิน สังวรินทร์, สุกีนันท์ อัญเชิญ และ นพมาศ วงศ์วิทยาเดชา. 2539. นาสซีวิทยา. ภาควิชาเภสัชวิทยา คณะวิทยาศาสตร์ มหาวิทยาลัยมหิดล. 611-618.
- ศรีจันทร์ พรจิราศิลป์, จุฑามณี สุทธิสีสังข์ และ รัชนี เมฆมนี. 2542. นาสซีวิทยา. ภาควิชาเภสัชวิทยา คณะเภสัชศาสตร์ มหาวิทยาลัยมหิดล. 420-421.
- Alves, A.P., Mulloy, B., Diniz, J.A. and Mouroa, P.A.S. 1997. Sulfated polysaccharide from the egg jelly layer are species-specific inducers of acrosome reaction in sperms of sea urchins. *J.Biol. Chem.* 272: 6965-6971.
- Bilan, M.I., Grachev, A.A., Ustuzhanina, N.E., Shashkov, A.S., Nifantiev, N.E. and Usov, A.I. 2002. Structure of fucoidan from the brown seaweed *Fucus evanescens*. *Carbohydrate Research*. 337: 719-730.
- Bilan, M.I., Grachev, A.A., Ustuzhanina, N.E., Shashkov, A.S., Nifantiev, N.E. and Usov, A.I. 2004. A high regular fraction of a fucoidan from the brown seaweed *Fucus distichus* L. *Carbohydrate Research*. 339: 511-517.
- Cappuccino, J.P. 1986. Agar plate sensitivity method. In *Microbial A Laboratory Manual*, The Benjamin Cummings Publishing Company Inc, California, p. 260-261.

- Church, F.R., Meade, J.B., Treanor, R.E. and Whinna, H.C. 1989. Antithrombin activity of fucoidan. *J. Biol. Chem.* 264: 3618-3623.
- Chizhov, A.O., Dell, A., Morris, H.R., Haslam, S.M., McDowell, R.A., Shashkov, A.S. and Nifant, N.E. 1999. A study of fucoidan from the brown seaweed *Chorda filum*. *Carbohydrate Research*. 320: 108-119.
- Chevolet, L., Mulloy, B., Ratiskol, J., Foucault, A. and Collice-Jouault, S. 2001. A disaccharide repeat unit is the major structure in fucoidans from two species of brown algae. *Carbohydrate Research*. 330: 529-535.
- Chotigeat, W., Tongsupa S., Supamataya K. and Phongdara, A. 2004. Effect of fucoidan on disease resistance of black tiger shrimp. *Aquaculture*. 233: 23-30.
- Collicec, S., Vidal, C.B. and Jozefonviez, J. 1994. A low molecular weight fucoidan fraction from the brown seaweed *Pelvetia canaliculata*. *Phytochemistry*. 35: 697-700.
- Dische, Z. 1947. A new specific colour reaction of hexuronic acid. *J. Biol. Chem.* 167: 189-198.
- Dodgson, K.S. 1961. Determination of Inorganic sulphate in studies on the enzymic and non-enzymic hydrolysis of carbohydrate and other sulphate esters. *Biochem. J.* 78: 312-319.
- Doner, L.W. and Whistler, R.L. 1973. Fucoidan. *In Industrial gums polysaccharide and their derivatives*. Academic prees Inc. New York. p115-120.
- Duarte, M.E.R., Cardoso, M.A., Noseda, M.D. Cerezo, A.S. 2001. Structural studies on fucoidans from the brown seaweed *Sargassum stenophyllum*. *Carbohydrate research*. 333: 281-293.

- Fabregas, J., Garcia, D., Fernandes, A.M., Rocha, A.I., Gomez, P., Escribano, J.M., Otero, A. and Coll, J.M. 1999. In vitro inhibition of Replication of Haemorrhagic Septicaemia Virus (VHSV) and African Swine Fever Virus (ASFV) by extracts from marine microalgae. *Antiviral Res.* 44: 67-73.
- Farndale, R.W., Buttle, D.J., Barrett, A.J. 1986. Improved quantitative and discrimination of sulfated glycosaminoglycans by use of dimethylmethylene blue. *Biochimica et Biophysica Acta.* 883: 173-177.
- Freshney, R.I. 1994. Culture of animal cells: A manual of basic technique. 3rd ed. Department of medical oncology CRC beatson laboratories university of glasgow.Wiley-Liss Inc. New York. p. 296-298.
- Fuiihara , M., Iizima, N., Yamamoto, I. And Nagumo, T. 1984. Purification and chemical and physical characterization of an antitumour polysaccharide from the brown seaweed *Sargassum fulvellum* . *Carbohydrate research.* 125: 97-106.
- Hellio, C., De La Broise, D., Dufosse, L., Le Gal, Y., Bourgougnon, N. 2001. Inhibition of marine bacteria by extracts of macroalgae: potential use for environmentally friendly antifouling paints. *Marine Environmental research.* 52: 231-247.
- Haroun-Bouhedja, F., Ellouali M., Sinquin C. and Boisson-Vidal, C. 2000. Relationship between sulfate groups and biological activities of fucans. *Thrombosis Research.* 100: 453-459.
- Itoh, H., Noda, H., Amano,H., Zhuaug, C., Mizuno, T. and Ito, H.1993. Antitumor activity and immunological properties of marine agal polysaccharides, especially fucoidan, prepared from *Sargassum thunbergii* of Phaeophyceae. *Anticancer Research.* 13: 2045 – 2052.

- Koyanagi, S., Tanigawa, N., Nakagawa, H., Soeda, S and Shimeno, H. 2002. Oversulfation of fucoidan enhances its anti-angiogenic and antitumor activities. *Biochemical Pharmacology.* 7478: 1-7.
- Lennette, EH., Spaulding, E.H. and Truant, J.P. 1974. Dilution test procedures. In *Manual of clinical microbiology.* American society for microbiology, Washington D.C., p. 410-417.
- Lowry, O.H., Rosebrough, N.T., Farr, A.L. and Randall, R.T. 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193: 265-275.
- Mabeau, S., Kloareg, B. and Joseleau, J.P. 1990. Fractionation and analysis of fucans from brown algae. *Phytochemistry.* 29: 2441-2445.
- Maimone, M. and Tollefson, D.M. 1990. Structure of a dermatan sulfate hexasaccharide that binds to heparin cofactor II with high affinity. *J. Biol. Chem.* 265: 18263 -18271.
- Matsuda, M., Shigeta, S. and Okutani, K. 1999. Antiviral activities of marine *Pseudomonas* polysaccharides and their oversulfated derivatives. *Mar.Biotechnol.* 1: 68-73.
- Minix, R. and Doctor, V.M. 1997. Interaction of fucoidan with proteases and inhibitors of coagulation and fibrinolysis. *Thrombosis research.* 87: 419-429.
- Mourao, P.A.S., Boisson-Vidal, C., Tapon-Bretaudiere, J., Drouet, B., Bros, A. and Fischer, A.M. 2001. Inactivation of Thrombin by a fucosylated chondroitin sulfate from echinoderm. *Thrombosis Research.* 102: 167-176.

- Mulloy, B. Mourao, P.A.S. and Gray, E. 2000. Structure/function studies of anticoagulant sulfated polysaccharide using NMR. Journal of biotechnology. 77: 123-135.
- Patankar, M.S., Oehninger, S., Barnett, T., Williams, R.L. and Clark, G.F. 1993. A revised structure for fucoidan may explain some of its biological activity. J. Biol. Chem. 268: 21770-21776.
- Pereira, M.S., Mulloy, B. and Mourao, P.A.S. 1999. Structure and anticoagulant activity of sulfated fucans. J. Biol. Chem. 274: 7656-7667.
- Preeprame, S., Hayashi, K., Lee, J.B., Sankava, U. and Hayashi, T. 2001. A novel antivirally active fucan sulfated derived from an- edible brown alga. *Sargassum horneri*. Chemical. Pharm. Bull. 44: 484-485.
- Ponce, N.M.A., Pujol, C.A., Damonte, E.B., Flores, M.L. and Stortz, C.A. 2003. Fucoidans from the brown seaweed *Adenocystis utricularis*: extraction methods, antiviral activity and structure studies. Carbohydrate Research. 338: 153-165.
- Rao, P.S. and Perekh, K.S. 1981. Antibacterial activity of Indian seaweed extracts. BOT. MAR. 24: 577-582.
- Religa, P., Kazi, M., Thyberg, J., Gaciong, Z., Swedenborg, J. and Hedin, U. 2000. Fucoidan inhibits smooth muscle cell proliferation and reduces mitogen activated protein kinase activity. European Journal of Vascular and Endovascular Surgery. 20: 419 – 426.
- Rocha, H.A.O., Franco, C.R.C., Trindade, E.S., Veiga, S.S., Leite, E.L., Dietrich, D.P. and Nader, H.B. 2001. A fucan from the brown seaweed *Spatoglossum schroederi* inhibits Chinese hamster ovary cell adhesion

to several extracellular matrix proteins. Brazilian Journal of Medicine and Biological Research. 34: 621-626.

Takahashi, Y., Uehara, K., Watanobe, R., Okumura, T., Yamashita, T., Omura, H., Yomo, T., Kanemitsu, A., Kawano, T., Narasaka, H., Suzuki, N. and Itami, T. 1998. Efficacy of oral administration shrimp in Japan. In Flegal TW (ed) Advances in Shrimp Biotechnology . National Center for Genetic Engineering and Biotechnology, Bangkok, p. 171-173.

Winzler, R.J. 1971. Determination of serum glycoproteins. Methods Biochem. Analyt. 2: 294-276.

Wityrouw, R. and De Clercq, E. 1997. Sulfated polysaccharide extracted from sea algae as potential antiviral drugs. Gen. Pharmacol. 29: 497-511.

Yoon, S.J., Pereira, M.S., Pavao, M.S.G., Hwang, J.K., Pyun, Y.R. and Mourao, P.A.S. 2002. The medicinal plant *Porana volubilis* contains polysaccharides with anticoagulant activity mediated by heparin cofactor II. Thrombosis research. 106: 51-58.

Zierer, M.S. and Mourao, P.A.S. 2000. A wide diversity of sulfated polysaccharides are synthesized by different species of marine sponges. Carbohydrate research. 328: 209-216.

<http://www.ku.ac.th/Agr Info/thaifish/aqplant/AQPT112>

http://www.takara-bio.co.jp/english/news_e/1996/06/17.htm

http://www.takara-bio.co.jp/english/news_e/2000/03/14.htm

http://www.takara.co.jp/english/bio_e/news_e/1996/06/17.htm

<http://www.slim4 health.com/limu/fucoidan2.html>

http://www.some-limu.info/fucoidan_research.htm