

เอกสารอ้างอิง

- Anuntagool, N., Intachote, P., Wuthiekanun, V., White, N.J. and Sirisinha, S., 1998. Lipopolysaccharide from non-virulent Ara⁺ *Burkholderia pseudomallei* isolates is immunologically indistinguishable from lipopolysaccharide from virulent Ara⁻ clinical isolates. *Clin. Diag. Lab. Immunol.* 5, 225–229.
- Anuntagool, A., Intachote, P., Naigowit, P. and Sirisinha, S. 1996. Rapid antigen detection assay for identification of *Burkholderia (Pseudomonas) pseudomallei* infection. *J. Clin. Microbiol.* 34:975-976.
- Anuntagool, N., Rugdech, P. and Sirisinha, S. 1993. Identification of specific antigens of *Pseudomonas pseudomallei* and evaluation of their efficacies for diagnosis of melioidosis. *J. Clin. Microbiol.* 31:1232-1236.
- Appassakij, H., Silpapojakul, K.R., Wansit, R. and Pornpatkul, M. 1990. Diagnostic value of the indirect hemagglutination test for melioidosis in an endemic area. *Am. J. Trop. Med. Hyg.* 42:248-253.
- Blocker, A., Komoriya, K. and Aizawa, S. 2003. Type III secretion systems and bacterial flagella: insights into their function from structural similarities. *Proc. Natl. Acad. Sci. USA.* 100:3027-30.
- Brett, P.T., DeShazer, D. and Woods, D.E. 1998. *Burkholderia thailandensis* sp. nov., a *Burkholder pseudomallei*-like species. *Int. J. Syst. Bacteriol.* 48:317-320.
- Brown, A.E., Dance, D.A., Suputtamongkol, Y., et al. 1991. Immune cell activation in melioidosis: increased serum levels of interferon-gamma and soluble interleukin-2 receptors without change in soluble CD8 protein. *J. Infect. Dis.* 163:1145-1148.
- Bryan, L.E., Wong, S., Woods, D.E., Dance, D.A.B. and Chaowagul, W. 1994. Passive protection of diabetic rat with antisera specific for the polysaccharide portion of the lipopolysaccharide isolated from *Pseudomonas pseudomallei*. *Can. J. Infect. Dis.* 5:170-178.
- Chaowagul, W., N. J. White, D. A. Dance, Y. Wattanagoon, P. Naigowit, T. M. Davis, S. Looareesuwan, and N. Pitakwatchara. 1989. Melioidosis: a major cause of community-

- acquired septicemia in northeastern Thailand. *J. Infect. Dis.* 159:890–899.
- Cheng, A. C. and Currie, B. J. 2005. Melioidosis: Epidemiology, Pathophysiology, and Management. *Clin Microbiol Rev.* 18(2): 383–416.
- Cheng, A. C., Peacock, S. J., Limmathurotsakul, D., Wongsuvan, G., Chierakul, W., Amornchai, P., Getcharat, N., Chaowagul, W., White, N. J., Day, N. P. and Wuthiekanun, V. 2006. Prospective evaluation of a rapid immunochromogenic cassette test for the diagnosis of melioidosis in northeast Thailand. *Trans R Soc Trop Med Hyg* 100:64-7.
- Dian, C., Eshaghi, S., Urbig, T., McSweeney, S., Heijbel, A., Salbert, G. and Birse, D. 2002. Strategies for the purification and on-column cleavage of glutathione-S-transferase fusion target proteins. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 769(1):133-44.
- Dharakul, T., Songsivilai, S., Anuntagool, N., Chaowagul, W., Wongbunnate, S., Intachote, P. and Sirisinha, S. 1997. Diagnostic value of an antibody enzyme-linked immunosorbent assay using affinity-purified antigen in an area endemic for melioidosis. *Am. J. Trop. Med. Hyg.* 56:418-423.
- Egan, A.M. and Gordon, D.L. 1996. *Burkholderia pseudomallei* activates complement and is ingested but not killed by polymorphonuclear leukocytes. *Infect. Immun.* 64:4952-4959.
- Haase, A., Brennan, M., Barrett, S., Wood, Y., Huffam, S., Brien, D. O., and B. Currie. 1998. Evaluation of PCR for diagnosis of melioidosis. *J. Clin. Microbiol.* 36:1039–1041.
- Holden, M.T., Titball, R.W., Peacock, S.J., Cerdeno-Tarraga, A.M., Atkins, T., Crossman, L.C., Pitt, T., Churcher, C., Mungall, K., Bentley, S.D., Sebahia, M., Thomson, N.R., Bason, N., Beacham, I.R., Brooks, K., Brown, K.A., Brown, N.F., Challis, G.L., Cherevach, I., Chillingworth, T., Cronin, A., Crossett, B., Davis, P., DeShazer, D., Feltwell, T., Fraser, A., Hance, Z., Hauser, H., Holroyd, S., Jagels, K., Keith, K.E., Maddison, M., Moule, S., Price, C., Quail, M.A., Rabinowitsch, E., Rutherford, K., Sanders, M., Simmonds, M., Songsivilai, S., Stevens, K., Tumapa, S., Vesaratchavest, M., Whitehead, S., Yeats, C., Barrell, B.G., Oyston, P.C. and Parkhill, J. 2004. Genomic plasticity of the causative agent of melioidosis, *Burkholderia pseudomallei*. *Proc. Natl. Acad. Sci. USA* 101:14240–14245.
- Inglis, T.J., Chiang, D., Lee, G.S. and Chor Kiang, L. 1998. Potential misidentification of *Burkholderia pseudomallei* by API 20NE. *Pathology.* 30:62-64.

- Ismail, G., Noor, Embi, M., Omar, O., Allen, J.C. and Smith, C.J. 1987. A competitive immunosorbent assay for detection of *Pseudomonas pseudomallei* exotoxin. J. Med. Microbiol. 23:353-357.
- Jitsurong, S., Chirathaworn, C., Brown, N.F., Beacham, I.R. and Tumwasorn, S. 2003. Searching for virulence *Burkholderia pseudomallei* genes by immunoscreening the lambda ZAPII expressed genomic library. Southeast Asian J. Trop. Med. Public. Health. 34:810-21.
- Jones, A.L., Beveridge, T.J. and Woods, D.E. 1996. Intracellular survival of *Burkholderia pseudomallei*. Infect. Immun. 64:782-790.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein Measurement With The Folin Phenol Reagent. J. Biol. Chem. 193, 265–275.
- Miyagi, K., Kawakami, K. and Saito, A. 1997. Role of reactive nitrogen and oxygen intermediates in gammainterferon-stimulated murine macrophage bactericidal activity against *Burkholderia pseudomallei*. Infect. Immun. 65:4108-4113.
- Petkanjanapong, V., Naigowit, P., Kondo, E. and Kanai, K. 1992. Use of endotoxin antigens in enzyme-linked immunosorbent assay for the diagnosis of *P. pseudomallei* infections (Meliodosis). Asian Pacific J. Allergy Immunol.10:145-150.
- Rugdech, P., Anuntagool, N. and Sirisinha, S. 1995. Monoclonal antibodies to *Pseudomonas pseudomallei* and their potential for diagnosis of melioidosis. Am. J. Trop. Med. Hyg. 52: 231-235.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. 1989. Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Santanirand, P., Harley, V.S., Dance, D.A., Drasar, B.S. and Bancroft, G.J. 1999. Obligatory role of gamma interferon for host survival in a murine model of infection with *Burkholderia pseudomallei*. Infect. Immun. 67:3593-3600.
- Serm Swan, R.W., Wongratanacheewin, S., Tattawasart, U. and Wongwajana, S. 1994. Construction of a specific DNA probe for diagnosis of melioidosis and use as an epidemiological marker of *Pseudomonas pseudomallei*. Mol. Cell. Probes 8:1-9.
- Sirisinha, S., Anuntagool, N., Dharakul, T., Ekpo, P., Wongratanacheewin, S., Naigowit, P., Petchelai, B., Thamlikitkul, V. and Suputtamongkol, Y. 2000. Recent developments in laboratory diagnosis of melioidosis. Acta. Tropica. Vol 74:235-245.

- Smith, C.S., Allen, J.C., Embi, M.N., Othman, O., Razak, N. and Ismail, G. 1987. Human melioidosis and emerging medical problem. *MIRCEN. J.* 3: 343-366.
- Smith, M.D., Angus, B.J., Wuthiekanun, V. and White, N.J. 1997. Arabinose assimilation defines a nonvirulent biotype of *Burkholderia pseudomallei*. *Infect. Immun.* 65:4319-4321.
- Stevens, M.P., Wood, M.W., Taylor, L.A., Monaghan, P., Hawes, P., Jones, P.W., Wallis, T.S. and Galyov, E.E. 2002. An Inv/Mxi-Spa-like type III protein secretion system in *Burkholderia pseudomallei* modulates intracellular behaviour of the pathogen. *Mol. Microbiol.* 46: 649-59.
- Tiangpitayakorn, C., Songsivilai, S., Piyasangthong, N. and Dharakul, T. 1997. Speed of detection of *Burkholderia pseudomallei* in blood cultures and its correlation with the clinical outcome. *Am. J. Trop. Med. Hyg.* 57:96-99.
- Walsh A. L. and Wuthiekanun V. 1996. The laboratory diagnosis of melioidosis. *Br. J. Biomed. Sci.* 53(4):249-53.
- Warawa, J. and Woods, D.E. 2005. Type III secretion system cluster 3 is required for maximal virulence of *Burkholderia pseudomallei* in a hamster infection model. *FEMS Microbiol. Lett.* 242(1):101-108.
- White, N.J. 2003. Melioidosis. *The Lancet.* 361:1715-1722.
- Wiersinga, W.J., Tom van der Poll, White, N.J., Day, N.P. and Peacock, S.J. 2006. Melioidosis: insights into the pathogenicity of *Burkholderia pseudomallei*. *Nat. Rev. Microbiol.* 4: 272-282.
- Wild, D. ed. 2001. *The immunoassay handbook*. 2nd ed. London. Nature Publishing.
- Wongratanacheewin, S., Tattawasart, U. and Lulitanond, V. 1990. An avidin-biotin enzyme-linked immunosorbent assay for the detection of *Pseudomonas pseudomallei* antigens. *Trans. R. Soc. Trop. Med. Hyg.* 84:429-430.
- Wongratanacheewin, S., Tattawasart, U., Lulitanond, V., Wongwajana, S., Sermswan, R.W., Sookpranee, M. and Nuntirooj, K. 1993. Characterization of *Pseudomonas pseudomallei* antigens by SDS-polyacrylamide gel electrophoresis and western blot. *Southeast Asian J. Trop. Med. Public. Health.* 24:107-113.
- Wuthiekanun, V., Amornchai, P., Chierakul, W., Cheng, A.C., White, N.J., Peacock, S.J. and Day, N.P. 2004. Evaluation of immunoglobulin M (IgM) and IgG rapid cassette test kits for

diagnosis of melioidosis in an area of endemicity. *J. Clin. Microbiol.* 42:3435-3437.

Wuthiekanun, V., Dance, D., Chaowagul, W., Suputtamongkol, Y., Wattanagoon, Y. and White, N. 1990. Blood culture techniques for the diagnosis of melioidosis. *Eur. J. Clin. Microbiol. Infect. Dis.* 9:654-658.

Wuthiekanun, V., Smith, M.D., Dance, D.A., Walsh, A.L., Pitt, T.L. and White, N.J. 1996. Biochemical characteristics of clinical and environmental isolates of *Burkholderia pseudomallei*. *J. Med. Microbiol.* 45:408-412.

<<http://www.cdc.gov/ncidod/dvbid/plague/bacterium.htm>> (Feb. 26, 2007)

<http://www.hpa.org.uk/infections/topics_az/deliberate_release/Glanders/Homepage.asp#row1>
(Feb. 26, 2007)