

บรรณานุกรม

จินตนา สิริพิทยานนท์. 2537. การวิเคราะห์โครงสร้างผลึก. ภาควิชาเคมี คณะวิทยาศาสตร์ มหาวิทยาลัยเชียงใหม่.

บุญยา รัตนสุภา. 2546. “การเตรียมและศึกษาโครงสร้างสารประกอบเชิงซ้อนของ คอปเปอร์(I)”, รายงานโครงการทางเคมี ภาควิชาเคมี คณะวิทยาศาสตร์ มหาวิทยาลัยสงขลานครินทร์.

ล้ม้าย สิงหโภกม. 2543. “การศึกษาโครงสร้างผลึกและการวิเคราะห์ทางความร้อนของสาร ประกอบเชิงซ้อนคอปเปอร์(I) ไดฟินิดิโซอูเรีย”, วิทยานิพนธ์วิทยาศาสตร์มหาบัณฑิต สาขาวิชาเคมีศึกษา มหาวิทยาลัยสงขลานครินทร์.

วรณี รัตนวี. 2543. “โครงสร้างผลึกของสารประกอบเชิงซ้อนชิโลเวอร์(I) เอทิลีน ไช ไอโซเรียบางต้า”, วิทยานิพนธ์วิทยาศาสตร์มหาบัณฑิต สาขาวิชาเคมีศึกษา มหาวิทยาลัยสงขลานครินทร์.

สัมพันธ์ วงศ์นาวา. 2535. การเรื่องรังสีเอกซ์แบบกระจายพลังงานเบื้องต้น. ภาควิชาเคมี มหาวิทยาลัยสงขลานครินทร์.

Albano, V.G., Bellon, P.L., Ciani, G. and Manasser, M. 1972. “Crystal and molecular structure of monoclinic di- μ -chloro-tris(tri-phenylphosphine)dicopper(I), Cu₂Cl₂(PPh₃)₃”, J.C.S. Dalton. (1972), 171-175.

Aslanidis, P., Hadjikakou, S.K. and Karagiannidis, P. 1993. “Four-coordinate copper(I) iodide complexes with triphenylphosphine heterocyclic thiones ligands. The crystal structure of [Cu(PPh₃)₂(pymtH)I]”, Polyhedron. 12(1993), 2221-2226.

Aslanidis, P., Hadjikakou, S.K. and Karagiannidis, P. 1994. “Preparation and spectral studies of dinuclear mixed-ligand copper(I) complexes. The crystal structure of bis [μ -S (pyridine-2-thione)(tmp) copper(I) bromide]”, Polyhedron. 13(1994), 3119-3125.

Aslanidis, P., Hadjikakou, S.K., Karagiannidis, P., and Cox,P.J. 1998. "Synthesis and characterization of copper(I) complexes with triphenylphosphine and heterocyclic thiones ligands: The structure of (thiazolidine-2-thione) (bis(triphenylphosphine)copper (I) chloride", Inorganica Chimica Acta. 271(1998), 243-247.

Aslanidis, P., Cox, P.J., Divanidis, S. and Karagiannidis, P. 2004. "Trans-1,2-bis (diphenylphosphino)ethane as bridging ligand in thione-S-ligated dimeric copper(I) chloride complexes", Inorganica Chimica Acta. 357(2004), 4231-4239.

Aslanidis, P., Cox, P.J., Divanidis, S. and Karagiannidis, P. 2004. "Copper(I) halide complexes from *cis* -1,2-bis(diphenylphosphino)ethylene and some heterocyclic thiones", Inorganica Chimica Acta. 357(2004), 1063-1076.

Aslanidis, P., Cox, P.J., Divanidis, S. and Tsipis, A. 2002. "Copper(I) halide complexes with 1,3-propanebis(diphenylphosphine) and heterocyclic thione ligands: Crystal and electronic structures (DFT) of [CuCl(pymtH)(dppp)], [CuBr(pymtH)(dppp)] and [Cu(μ -I) (dppp)₂]", Inorganic Chemistry. 41(2002), 6875-6886.

Bowmaker, G. A. 1994. "Vibrational spectra and crystal structure of tris- and trtrakisis(ethylenethiourea)copper(I) systems", Aust.J. Chem. 47(1994), 15-24.

Cotton, F.A. and Wilkinson, G. 1988. Advanced Inorganic Chemistry, 5th ed., New york : John Wiley & Sons.

Bukong, S. 2004. "Crystal structures of copper oxyanion complexes containing diethyl thiourea and ethylenethiourea ", M.Sc.Physical Chemistry, Prince of Songkla University.

Cox, P.J., Aslanidis, P., Hadjikakou, S.K. and Karagiannidis, P. 1999. "Synthesis, spectroscopic and computational studies plus crystal structure of [(bis-benzo-1,3-thiazolidine-2-thione)(bis-triphenylphosphine)copper(I)] [Chloro(benzothiazolidine-2-thione)(bis-triphenylphosphine)copper(I)]", Polyhedron. 18(1999), 1501-1506.

Divanidis, S., Cox, P.J., Karagiannidis, P. and Aslanidis, P. 2005. "Linking thione-ligated copper (I) centres with trans-1,2-bis(diphenylphosphino)ethane (trans-dppen): crystal structure of the pyrimidine-2-thione derivatives $[\text{CuBr}(\mu_2\text{-trans-dppen})(\text{pymtH})]_2$ and $[\text{CuI}(\mu_2\text{-trans-dppen})(\text{pymtH})]$ ", Polyhedron. 24(2005), 351-358.

Hadjikakou, S.K., Aslanidis, P. and Karagiannidis, P. 1991. "Synthesis and photolysis of mixed copper(I) complexes with thiones and tri-p-tolylphosphine or triphenylphosphine; X-ray crystalstructure of bis[copper(I)(1,3-thiazolidine-2-thione)(tri-p-tolylphosphine) chloride]", Polyhedron. 10(1991), 935-940.

Hadjikakou, S.K., Aslanidis, P., Akrivos, P.D. and Karagiannidis, P. 1992. "Study of mixed ligand copper(I) complexes with tri-m-tolylphosphine (tmtp) and heterocyclic thiones. Crystal structures of bis[μ -S(benzimidazoline-2-thione)(tmtp) copper(I) chloride] and bis[μ -Br(benzimidazoline-2-thione)(tmtp) copper(I)]", Inorganica Chimica Acta. 197 (1992), 31-38.

Hadjikakou, S.K., Aslanidis, P. and Karagiannidis, P. 1991. "Synthesis and photolysis of a new series of Cu(I) complexes with tri-*m*-tolylphosphine and heterocyclic thiones. The crystal structure of bis[μ -iodo(pyrimidine-2-thione)(tri-*m*-tolylphosphine) copper(I)]", Inorganica Chimica Acta. 184(1991), 161-166.

Hadjikakou, S.K., Aslanidis, P. and Karagiannidis, P. 1991. "Synthesis and photolysis of a new series of Cu(I) complexes with tri-*o*-tolylphosphine and heterocyclic thiones as ligands. The crystal structure of (thiazolidine-2-thione)(tri-*o*-tolylphosphine) copper(I) bromide", Inorganica Chimica Acta. 186(1991), 199-204.

Isab, A.A., Fettouhi, M., Ahmad, S. and Ouahab, L. 2003. "Mixed ligand gold(I) complexes of phosphines and thiourea and X-ray structure of (thiourea-KS)(tricyclohexylphosphine) gold(I) chloride", Polyhedron. 22(2003), 1349-1354.

Jianping Lang. and Kazuyumi Tatsumi. 1996. "Synthesis and crystal structure of a binuclear copper (I) bromide complex of benz-1,3-thiazolidine-2-thione and triphenylphosphine, $[\text{Cu}(\text{bztdtH})(\text{PPh}_3)\text{Br}]_2$ ", Polyhedron. 15(1996), 981-986.

Karagiannidis, P., Aslanidis, P. and Papastefanou, S. 1989. "Cu(I) complexes with heterocyclic thiones and triphenylphosphine as ligands the crystal structure of $[\text{Cu}(\text{tzdtH})_2(\text{PPh}_3)_2]\text{NO}_3$ ", Inorganica Chimica Acta. 156(1989), 265-270.

Karagiannidis, P., Aslanidis, P. and Papastefanou, S. 1990. "Synthesis and characterization of copper(I) halide complexes with heterocyclic thiones and triphenylphosphine as ligands. The X-ray crystal structure of copper(I)1-methyl-1,3-imidazoline-2-thione bis (triphenylphosphine)bromide, $[\text{Cu}(\text{PPh}_3)_2\text{meimtH}\text{Br}]$ ", Polyhedron. 9(1990), 981-986.

Karagiannidis, P., Hadjikakou, S.K., Aslanidis, P. and Hountas, A. 1990. "Synthesis and photochemical study of Cu(I) complexes with tri-p-tolylphosphine and heterocyclic thiones. The crystal structure of $[\text{CuCl}(\text{pymtH})(\text{p}-\text{CH}_3\text{C}_6\text{H}_4)_3\text{P}]_2$ ", Inorganica Chimica Acta. 178(1990), 27-34.

Lemos, S.S., Carmargo, M.A., Cardoso, Z.Z., Deflon, V.M., Forsterling, F.H. and Hagenbah, A. 2001. "Copper(I) pseudohalide complexes with 4,6-dimethylpyridine-2(1H)-thione and triphenylphosphane as ligands. The X-ray crystal structures of $[\text{Cu}(\text{N}_3)(\text{dmpytmH})(\text{PPh}_3)_2]$ and $[\text{Cu}(\text{NCS})(\text{dmpytmH})(\text{PPh}_3)_2]$ ", Polyhedron. 20(2001), 849-854.

Lecomte, C. and Skoulika, St. 1989. "Copper(I) bromide complexes with heterocyclic thiones and triphenylphosphine as ligands. The X-ray crystal structure of copper(I) pyrimidine-2-

thione bis(triphenylphosphine)bromide [Cu(PPh₃)₂(PymtH)Br]”, Polyhedron. 8 (1989), 1103-1109.

Lobana, T.S. and Castineiras, A. 2002. “Metal-heterocyclic thione interactions. 13. Pyridine-2-thione derivatives of copper(I): crystal structure of dinuclear [bromo(pyridine-2-thione)(tri-*p*-tolylphosphine)copper(I)]₂ complex”, Polyhedron. 21(2002), 1603-1611.

Okaya, Y. and Knobler, C. B. 1964. “Refinement of the crystal structure of tris(thiourea) copper (I) chloride”, Acta. Cryst. 17(1964), 928-930.

Ramsharan Singh. and Dikshit, K. 1995. “Synthesis and characterization of mixed ligand copper (I) complexes containing halide,triphenylarsine and N,N-dimethyl-N-phenylthiourea (dmptH), N,N-dibutyl-N-phenylthiourea (dbptH) or 1,3-thiazolidine-2-thione (tzdtH). The X-ray crystal structure of [Cu(PPh₃)₂(dmptH)Cl]”, Polyhedron. 14(1995), 1799-1995.

Rout, G.C., Seshasayee, M., Aravamudan, G. and Sowrirajan, S. 1984. “Synthesis and structure of tetrakis(phenylenthiourea) tellurium(II)chloride dihydrochloride, C₂₈H₂₆N₈S₄Cl₄Te”, Polyhedron. 3(1984), 921-927.

Saithong, S. 2003. “Crystal structures of silver(I) complexes containing ethylenethiourea and diethyl thiourea ligands”, M.Sc.Physical Chemistry, Prince of Songkla University.

Satyanarayana, S. and Nagasundara, K.R. 2004. “Synthesis and spectral properties of the complexes of cobalt(II), nickel(II), copper(II), Zinc(II), and cadmium(II) with 2-(thiomethyl-2-benzimidazolyl-benzimidazole”, Synthesis and reactivity in inorganic and metal-organic chemistry. 34(2004), 883-895.

Schmidbaur, R. Reber, G., Schier, A., Wagner, F.E. and Muller, G. 1988. “Structural correlation between *trans*- and *cis*-bis(diphenylphosphino)ethane, bis(diphenylphosphino)methane and their chlorogold(I) complexes”, Inorganica Chimica Acta. 147(1998), 143-150.

Skoulika, S. and Aubry, A. 1991. "New copper(I) chloride complexes with heterocyclic thiones and triphenylphosphine as ligands. Crystal structure of $[\text{Cu}(\text{PPh}_3)_2(\text{bzimtH}_2)\text{Cl}]$ CH_3COCH_3 and $[\text{Cu}(\text{PPh}_3)_2(\text{nbzimtH}_2)\text{Cl}]$ ", *Inorganic Chimica Acta*. 183(1991), 207-211.