

เอกสารอ้างอิง

วัฒนาลัย ปานบ้านเกร็ด และ สรวง อุดมภารกุณฯ. 2536. เทคนิคทางเอนไซม์พันธุศาสตร์และพันธุวิกรรม. สมาคมเทคโนโลยีชีวภาพแห่งประเทศไทย.

อรยา สุตเชียรกุล. 2541. โรคติดเชื้อ. พิมพ์ครั้งที่2. กรุงเทพฯ: ไฮลิสติก พับลิชชิ่ง จำกัด.

Abbott, S.L. and Janda, J.M. 1994. Severe gastroenteritis associated with *Vibrio hollisae* infection: Report of two cases and review. Clin. Infect. Dis. 18: 310-312.

American Public Health Association: American water works association and water pollution control federation: Standard methods for the examination of water and waste water, 1981. 15th ed., Washington.

Atlas, R.M. 1997. Handbook of Microbiological media. 2nd ed. CRC Press.

Arnheim, N. and Erlich, H. 1992. Polymerase chain reaction strategy. Annu. Rev. Biochem. 61:131-156.

Birnboim, H.C. and Doly, J. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. Nucleic Acids Res. 7:1513-1523.

Blake, P.A. 1983. Vibrios on the half shell: what the walrus and the carpenter didn't know. Ann Intern Med. 99: 558-559.

Blake, P.A., Weaver, R.E. and Hollis, D.G. 1980. Diseases of humans other than cholera caused by vibrios. Annu. Rev. Microbiol. 34:341-367.

Cakir, I., Dogan, H.B., Halman, A.K., Worobo, R.W. 2001. An alternative approach for enumeration of *Escherichia coli* in food. Int. J. Food. Microbiol. 68:217-223.

Carnahan, A.M., Hardling, J., Watsky, D. and Hansman, S. 1994. Identification of *Vibrio cholerae* associated with severe gastroenteritis after consumption of raw oyster. J. Clin. Microbiol. 32:1805-1806.

Chang, B., Taniguchi, H., Miyamoto, H. and Yoshida, S. 1998. Filamentous bacteriophage of *Vibrio parahaemolyticus* as a possible clue to genetic transmission. J. Bacteriol. 180:5094-5101.

Chang, B., Yoshida, S., Miyamoto, H., Ogawa, M., Horikawa, M., Ogata, K., Nishibuchi, M. and Taniguchi, H. 2000. A unique and common restriction fragment pattern of the nucleotide sequences homologous to the genome of Vf33, a filamentous bacteriophage, in pandemic strains of *Vibrio parahaemolyticus* O3:K6, and O1:K untypeable. FEMS. Microbiol. Lett. 192:231-236.

Danikina, E.B., Smolikova, L.M., Gurleva, G.G., Voronezhskaya, L.G., Kharabadzhakhian, G.D. and Bogdanova, M. 1995. Choice of commercial selective nutrient media for the isolation of pathogenic vibrios of different species. Klinicheskaya. Laboratornaia. Diagnostika. 4:50-53.

DiRita, V.J. 1992. Co-ordinated expression of virulence gene by ToxR in *Vibrio cholerae*. Mol. Microbiol. 6:451-458.

- Echeverria, P., Pitarangsi, C., Eampokalap, B., Vibulbandhitkit, S., Boonthai, P. and Rowe, B. 1983. A longitudinal study of the prevalence of bacterial enteric pathogens among adults with diarrhea in Bangkok, Thailand. Diagnostic Microbiol and Infect Dis. 1:193-204.
- Fabbri, A., Falzano, L., Frank, C., Donelli, G., Matarrese, P., Raimondi, F., Fasano, A. and Fiorentini, C. 1999. *Vibrio parahaemolyticus* thermostable direct hemolysin modulates cytoskeletal organization and homeostasis in intestinal cultured cells. Infect. Immun. 67:1139-1148.
- Fasano, A.B., Baudry, B., Pumplin, D.W., Wasserman, S.S., Tall, B.D., Ketley, J.M. and Kaper, J.B. 1991. *Vibrio cholerae* produce a second enterotoxin, which affects intestinal tight junctions. Proc. Natl. Acad. Sci. USA. 88:5242-5246.
- Hanchalay, S., Seriwatana, J., Echeverria, P., Holmgren, J., Tirapat, C., Moseley, S.L. and Taylor, D.N. 1985. Non-O1 *Vibrio cholerae* in Thailand: homology with cloned cholera toxin genes. J. Clin. Microbiol. 21:288-289.
- Haughes, J.M., Hollis, D.G., Gangarosa, E.J. and Weaver, R.E. 1978. Non-cholera vibrio infection in the United States. Clinical, epidemiologic and laboratory features. Ann. Intern. Med. 88:602-606.
- Henson, J.M. and French, R. 1993. The Polymerase chain reaction and plant disease diagnosis. Annu. Rev. Phytopathol. 31:81-109.
- Hickman, F.W., Farmer, J.J., Hollis, D.G., Fanning, G.R., Steigerwalt, A.G., Weaver, R.E. and Brenner, D.J. 1982. Identification of *Vibrio hollisae* sp. Nov. from patients with diarrhea. J. Clin. Microbiol. 15:395-401.

- Hlady, W.G. 1997. *Vibrio* infections associated with raw oyster consumption in Florida, 1981-1994. *J. food Protect.* 60:353-357.
- Hoge, C.W., Watsky, D., Peeler, R.N., Libonati, J.P., Isarel,E. and Moris, J.G. 1989. Epidemiology and spectrum of Vibrio infection in a Chesapeake Bay community. *J. Infect Dis.* 160:985-993.
- Honda, T., Goshima, K., Takeda, Y., Sugino, Y. and Miwatani, T. 1976. Demonstration of the cardiotoxicity of the thermostable direct hemolysin (lethal toxin) produced by *Vibrio parahaemolyticus*. *Infect. Immun.* 13:163-171.
- Honda, T. and Iida, T. 1993. The pathogenicity of *Vibrio parahaemolyticus* and the role of the thermal stable direct hemolysin and related hemolysins. *Rev. Med. Microbiol.* 4: 106-113.
- Huntley, J.S., Hall, A.C., Sathyamoorthy, V. and Hall, R.H. 1993. Cation flux studies of the lesion induced in human erythrocyte membranes by the thermostable direct hemolysin of *Vibrio parahaemolyticus*. *Infect. Immun.* 61:4326 - 4332.
- Kar, S., Ghosh, R.K., Ghosh, A.N. and Ghosh, A. 1996. Integration of the DNA of novel filamentous bacteriophage VSK from *Vibrio cholerae* O139 into the host cromosomal DNA. *FEMS. Microbiol. Lett.* 145:17-22.
- Lee, S.E., Shin, S.H., Kim, Y.S., Kim, Y.R., Shin, D.H., Chung, S.S., Lee, Z.H., Lee, J.Y., Jeong, K.C., Chol, S.H. and Rhee, J.H. 2000. *Vibrio vulnificus* has the transmembrane transcription activator ToxRS stimulating the expression of the hemolysin gene *vvha*. *J. Bacteriol.* 182:3405-3415.

- Levine, W.C. and Griffin, P.M., 1993. Vibrio infection on the gulf coast: results of first year of regional surveillance. *J. Infect Dis.* 167:479-483.
- Lichstein, H.C., Synder, M.I. 1941. The inhibition of the spreading growth of *Proteus* and other bacteria to permit the isolation of associated Streptococci. *J. Bact.* 42:653-664.
- Lin, Z., Kumagai, K., Baba, K., Mekalanos, J.J. and Nishibuchi, M. 1993. *Vibrio parahaemolyticus* has a homolog of the *Vibrio cholerae* *toxRS* operon that mediates environmentally induced regulation of the thermostable direct hemolysin gene. *J. Bacteriol.* 175:3844-3855.
- Lowry, P.W., Mcfarland, L.M. and Threelfoot, H.K. 1986. *Vibrio hollisae* septicemia after consumption of catfish. *J. Infect Dis.* 154:730-731.
- Marvin, D.A. 1974. Filamentous bacterial viruses. XII, Molecular architecture of the class I (fd, Ifl, Ike) virion. *J. Mol. Biol.* 88:581-600.
- Marvin, D.A., Wiseman, R.L. and Wachtel, E.J. 1974. Filamentous bacterial viruses. XI, Molecular architecture of the class II (pfl, xf) virion. *J. Mol. Biol.* 82:121-138.
- Miller, V.L., Taylor, R.K. and Mekalanos, J.J. 1987. Cholera toxin transcriptional activator ToxR is transmembrane DNA binding protein. *Cell.* 48:271-279.
- Miliotis, M.D., Tall, B.D. and Gray, R.T. 1995. Adherence to and invasion of tissue culture cells by *Vibrio hollisae*. *Infect. Immun.* 63:4956-4963.

- Miyamoto, Y., Kato, T., Obara, Y., Akiyama, S., Takizawa, K. and Yamai, S. 1969. In vitro hemolytic characteristic of *Vibrio parahaemolyticus*: its close correlation with human pathogenicity. J. Bacteriol. 100:1147-1149.
- Mizunoe, Y., Wai, S.N. and Takeda, A. 1999. Restoration of culturability of starvation-stressed and low-temperature-stressed *Escherichia coli* O157 cells by using H₂O₂-degenerating compounds. Arch Microbiol. 172:63-67.
- Morris, J.G., Wilson, R., Hollis, D.G., Weaver, R.E., Miller, H.G., Tacket, C.O., Hickman, F.W. and Blake, P.A. 1982. Illness caused by *V. damsela* and *V. hollisae*. Lancet. 1294-1296.
- Myatt, D.C. and Davis, G.H. 1989. Isolation of medically significant *Vibrio* species from riverine sources in southeast Queensland. Microbios. 60:111-123.
- Naim, R., Yanagihara, I., Lida, T. and Honda, T. 2001. *Vibrio parahaemolyticus* thermostable direct hemolysin can induce an apoptotic cell death in Rat-1 cells inside and outside of the cells. FEMS. Microbiol. Letter. 195:237-244.
- Nishibuchi, M. and Kaper, J.B. 1995. Thermostable direct hemolysin gene of *Vibrio parahaemolyticus*: a virulence gene acquired by marine bacterium. Infect. Immun. 63:2093-2099.
- Nishibuchi, M., Doke, S., Toizumi, S., Umeda, T., Yoh, M. and Miwatani, T. 1988. Isolation from a coastal fish of *Vibrio hollisae* capable of producing a hemolysin similar to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. Appl. Environ. Microbiol. 58:2144-2146.

- Nishibuchi, M., Hill, W.E., Zon, G., Payne, W.L. and Kaper, J.B. 1986. Synthetic oligonucleotide probes to detect Kanagawa phenomenon-positive *Vibrio parahaemolyticus*. Infect. Immun. 23:1091-1095.
- Nishibuchi, M., Ishibashi, M., Takeda, Y. and Kaper, J.B. 1985. Detection of the thermostable direct hemolysin gene and related DNA sequence in *Vibrio parahaemolyticus* and other *Vibrio* species by the DNA colony hybridization test. Infect. Immun. 49:481-486.
- Nishibuchi, M., Janda, J.M. and Ezaki, T. 1996. The thermostable direct hemolysin gene (TDH) of *Vibrio hollisae* is dissimilar in prevalence to and phylogenetically distant from the *tdh* genes of other Vibrios: Implication in the horizontal transfer of the *tdh* gene. Microbiol. Immun. 40:59-65.
- Nishibuchi, M., Khaeomanee-iam, V., Honda, T., Kaper, J.B. and Miwatani, T. 1990. Comparative analysis of the hemolysin genes of *Vibrio Cholerae* non-O1, *V. mimicus* and *V. hollisae* that are similar to the *tdh* gene of *V. parahaemolyticus*. FEMS. Microbiol. Lett. 55:251-256.
- O'Brien, M. and Cowell, R. 1985. Modified taurocholate-tellurite-gelatin agar for improved differentiation of *Vibrio species*. Clin. Micro. 22:1011-1013.
- Osario, C.R. and Klose, K.E. 2000. A region of the transmembrane regulatory protein ToxR that tethers the transcriptional activation domain to the cytoplasmic membrane displays wide divergence among *Vibrio* species. J. Bacteriol. 182:526-528.

- Pearson, G.D., Woods, A., Chiang, S.L. and Mekalanos, J.J. 1993. CTX genetic element encodes a site-specific recombination system and an intestinal colonization factor. Proc. Natl. Acad. Sci. USA. 90:3750-3754.
- Pfau, J.D. and Taylor, R.K. 1998. Mutation in toxR that separate transcriptional activation from DNA binding at cholera toxin gene promoter. J. Bacteriol. 180:4724-4733.
- Rank, E.L., Smith, I.B. and Langer, M. 1988. Bacteremia caused by *Vibrio hollisae*. J. Clin. Microbiol. 26:375-376.
- Raimondi, F., Kao J.P., Kaper, J.B., Guandalini, S. and Fasano, A. 1995. Calcium-dependent intestinal chloride secretion by *Vibrio parahaemolyticus* thermostable direct hemolysin in a rabbit model. Gastroenterol. 109: 381-386.
- Raimodi, F., Kao, J. P.Y., Florentini, C., Fabbri, A., Donelli, G., Gasparini, N., Rubino, A. and Fasano, A. 2000. Enterotoxicity and cytotoxicity of *Vibrio parahaemolyticus* thermostable direct hemolysin *in vitro* system. Infect. Immun. 68:3180-3185.
- Reich, K.A. and Schoolnik, G.K., 1994. The light organ symbiont *Vibrio fisheri* possesses a homolog of the *Vibrio cholerae* transmembrane transcriptional activator ToxR. J. Bacteriol. 176:3085-3088.
- Rouzet, S.G., Donnio, P.Y., Juguet, F.J., Plessis, P., Minet, J. and Avril, J.L. 1996. First European case of gastroenteritis and bacteremia due to *Vibrio hollisae*. Euro. J. Clin. Microbiol. Infect. Dis. 15:864-866.

Sakazaki, R., Tamura, K., Kato, T., Obara, Y., Yamai, S. and Hobo, K. 1968. Studies of the enteropathogenic, facultatively halophilic bacteria, *Vibrio parahaemolyticus*. III. Enteropathogenicity. Jpn. J. Med. Sci. Biol. 21:325-331.

Sakazaki, R., Tamura, K., Nakamura, A., Kurata, T. and Goda, A. 1974. Studies on enteropathogenic activity of *Vibrio parahaemolyticus* using ligated gut loop model in rabbits. Jpn. J. Med. Sci. Biol. 27:35-43.

Sakurai, J., Honda, T., Jinguji, Y., Arita, M. and Miwatani, T. 1976. Cytotoxic effect of the thermostable direct hemolysin produced by *Vibrio parahaemolyticus* on FL cells. Infect. Immun. 13:876-883.

Sambrook, J., Fritsch, F. and Maniatis, T. 1989. Molecular cloning: a laboratory manual. 2nd ed. Cold Spring Harbor Laboratory Press, New York.

Takeda, Y. 1988. Thermostable direct hemolysin of *Vibrio parahaemolyticus*. *Methods Enzymol.* 165: 189-193.

Tang, G.Q., Iida, T., Yamamoto, K. and Honda, T. 1994. A mutant toxin of *Vibrio parahaemolyticus* thermostable direct hemolysin which has lost hemolytic activity but retain ability to bind to erythrocytes. Infect. Immun. 62:3299-3304.

Tang, G.Q., Iida, T., Yamamoto, K. and Honda, T. 1995. Ca²⁺ independent cytotoxic of *Vibrio parahaemolyticus* thermostable direct hemolysin (TDH) on intestine 407, a cell derived from human embryonic intestine. FEMS. 134:233-238.

Taniguchi, H., Sato, K., Ogawa, M., Udou, T. and Mizuguchi, Y. 1984. Isolation and characterization of a filamentous phage, Vf33, specific for *Vibrio parahaemolyticus*. *Microbiol. Immunol.* 28:327-337.

Thamlikitkul, V. 1990. Vibrio bacteremia in Siriraj Hospital. *Thailand J. Med. Assoc.* 73:136-139.

Tortora, G.J., Funke, B.R. and Case, C.L. 1997. *Microbiology. An Introduction.* 6th ed. Wesley Longman, Inc.

Trucksis, M., Galen, J.E., Michalski, J., Fasano, A. and Kaper, J.B. 1993. Accessory cholera enterotoxin (Ace), the third toxin of a *Vibrio cholerae* virulence cassette. *Proc. Natl. Acad. Sci. USA.* 90:5267-5271.

Van Schothorst, M. and Renaud, A.M. 1985. Malachite green pre-enrichment medium for improved salmonella isolation from heavily contaminated samples. *Appl. Bact.* 59:223-230.

Vuddhakul, V., Chowdhury, A., Laohaprerthisan, V., Pungrasamee, P., atararungrong, N., Thianmontri, P., Ishibashi, M., Matsumoto, C. and Nishibuchi, M. 2000. Isolation of a pandemic O3:K6 clone of a *Vibrio parahaemolyticus* strain from environmental and clinical sources in Thailand. *Appl. Environ. Microbiol.* 66:2685-2689.

- Vuddhakul, V., Nakai, T., Matsumoto, C., Oh, T., Nishino, T., Chen, C.H., Nishibuchi, M. and Okuda, J. 2000. Analysis of *gyrB* and *toxR* gene sequences of *Vibrio hollisae* and development of *gyrB*-and *toxR*-targeted PCR methods for isolation of *Vibrio hollisae* from the environment and its identification. *Appl. Environ. Microbiol.* 66:3506-3514.
- Vuddhakul, V., Patararungrong, N., Pungrasamee, P., Jitsurong, S., Morigaki, T., Asai, N. and Nishibuchi, K. 2000. Isolation and characterization of *Escherichia coli* O157 from retail beef and bovine feces in Thailand. *FEMS. Microbiol. Lett.* 182:343-347.
- Waldor, M.K. and Mekalanos, J.J. 1996. Lysogenic conversion by a filamentous phage encoding cholera toxin. *Science*. 272:1910-1914.
- Waldor, M.K., Rubin, E.J., Pearson, G.D., Kimsey, H. and Mekalanos, J.J. 1997. Regulation, replication, and integration functions of the *Vibrio cholerae* CTX ϕ are encoded by region RS2. *Mol. Microbiol.* 24:917-926.
- Wong, H.C., Chen, M.C., Liu, S.H. and Liu, D.P. 1999. Incidence of highly genetically diversified *Vibrio parahaemolyticus* in seafood imported from Asian countries. *International. Food. Microbiol.* 52:181-188.
- Wright, R.C. 1984. A new selective and differential agar medium for *Escherichia coli* and coliform organisms. *Apply. Bacteriol.* 56:381-388.
- Yamasaki, S., Shirai, H., Takeda, Y. and Nishibuchi, M. 1991. Analysis of the gene of *Vibrio hollisae* encoding the hemolysin similar to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. *FEMS. Microbiol. Lett.* 64:259-263.

Yoh, M., Honda, T. and Miwayami, T. 1986. Purification and partial characterization of *Vibrio hollisae* hemolysin that relates to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. Can. J. Microbiol. 32:632-636.

Yoh, M., Honda, T. and Miwatani, T. 1988. Comparison of hemolysins of *Vibrio cholerae* non-O1 and *Vibrio hollisae* with thermostable direct hemolysin of *Vibrio parahaemolyticus*. Can J. Microbiol. 34:1321-1324.

Yoh, M., Honda, T. and Miwatani, T. 1989. Homogeneity of a hemolysin (Vh-rTDH) produced by clinical and environment isolates of *Vibrio hollisae*. FEMS. Microbiol. Lett. 52:171-175.

Yoh, M., Honda, T., Miwatani, T., Tsunasawa, S. and Sakiyama, F. 1989. Comparative amino acid sequence analysis of hemolysins produced by *Vibrio hollisae* and *Vibrio parahaemolyticus*. J. Bacteriol. 171:6859-6861.