

## References

1. Boylan, J.C. et al, Handbook of pharmaceutical excipients 1986,APhA, USA
2. Cheong, L.W.S. et al, Relationship between polymer viscosity and drug release from a matrix system Pharm. Res. 1992 ; 9(11) 1510-1514
3. Colombo,P. et al,Drug release modulation by physical restrictions of matrix swelling, Int.J.Pharm. 1990; 63, 43-48
4. Colombo, P. et al, Swelling characteristics of hydrophilic matrices for controlled release, Int.J.Pharm. 1992 ; 88 ,99-109

5. Gennaro, A.R. et al, Remington's Pharmaceutical Sciences 1985, Mack publishing company
6. Herman, J. and Remon, J.P., Modified starches as hydrophobic matrix for controlled oral delivery, *Int.J.Pharm.* 1989 ; 56, 65-75
7. Lin, S.Y. and Ayres, J.W., Calcium alginate beads as core carriers of 5-aminosalicylic acid, *Pharm. Res.* 1992 ; 9(9) 1128-1131
8. Lin, S.Y., Kao, Y.H., Effect of Eudragit resins and dibasic calcium phosphate on the compaction and dissolution behavior of directly compressible controlled release theophylline tablet, *Drug Dev. Ind. Pharm.* 1991 ; 16(5) , 855-874
9. Lu, M.F. et al, Xanthan gum and alginate based controlled release theophylline formulations, *Drug Dev. Ind. Pharm.* 1991 ; 17(14), 1987-2004
10. Marini, J.O. et al, Some factors affecting the release of drug from membrane coated slow release tablets, 1991; 17(6) 865-876
11. Martinsen, A. and Storrø, G., Alginates as immobilization material: III diffusional properties , *Biotech. Bioeng.* , 1992 ; 39, 186-194
12. Peppas, N.A., Sahlin, J.J., A simple equation for the description of solute release. III Coupling of diffusion and relaxation . *Int. J.Pharm.*, 57(1989) 169-172
13. Shah, A.C. et al, Gel-matrix systems exhibiting bimodal controlled release for oral drug delivery *J. Control. Release*, 9(1989) , 169-175.
14. Tekeuchi, H. et al , Controlled release theophylline tablet with acrylic polymers prepared by spray drying technique , *Drug Dev. Ind. Pharm.* 1989 ; 15(12) 1999-2016