

References

1. Shibouta Y, Inada Y, Ojima M et al. Pharmacological profile of a highly potent and long-acting angiotensin II receptor antagonist, 2-ethoxy-1-{[2'-(1H-tetrazol-5-yl)biphenyl-4-yl]methyl}-1H-bezimidazole-7-carboxylic acid (CV-11974), and its product, (\pm)-1-(cyclohexyloxycarbonyloxy)-ethyl2-ethoxy-1-{[2'-(1H-tetrazol-5-yl)biphenyl-4-yl]methyl}-1H-benzimidazole-7-carboxylate (TCV-116). *J Pharmacol Exp Ther* 1993;266:114-120.
2. Cervenka L, Wang C-T, Navar LG. Effects of acute AT1 receptor blockade by candesartan on arterial pressure and renal function in rats. *Am J Physiol* 1998;274:F940-F945.
3. Tamaki T, Nishiyama A, Yoshida H et al. Effects of EXP3174, a non-peptide angiotensin II receptor antagonist, on renal hemodynamics and renal function in dogs. *Eur J Pharmacol* 1993;236:15-21.
4. Li XC, Widdop RE. Regional hemodynamic effects of the AT1 receptor antagonist CV-11974 in conscious renal hypertensive rats. *Hypertension* 1995;26:989-997.
5. Thomsen K. Lithium clearance: a new method for determining proximal and distal tubular reabsorption of sodium and water. *Nephron* 1984;37:217-223.
6. Thomsen K, Leyssac PP. Effect of dietary sodium content on renal handling of lithium: Experiments in conscious diabetes insipidus rats. *Pflüger Arch* 1986;407:55-58.
7. Kirchner KA. Lithium as a marker for proximal tubular delivery during low salt intake and diuretic infusion. *Am J Physiol* 1987;253:F188-F196.
8. Koomans HA, Boer WH, Dorhout-ME. Evaluation of lithium clearance as a marker of proximal tubule sodium handling. *Kidney Int* 1989;36:2-12.
9. Fuhr J, Kaczmarczyk J, Krutten CD. Eine einfache colorimetrische Methode zur Inulinbestimmung für Nieren-Clearance-Untersuchungen bei Stoffwechselgesunden und Diabetikern. *Klin Wochensch* 1955;33:729-730.
10. Smith HW, Finkelstein N, Aliminosa L. The renal clearances of substituted hippuric acid derivatives and other aromatic acids in dogs and man. *J. Clin. Invest* 1945;42: 388-404.
11. Smart ML, Hiranyachattada S, Harris PJ. Effects of angiotensin II receptor blockade on proximal fluid uptake in the rat kidney. *Br J Pharmacol* 1999;126:697-700.
12. Zhou J, Thomas D, Harris PJ, Skinner SL. The role of endogenous angiotensin II in the regulation of renal hemodynamics and proximal fluid reabsorption in the rat. *J Physiol* 1992;453:1-13.

13. Brown GP, Douglas JG. Angiotensin II binding sites on isolated rat renal brush border membranes. *Endocrinology* 1982;111:1830-1836.
14. Thekkumkara TJ, Cookson, R, Linas S. Angiotensin (AT1) receptor-mediated increases in transcellular sodium transport in proximal tubule cells. *Am J Physiol* 1998;274:F897-F905.
15. Harris PJ, Hiranyachattada S, Antoine AM et al. Regulation of renal tubular sodium transport by angiotensin II and atrial natriuretic factor. *Clin Exp Pharmacol Physiol* 1996;Suppl3:S112-S118.
16. Garvin JL. Angiotensin stimulates bicarbonate transport and Na^+/K^+ ATPase in rat proximal straight tubules. *J Am Soc Nephrol* 1991;1:1146-1152.
17. Aperia A, Holtback U, Syren ML et al. Activation and deactivation of renal Na^+/K^+ ATPase: a final common pathway for regulation of natriuresis. *FASEB J* 1994;8:436-439.
18. Bomsztyk K, Wright FS. Dependence of ion fluxes on fluid transport by rat proximal tubule. *Am J Physiol* 1986;250:F680-F689.
19. Li XC, Widdop RE. Angiotensin type I receptor antagonists CV-11974 and EXP 3174 cause selective renal vasodilatation in conscious spontaneously hypertensive rats. *Clin Sci* 1996;91:147-154.
20. Foidart J, Sraer J, Delarue F, et al. Evidence for mesangial glomerular receptors for angiotensin II linked to mesangial cell contractility. *FEBS Lett* 1980;121:333-339.
21. Ichikawa I, Brenner BM. Glomerular actions of angiotensin II. *Am J Med* 1984;76:43-49.
22. Ardaillou R, Sraer J, Chansel D, et al. The effects of angiotensin II on isolated glomeruli and cultured glomerular cells. *Kidney Int* 1987;31(suppl 20):S74-S80.
23. Burnier M, Brunner HR. Angiotensin receptor antagonist in hypertension. *Kidney Int* 1998;54 (suppl 68):S107-S111.