

References

- Al-Meshal, M.A. and Bayomi, A., 1995, Evaluation of egg albumin as a filler for prolonged release direct compressed tablets. *Drug Dev. Ind. Pharm.* 21 (6), 739-745.
- Aubry, A. and Wainer, I.W., 1993. An in vitro study of the stereoselective dissolution of (rac)-verapamil from two sustained release formulations. *Chirality*, 584-590.
- Cabral Marques, H.M., Hardgarft, Kellaway, I.W. and Pugh, W.J., 1990. Studies of cyclodextrin inclusion complexes. II. Molecular modelling and ¹H-NMR evidence for the salbutamol- β -cyclodextrin complex. *Int. J. Pharm.* 63, 267-274.
- Carr, R.A., Pasutto, F.M., Longstreth, J.A. and Foster, R.T., 1993. Stereospecific determination of the in vitro dissolution of modified release formulations of (\pm)-verapamil. *Chirality*, 5, 443-447.
- Chapman, I.D., Buchheit, K.H., Manley, P., Morley, J., 1992. Active enantiomers may cause adverse effects in asthma. *Trends Pharmacol. Sci.* 13, 231-232.
- Boulton, D.W. and Fawcett, J.P., 1997. Pharmacokinetics and pharmacodynamic of single oral doses of albuterol and its enantiomers in humans. *Clin. Pharmacol. Ther.* 62, 138-144.
- Dong, Z.C. and Jiang, J.T., 1996. Mechanism of soluble drug release from hydrophilic matrices, the effect of hydroxypropyl methylcellulose (HPMC) content. *Acta Pharm. Sinica*. 31, 43-47.
- Mora, M. and Pato, J., 1995. Preparation of egg albumin microparticles for oral applications. *J. Contr. Rel.* 35, 107-112.

- Duddu, S.P., Vakilynejad, M., Jamali, F. and Grant, D.J.W., 1993. Stereoselective dissolution of propranolol hydrochloride from hydroxypropyl methylcellulose matrices. *Pharm. Res.* 101, 648-1653.
- Esquisabel, A, Hernandez, RM, Gacon, AR, Igartua, M. and Pedraz, JT, 1997. Determination of salbutamol enantiomers by high performance capillary electrophoresis and its application to dissolution assays. *J. Pharm. Biomed. Anal.* 16, 357-366.
- Higuchi, T. and Connors, K.A., 1965. Phase solubility techniques. *Avd. Anal. Chem. Instr.* 4, 117-212.
- Jones, S.P. and Parr, G.D., 1987. Acetoluides as models for studying cyclodextrin inclusion complexes. *Int. J. Pharm.* 36, 223-231.
- Jun, H.N. and Lai, J.W., 1983. Preparation and in vitro dissolution tests of egg albumin microcapsules of nitrofurantoin. *Int. J. Pharm* 16, 65-77.
- Li, S. and Purdy, W.C., 1992. Circular dichroism, ultraviolet, and proton nuclear magnetic resonance spectroscopic studies of the chiral recognition mechanism of β -cyclodextrin. *Anal. Chem.* 641, 405-1412.
- Maggi, L., Massolini, G., de Lorenzi, E., Conte, U. and Caccialanza, G., 1996. Evaluation of stereoselective dissolution of verapamil hydrochloride from matrix tablets press-coated with chiral excipients. *Int. J. Pharm.* 13, 943-52.
- Miya, T., Miyakawa, T., Kawayano, M. and Mivake, Y., 1987. Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds. *J. Chromatogr.* 408, 316-322.
- Mora, M. and Pato, J., 1993. Preparation of egg albumin microparticles for oral applications. *J. Contr. Rel.* 25, 107-113.

- Morley, J., Chapman, I.D., Foster, A., Hoshiko, K. and Mazzoni, L., 1991. Effects of (+) and racemic salbutamol on airway response in the guinea-pig. Br. J. Pharmacol. 104, 295P.
- Oda, Y., Asakawa, N., Kajima, T., Yoshida, Y. and Sato, T., 1991a. Column-switching high-performance liquid chromatography for on-line simultaneous determination and resolution of enantiomers of verapamil and its metabolites in plasma. Pharm. Res. 8, 997-1001.
- Oda, Y., Asakawa, N., Kajima, T., Yoshida, Y. Sato, T., 1991b. On-line determination and resolution of verapamil enantiomers by high-performance liquid chromatography with column-switching. J. Chromatogr. 541, 411-418.
- Pesola, G.R. and Walle, T., 1992. Enantiomeric interaction in the sulfate conjugation of the β_2 -agonist drug albuterol by the human liver. Res. Commun. Chem. Pathol. Pharmacol. 75, 125-128.
- Suedee, R., Brain, K.R. and Heard, C.M., 1997. Enantioselective retardation of rac-propranolol from matrices containing cellulose derivatives. Chirality, 9, 307-312.
- Suzuki, M. and Sasaki, Y., 1979. Inclusion compound of cyclodextrin and azo dyes 1. Methyl orange. Chem. Pharm. Bull. 27, 609-619.
- Szejtli, J. ed., 1982. Cyclodextrins and their inclusion complexes. Budapest, Akademai, Kaido.
- Torrado, S., Torrado, J.J. and Cadorniga, R., 1992. Topical application of albumin microspheres containing vitamin A: Drug release and availability, Int. J. Pharm. 86, 147-152.

Uekama, K. Narisawa, S., Hirayama, F. and Otagiri, M., 1983. Improvement of dissolution and absorption characteristics of benzodiazepines by cyclodextrin complexation. Int. J. Pharm. 16, 327-338.

Vakily, M. and Jamali, F., 1992. Stereoselective dissolution and human pharmacokinetic of tiaprofenic acid. Pharm. Res, 9, S-325.

Vakily, M. and Jamali, F., 1994. Human pharmaceutics of tiaprofenic acid after regular and sustained release formulations: Lack of chiral inversion and stereoselective release. J. Pharm. Sci. 83, 495-499.

Vakily, M., Pasutto, F.M., Daneshtalab, M. and Jamali, F., 1995. Inclusion complexation of heptakis (2,6-di-O-ethyl)- β -cyclodextrin with tiaprofenic acid: pharmacokinetic consequence of a pH-dependent release and stereoselective dissolution. J. Pharm.Sci., 84 1014-19.