

8. เอกสารอ้างอิง

- กรรมการ กาญจนชาตรี, กนกธร ปิยธรรมรัตน์ และนิกร อินทรเจริญ. 2546. ผลของความลึกของ
ระดับน้ำทะเลและขนาดของหอยมุกกล้วยหา (*Pteria penguin*) ต่อการเกิดมุก. ว.สงขลา
นครินทร์ วทท. 25(5) : 659-671.
- จินตนา นักรณนาด. 2536. การเพาะเลี้ยงหอยมุกในประเทศไทย. รายงานเสนอในการประชุมวิชา
การวิทยาศาสตร์และเทคโนโลยีแห่งประเทศไทย ครั้งที่ 19. สงขลา.
- จินตนา นักรณนาด. 2537. “การเพาะเลี้ยงหอยมุกในประเทศไทย”. ประมงเศรษฐกิจ. 3 (31):
63-67.
- สุวรรณ นิคม 2540. การศึกษาการผลิตพันธุ์หอยมุกกระแส[(*Pinctada fucata* (Gould,1850))]
วิทยานิพนธ์ปริญญาโท.บัณฑิตวิทยาลัย มหาวิทยาลัยเกษตรศาสตร์, กรุงเทพมหานคร.
- สุชาติ อุปถัมภ์, มาลียา เครือตรากู, เยาวลักษณ์ จิตราวนวงศ์ และศิริวรรณ จันทเตมีร์. 2538. สังเข
วิทยา. คณะวิทยาศาสตร์ มหาวิทยาลัยมหิดล. กรุงเทพมหานคร. 517 น.
- Alagarswami,K.; Dharmaraj,S.;Vellayudham,T.S.;Chellam,A.;Victor,A.C.C. 1983a. Embryonic
and early larval development of the pearl oyster *Pinctada fucata* (Gould). Proc.Symp.
Coastal Aquac.Mar.Bio.Ass. India. Pt. 2 pp. 598-603.
- Alagarswami,K.; Dharmaraj,S.;Vellayudham,T.S.;Chellam,A.;Victor,A.C.C.;Gandhi, A.D.
1983b. Larval rearing and production of spat of pearl oyster *Pinctada fucata* (Gould).
Aquaculture. 34 : pp287-301.
- Alagarswami,K.; Dharmaraj,S.; Chellam,A;Vellayudham,T.S.1989.Larva and Juvenile rearing
of black- lip pearl oyster, *Pinctada margaritifera* (Linnaeus). Aquaculture.76: pp 43-56.
- Araya-Nunez,O.; Ganning,B.; Bueckle-Ramirez,F. (1995)Embryonic development, larval
culture, and settling of American pearl-oyster (*Pteria sterna*, Gould)spat.CALIF-FISH-
GAME.81(1): pp.10-21.
- Borrero,F.T.(1995).Experimental culture of pearl oysters (*Pinctada imbricata* and *Pteria
colymbus*)(Bivalvia: Pteriidae)on the Caribbean coast of Columbia .Journal World –
Aquaculture.14(1) 261 p.
- Carpenter,K.E. and Niem,V.H. 1998. The living Marine Resources of the Western Central
Pacific. FAO species identification guide for Fishery purposes. (1): 181-183.
- Chen,G.;Chen,J.and Chen,P. 1986. Studies on the natural pearl and shell of *Pinctada martensii*
Journal Marine Drugs Haiyang Yaowu. 5(3) : 4-7

- Choi,Y.H and Chang ,Y. J. 2002. Gametogenic cycle of the transplanted-cultured pearl oyster,*Pinctada fucata martensii* (Bivalvia : Pteriidae)in Korea. Aqauaculture.pp1-15.
- Choi,Y.H and Chang ,Y. J.2003. The influence of cooling rate, developmental stage, and the addition of sugar on cryopreservation of larvae of the pearl oyster *Pinctada fucata martensii* .Department of Aquaculture .46 : pp.190-193.
- Dharmaraj,S.;Vellayudham,T.S.;Chellam,A.;Victor,A.C.C.;Gopinathan,C.P.1991. Hatchery production of pearl oyster spat *Pinctada fucata* Bull.Cent.Mar.Fish.Res.Inst.(Cochin, India)49, Special Publication, 36 pp.
- Doroudi,M.S. 1996. Infestation of pearl oysters by boring and fouling organisms in the northern Persian Gulf. Indian Journal of Marine Sciences. 25(2): 168-169.
- Ellis,S.and Haws, M.(2000).Producing Pearl Using the Black- lip Pearl Oyster(*Pinctada margaritifera*).Center for tropical and Subtropical Aquaculture (CTSA) Aquafarmer Information . 141.8 pp.
- Friedman,K.J and Bell,J.D. 2000. Shorter immersion times increase yields of the blacklip pearl oyster *Pinctada margaritifera* (Line.) from spat collectors in Solomon Island. Aqauaculture. 187(3-4) :pp 299-313.
- Gaytan-Mondragon,I.;Caceres-Martinez,C. and Tobias-Sanchez,M.(1993).Growth of the pearl Oysters *Pinctada mazatlanica* and *Pteria sterna* in different culture at La Paz Bay,Baja California Sur, Mexico.Journal World Aquaculture 24(4):pp.541-546.
- Haws, M. and Ellis,S.(2000).Collecting Black-lip Pearl Oyster Spat. Center for tropical and Subtropical Aquaculture (CTSA) Aquafarmer Information . 144.15 pp.
- Komaru,A.;Matsuda,H.;Yamakawa,T. and Wada, K.T. 1990. Meiosis and fertilization of the Japannese pearl oyster eggs at different temperature observed with a fluorescence microscope.Nippon Suisan Gakkaishi Bull. Jan. Soc. Sci.Fish. 56(3): pp 425-430.
- Laurence,J.G.1996. An investigation into pearl culture in the pinto abalone *Haliotis kamtschatkana* (Jonas) by examining its repaired shell and factors that influence its repair rate.Mai 34(2) : p670.
- Luong-Van,J. 1994.Tropical Australia microalgae as live food for pearl oyster, *Pinctada maxima*,spat.Fisheries research and development. 95 : p 131.

- Masahiro Ito. 1998.Commercial spat production of *Pinctada maxima* at Roko Island hatchery, North-East Australia, with in vitro fertilisation technique.Pearl Oyster information Bulletin. 11 : pp. 3-5.
- Monteforte,-M.;Kappelman-Pina,-E. and Lopez-Espinosa,B.1995. Spat fall of pearl oyster,*Pteria sterna* (gould) on experimental collectors at Bahia de La Paz,South Baja California, Mexico.Aquaculture. 26 (7) : pp 497-511.
- Mazon-Suastegui. (1988). Conditioning and spawning of four bivale species ,fed with artificial diets. Aquaculture . (38):pp4-13.
- Okauchi,M.1990. Food value of *Isochrysis aff. galbana* for the growth of pearl oyster spat.Nippon Suisan Gakkaishi Bull. Jap.Soc.Sci.Fish. 56(8) :1343.
- Rahim-Aldin,S.A.(1990).Histology and histochemistry of the mantle of *Pinctada radiata* and *Pinctada margaritifera* from the Red Sea.Thesis –Abattract 50(1)165 pp.
- Rose,R.A.and Baker,S.B.(1994).Larval and spat culture of the Western Australian silver-or goldlip pearl oyster, *Pinctada maxima* Jamson (Mollusca : Pteriidae) Aquaculture . 126 (1-2):pp35-50.
- Saucedo,P.and Monteforte,M. (1997). In situ growth of pearl oyster *Pinctada mazatlanica* (Hanley 1856)and *Pteria sterna* (Gould 1851)under repopulation condition at Bahia de La Paz, Baja California Sur, Mexico. Aquaculture 28 (5):pp.367-378.
- Southgate,P.C. and Beer,A.C. 1997. Hatchery and early nursery culture of the blacklip pearl oyster (*Pinctada margaritifera*). Journal Shellfish Research. 16(2): 561–567.
- Southgate,P.C. and Rose,R.A. 1997. Fouling animals and their effect on the growth of silver-lip pearl oysters, *Pinctada maxima* (Jameson) in suspended culture. Aquaculture. 153(1-2) : pp 31-40.
- Southgate,P.C. ,Beer,A.C.,Duncan,P.F and Tamburri R. 1998. Assessment of the nutritional value of three species of tropical , dried *Tetraselmis* and a yeast – based diet for larvae of the blacklip pearl oyster, *Pinctada magaritifera*. Aquaculture 162(3-4): 245-259
- Southgate,P.C. 2000. Growth of blacklip pearl oyster (*Pinctada margaritifera*) juveniles using different nursery culture techniques. Aquaculture 187 (1-2) : pp 97-104.
- Sudo, S.; Fujikawa,T.; Nagakura,T.; Okhubo,T.; Sakaguchi,K.; Tanaka,M.; Nakashima,K.; Takahashi. T. 1997. "Structures of mollusc shell framework protein" Journal Nature. (6633): pp.563-564.

- Tanaka,Y.;Kumeta,E.1981. Successful artificial breeding of the silver-lip pearl oyster, *Pinctada maxima* (Jameson).Bull.Natl.Rea.Inst.Aquacult.(Jpn).2 : pp.21-28.
- Taylor,J.J., Rose,R.A. and Southgate,P.C. 1997a. Effects of stocking density on growth and survival of early juvenile silver-lip pearl oyster, *Pinctada maxima* (Jameson), held in suspended nursery culture. Aquaculture. 153(1): 41-49.
- Taylor,J.J., Southgate,P.C., Wing,M.S.and Rose R.A.1997.Assessment of the nutritional value of five species of microalgae for spat of the silverlip pearl oyster ,*Pictada maxima* (mollusca: pteriidae)(Jameson).Asian Fisheries Science. 10 (2) : p 27
- Taylor,J.J., Southgate,P.C. and Rose,R.A. 1998. Effects of mesh covers on the growth and survival of silver-lip pearl oyster (*Pinctada maxima*, Jameson) spat. Aquaculture. 162(3-4) : 243-248.
- Victor,A.C.C.;Chellam,A.;Dharmaraj,S.;Velayudhan,T.S.1995. Manual on pearl oyster seed production, frrming and pearl culture. Bull.Cent.Mar.Fish. Res. Inst.(Cochin, India)63, Special Publication, 53 pp.
- Wallace,R.K.(2001) Cultivating the Eastern Oyster, *Crassostrea virginica*. Southern Regional Aquaculture Center (SRAC) .432. 8 pp.
- Yukihira ,H.;Klumpp,D.W.;Lucas,J.S. (1999).Feeding adaptations of the pearl oysters *Pinctada margaritifera* and *P.maxima* to variations in natural particulates. Marine Ecology.182 :pp.161-173.