

เอกสารอ้างอิง

วรารุณ สุขมาวงศ์ .(2527). คู่มือการฝากครรภ์และการคลอด. พิมพ์ครั้งที่ 12 .คณะ

แพทยศาสตร์ โรงพยาบาลรามาธิบดี. มหาวิทยาลัยมหิดล.

Albers, L. (1998). Midwifery management of pain in labor: The CNM Data Group,

1996. Journal of Nurse-Midwifery, 43(2), 77-82.

Albers, L. L. (1999). The duration of labor in healthy women. Journal of

Perinatology, 19(2), 114-119.

Allen, S. (1998). A qualitative analysis of the process, mediating variables and

impact of traumatic childbirth. Journal of Reproductive and Infant Psychology, 16, 107-

131.

Ballard, C. G., Stanley, A. K., & Brockington, I. F. (1995). Post-traumatic stress

disorder (PTSD) after childbirth. British Journal of Psychiatric, 166(4), 525-528.

Blackburn, S. T., & Loper, D. L. (1992). Maternal, fetal, and neonatal physiology:

A clinical perspective. Philadelphia: W.B. Saunders Company.

Bonica, J. J. (1994). Labor pain. In P. D. Wall & R. Melzack (Eds.), Textbook of

pain (3rd ed., pp. 615-640). New York: Churchill Livingstone.

Bonica, J. J., & McDonald, J. S. (1990). The pain of childbirth. In J. J. Bonica

(Ed.), The management of pain volume II (2nd ed., pp. 1313-1343). Philadelphia: Lea &

Febiger.

Bloom, S. L., McIntire, D. D., Kelly, M. A., Beimer, H. L., Burpo, R. H., Garcia, M.

A. et al. (1998). Lack of effect of walking on labor and delivery. The New England

Journal of Medicine, 339(2), 76-79.

Briggs, M., & Closs, J. S. (1999). A descriptive study of the use of visual

analogue scales and verbal rating scales for the assessment of postoperative pain in

orthopedic patients. Journal of Pain and Symptom Management, 18(6), 438-446.

Briggs, G. G., Freeman, R. K., Yaffe, S. J. (1994). A reference guide to fetal and

neonatal risk: Drugs in pregnancy and lactation. Philadelphia: Williams & Wilkins.

Brook, E. L. (1984). Soothing music during the active phase of labor: Physiologic

effect on mother and infant. Unpublished master's thesis, University of Florida, Florida.

Brown, S. T., Campbell, D., & Kurtz, A. (1989). Characteristics of labor pain at

two stages of cervical dilation. Pain, 38(3), 289-295.

Burroughs, A. (1992). Maternity nursing: An introductory text (6th ed.).

Philadelphia:W. B. Saunders Company.

Burroughs, A., & Leifer, G. (2001). Maternity Nursing: An introductory text (8th

ed.). Philadelphia: W.B. Saunders company.

Callaghan, C. C. (1996). Pain, music creativity and music therapy in palliative

care. Complementary Medicine International, 3(2), 43-48.

Chlan, L. (1997). Music intervention. In B. M. Dossey (ed. pp 58-68). Core

curriculum for holistic nursing. Gaithersburg, MA: An Aspen.

Clark, M. E., McCorkle, R.R., & Williams, S.B. (1981). Music therapy-assisted

labor and delivery. Journal of Music Therapy, 18(2), 88-100.

Cohen, J. (1988). Statistical power analysis for the behavioral sciences (2nd ed.).

Hillsdale, NJ: Lawrence Erlbaum Associates.

Cook, T. D., & Campbell, D. T. (1979). Quasi-experimental: Design & Analysis

issues for field settings. Boston:Houghton Mifflin Company.

Dannenbring, D., Stevens, M. J., & House, A. E. (1997). Predictors of childbirth

pain and maternal satisfaction. Journal of Behavioral Medicine, 20(2), 127-142.

de Jong, P. R., Johanson, R. B., Baxen, P., Adrians, V. D., van der Westhuisen,

S., & Jones, P. W. (1997). Randomised trial comparing the upright and supine positions for the second stage of labour. British Journal of Obstetrics and Gynaecology, 104(5), 567-571.

Dick-Read, G. (1972). Childbirth without fear. New York: Harper & Row.

Fairlie, F. M., Marshall, L., Walker, J. J., & Elbourne, D. (1999). Intramuscular

opioids for maternal pain relief in labour: A randomized controlled trial comparing

pethidine with diamorphine. British Journal of Obstetric Gynaecology, 106(11), 1191-1187.

Faucher, M. A., & Brucker, M. C. (2000). Intrapartum pain: Pharmacological

management. Journal of Obstetric, Gynecologic, and Neonatal Nursing, 29(2), 169-180.

Fenwick, L., & Simkin, P. (1987). Maternal positioning to prevent or alleviate

dystocia in labor. Clinical Obstetrics and Gynecology, 30(1), 83-89.

Flynn, A. M., Kelly, J., Hollins, G., & Lynch, P. F. (1978). Ambulation in labour.

British Medical Journal, 2, 591-593.

Fones, C. (1996). Posttraumatic stress disorder occurring after painful childbirth.

Journal of Nervous and Mental Disease, 184(3), 195-196.

Fridh, G., Kopare, T., Gaston-Johansson, F., & Norvell, K.T. (1988). Factors

associated with more intense labor pain. Research in Nursing & Health, 11(2), 117-124.

Friedman, E. A. (1978). Labor: Clinical evaluation and management (2nd ed.).

New York: Appleton-Century-Crofts.

Fujinawa, A., Kawai, I., Ohashi, H., & Kimura, S. (1977). A case of musicogenic

epilepsy. Folia Psychiatrica et Neurologica Japonica, 31(3), 463-472. =

Gaston-Johansson, F., & Gustafsson, M. (1990). Rheumatoid

arthritis: Determination of pain characteristics and comparison of RAI and VAS in its

measurement. Pain, 41(1), 35-40.

Geden, E.A., Lower, M., Beattie, S., & Beck, N. (1989). Effects of music and

imagery on physiologic and self-report of analogued labor pain. Nursing Research,

38(1), 37-41.

Gonzalez, C. E. (1989). The music therapy-assisted childbirth program: A study

evaluation. Pre-and Peri-Natal Psychology, 4(2), 111-124.

Golub, M. S. (1996). Labor analgesia and infant brain development.

Pharmacology Biochemistry and Behavior, 55(4), 619-628.

Green, J. M. (1993). Expectations and experiences of pain in labor: Findings from a large prospective study. Birth, 20(2), 65-72.

Hanser, S. B. , Larson, S. C., & O'Connell, A. S. (1983). The effect of music on relaxation of expectant mothers during labor. Journal of Music Therapy, 20(2), 50-58.

Hanson, L. (1998). Second-stage positioning in nurse-midwifery practices.

Journal of Nurse-Midwifery, 43 (5), 320-325.

Harvey, P. (2001). Yoga for every body. New York: The Reader's Digest

Association, Inc.

Heardman, H. (1982). Relaxation and exercise for childbirth. New York: Churchill

Livingstone.

Herbst, A., Wolner-Hanssen, P., & Ingemarsson, I. (1995). Risk factors for fever in labor. Obstetrics and Gynecology, 86(5), 790-794.

Heyns, O.S. (1959). Abdominal decompression in the first stage of labour.

Journal of Obstetrics & Gynaecology of the British Empire, 66(2),220 -228.

- Hodnett, E. (1996). Nursing support of the laboring women. Journal of Obstetric, Gynaecologic, and Neonatal Nursing, 25(3), 257-264.
- Huskisson, E. C. (1983). Visual Analogue Scales. In R. Melzack (Ed.), Pain measurement and assessment (pp. 33-37). New York: Raven Press.
- Jacobson, H. (1991). Protecting the back during pregnancy. AAOHN Journal, 39(6), 286-291.
- Jensen, M. P., Karoly, P., & Braver, S. (1986). The measurement of clinical pain intensity: A comparison of six methods. Pain, 27(1), 117-126.
- Keiss, H. O. (1996). Statistical concepts for the behavioral sciences (2nd ed.). Boston: Allyn and Bacon.
- Kendall-Tackett, K. A., & Kantor, G. K. (1993). Postpartum depression: A comprehensive approach for nurses. Newbury Park, CA: Sage.
- King, T. (1997). Epidural anesthesia in labor. Journal of Nurse-Midwifery, 42(5), 377-387.

- Kusolleartjariya, S. (1997). A study of comparison between folk's music and classical music to decreased labour pain. Thailand Journal of Health Promotion and Environmental Health, 20(3), 32-42.
- Lanahan, C. C. (1978). Variables affecting pain perception during labor. In L. K. McNall & J. T. Galeener (Eds.), Current practice in obstetric and gynecologic nursing: Volume two. Saint Louis, MO: The C. V. Mosby Company.
- Lee, K. A., & Kieckhefer, G. M. (1989). Measuring human responses using visual analogue scales. Western Journal of Nursing Research, 11(1), 128- 132.
- Limprasirt, D. (1996). Effect of music on pain coping during labour in primiparas. Unpublished master's thesis, Chiang Mai University, Chiang Mai, Thailand.
- Lipsey, M. W. (1990). Design sensitivity: Statistical power for experimental research. Newbury Park, CA: Sage.
- Lowe, N. K. (1991). Critical predictors of sensory and affective pain during four phases of labor. Journal of Psychosomatic Obstetric and Gynaecology, 12(3), 193-208.
- Macario, A., Scibetta, W. C., Navarro, J., & Riley, E. (2000). Analgesia for labor pain: A cost model. Anesthesiology, 92 (3), 841-850.

Martin, E. J. (2002). Intrapartum management modules: A perinatal education

program. Philadelphia: Lippincott Williams & Wilkins.

Masten, Y. (1997). Obstetric nursing (2nd ed.). Aurora, CO: Skidmore-Roth.

McGuire, D. B. (1984). The measurement of clinical pain. Nursing Research,

33(3), 152-156.

Melzack, R. (1987). The short-form McGill Pain Questionnaire. Pain, 23(4), 345-

356.

Melzack, R., Bélanger, E., & Lacroix, R. (1991). Labor pain: Effect of maternal

position on front and back pain. Journal of Pain and Symptom Management, 6(8), 476-

480.

Melzack, R., Kinch, R., Dobkin, P., Lebrun, M., & Trenzer, P. (1984). Severity of

labour pain: Influence of physical as well as psychologic variables. Canadian Medical

Association Journal, 125(5), 357-363.

Melzack, R., Taenzer, P., Feldman, P., & Kinch, R. A. (1981). Labour is still

painful after prepared childbirth training. Canadian Medical Association Journal, 125(4),

357-363.

Melzack, R., & Torgerson, W.S. (1971). On the language of pain. Anesthesiology,

34(1), 50-59.

Melzack, R., & Wall, P.D. (1965). Pain mechanisms: A new theory. Science,

150(3699), 971-979.

Melzack, R., & Wall, P. D. (1996). Pain mechanisms: A new theory. Pain Forum,

5(1), 3-11.

Méndez-Bauer, C., Arroyo, J., Ramos, C. G., Lavilla, M., Izquierdo, F., Elizaga,

V., et al., (1975). Merriam-Webster's Collegiate Dictionary (10th), Springfield, MA:

Merriam-Webster.

Merskey, H., Bonica, J.J., Carmon, A. Dubner, R., Kerr, F.W.L., Lindblom, U., et

al. (1979). Pain terms:A list with definitions and notes on usage. Pain, 6(3), 249-252.

Molina, F. J., Solá, P. A., López, E., & Pires, C. (1997). Pain in the first stage of

labor: Relationship with the patient's position. Journal of Pain and Symptom

Management, 13(2), 98-103.

Niven, C. A. (1992). Psychological care for families: Before, during and after

birth. Oxford, Great Britain: Butterworth-Heinemann Ltd.

Niven, C., & Gijsbers, K. (1984a). A study of labor pain using the McGill Pain

Questionnaire. Social Science Medicine, 19(12), 1374-1351.

Niven, C. A., & Gijsbers, K. (1984b). Obstetric and non-obstetric factors related

to labour pain. Journal of Reproductive and Infant Psychology, 2, 61-78.

Olofsson, C., Ekblom, A., Ekman-Ordeberg, G., Hjelm, A., & Irestedt, L. (1996).

Lack of analgesic effect of systemically administered morphine or pethidine on labour

pain. British Journal of Gynaecology, 103(10), 968-972.

Olds, S. B., London, M. L., & Ladewig, P.A.W. (2000). Maternal-newborn nursing:

A family and community-based approach (6th ed.). Upper Saddle River, NJ: Prentice

Hall Health.

Pasternak, J. F., & Gorey, M. T. (1998). The syndrome of acute near-total

intrauterine asphyxia in the term infant. Pediatric Neurology, 18(5), 391-398.

Phumdoung, S., & Good, M. (2003). Music reduces sensation and distress of

labor pain. Pain Management Nursing, 4(2), 54-61.

Preston, R., Crosby, E.T., Kotarba, D., Dudas, H., Elliot, R. D. (1993). Maternal positioning affects fetal heart rate changes after epidural analgesia for labour. Canadian Journal of Anesthesia, 40 (12), 1136-1141.

Price, D. D., McGrath, P. A., Rafii, A., & Buckingham, B. (1983). The validity of visual analogues scales as ratio scale measures for chronic and experimental pain. Pain, 17(1),45-56.

Quine, L., Rutter, D. R., & Gowen, S. (1993). Women's satisfaction with the quality of the childbirth experience: a prospective study of social and psychological predictors. Journal of Reproductive and Infant Psychology, 11, 107-113.

Ranta, P., Jouppila, P., & Jouppila, R. (1996). The intensity of labor pain in grand multiparas. Acta Obstetricia et Gynecologica Scandinavica, 75 (3), 250-254.

Ranta, P., Jouppila, P., Spalding, M., & Jouppila, R. (1995). The effect of maternal obesity on labour and labour pain. Anesthesia, 50(4), 322-326.

Ranta, P., Spalding, M., Kangas-Saarela, T., Jokela, R., Hollmen, A., Jouppila, P., et al. (1995). Maternal expectations and experiences of labour pain-options of 1091 Finnish parturients. Acta Obstetricia et Gynecologica Scandinavica, 39(1), 60-66.

Revill, S. I., Robinson, M., Rosen, M., & Hogg, M. I. J. (1976). The reliability of a linear analogue for evaluating pain. Anesthesia, 31(9), 1191-1198.

Reynolds, J. L. (1997). Post-traumatic stress disorder after childbirth: The phenomenon of traumatic birth. Canadian Medical Association, 156(6), 831-835.

Roberts, J., Malasanos, L., & Mendez-Bauer, C. (1981). Maternal positions in labor: Analysis in relation to comfort and efficiency. Birth Defects: Original Article Series, 17(6), 97-128.

Robertson, A. (no year) Preparing for birth: Background notes for pre-natal classes. ACE Graphic.

Rohrbach, A., Viehweg, B., Kuhnert, I., Koster, A., & König, F. (2001). [Effect of peridural analgesia on labor progress.] Anaesthesiol Reanim, 26(2), 39-43.

Rosen, M. G., & Peisner, D. B. (1987). Effect of amniotic membrane rupture on length of labor. Obstetrics & Gynecology, 70(4), 604-607.

Ryding, E. L. (1993). Investigation of 33 women who demanded a cesarean section for personal reasons. Acta Obstetrica et Gynecologica Scandinavica, 72(4), 280-285.

Ryding, E. L., Wijma, B., Wijma, K., & Rydhström, H. (1998). Fear of childbirth

during pregnancy may increase the risk of emergency cesarean section. Acta

Obstetrica et Gynecologica Scandinavica, 77(5), 542-547.

Saisto, T., Ylikorkala, O., & Halmesmaki, E. (1999). Factors associated with fear

of delivery in second pregnancies. Obstetrics and Gynecology, 94(5), 679-682.

Sammons, L. N. (1984). The use of music by women during childbirth. Journal of

Nurse-Midwifery, 29(4), 266-270.

Scott, J., & Huskisson, E. C. (1976). Graphic representation of pain. Pain, 2(2),

175-184.

Sheiner, E., Sheiner, E. K., & Shoham-Vardi, I. (1998). The relationship between

parity and labor pain. International Journal of Gynecology & Obstetrics, 63(3), 287-288.

Sherwen, L. N., Scoloveno, M. A., & Weingarten, C. T. (1999). Nursing care of the

childbearing family (3rd ed.). Norwalk, CT: Appleton & Lange.

Simkin, P. (1991). Non-pharmacological methods of pain relief during labour. In

I. Chalmers, M. Enkin, M. Keirse. Effective care in pregnancy & Childbirth (pp. 893-912).

Oxford: Oxford University Press.

Sjögren, B., & Thomassen, P. (1997). Obstetric outcome in 100 women with severe anxiety over childbirth. Acta Obstetricia et Gynecologica Scandinavica, 76(10), 948-952.

Snaz D. C. (1998). [Advantages and disadvantages of analgesia during childbirth.] Revista De Enfermeria, 21(244), 17-21.

Stewart, D. E. (1982). Psychiatric symptoms following attempted natural childbirth. Canadian Medical Association Journal, 127(8), 713-716.

Stewart, P., & Calder, A. A. (1984). Posture in labour: patients' choice and its effect on performance. British Journal of Obstetrics and Gynaecology, 91, 1091-1095.

Sweet, B. R. (1997). Mayes' midwifery: A textbook for midwives (12th ed.). Philadelphia: Baillière Tindall.

Thorp, J. A., & Breedlove, G. (1996). Epidural analgesia in labor: An evaluation of risks and benefits. Birth, 23(2), 63-81.

Thune-Larsen, K-B., & Møeller-Pedersen, K. (1988). Childbirth experience and postpartum emotional disturbance. Journal of Reproductive and Infant Psychology, 6, 229-240.

Troy, N. W. (2003). Is the significance of postpartum fatigue. MCN, 1(4), 252-257.

Turley, G. M. (2000). Essential forces and factors in labor. In S. Mattson & J. E.

Smith (Eds.), Core curriculum for maternal-newborn nursing (2nd ed., pp. 204-240).

Philadelphia: W.B. Saunders Company.

Varma, T. R., & Curzen, P. (1973). The effects of abdominal decompression on

pregnancy complicated by the small-for-dates fetus. The Journal of Obstetrics and

Gynaecology, 80, 1086-1094.

Waldenström, U. (1999). Experience of labor and birth in 111 women. Journal of

Psychosomatic Research, 47(5), 471-482.

Waldenström, U., Bergman, V., & Vasell, G. (1996). The complexity of labor pain:

experiences of 278 women. Journal of Psychosomatic Obstetrics and Gynaecology,

17(4), 215-228.

Waldenström, U., & Gottvall, K. (1991). A randomized trial of birthing stool or

conventional semirecumbent position for second-stage labor. Birth, 18(1), 5-10.

- Webb, R. J., & Kantor, G. S. (1992). Obstetrical epidural anaesthesia in a rural Canadian hospital. Canadian Journal of Anaesthesia, 39(4), 390-393.
- Weisenberg, M., & Caspi, Z. (1989). Cultural and educational influences on pain of childbirth. Journal of Pain and Symptom Management, 4(1), 13-19.
- Wuitchik, M., Bakal, D., & Lipshitz, J. (1989). The clinical significance of pain and cognitive activity in latent labor. Obstetrics and Gynecology, 73(1), 35-42.
- Wuitchik, M., Hesson, K., & Bakal, D. A. (1990). Perinatal predictors of pain and distress during labor. Birth, 17(4), 186-191.
-