

เอกสารอ้างอิง

- Ader, J-L., Tran-Van, T. and Praddaude, F. (1987). Reduced urinary kallikrein activity in rats developing spontaneous hypertension. *Am. J. Physiol.* **252(21)**:F964—F969.
- Beierwaltes, W.H., Arendshorst, W.J. and Klemmer, P.J. (1982). Electrolyte and water balance in young spontaneously hypertensive rats. *Hypertension.* **4**:908—915.
- Cervenka, L., Wang, C-T. and Navar, L.G. (1998). Effects of acute AT1 receptor blockade by candesartan on arterial pressure and renal function in rats. *Am. J. Physiol.* **274(43)**:F940—F945.
- Dietz, R., Schomig, A., Haebara, H., Mann, J.F.E., Rascher, W., Luth, J.B., Grunherz, N. and Gross, F. (1978). Studies on the pathogenesis of spontaneous hypertension of rats. *Cir. Res.* **43(1)(Suppl I)**:I98—1106.
- Gibson, R.E., Thorpe, H.H., Cartwright, M.E., Frank, J.D., Schorn, T.W. Bunting, P.B. and Siegl, P.K.S. (1991). Angiotensin II receptor subtypes in renal cortex of rats and rhesus monkeys. *Am. J. Physiol.* **261**:F512—F518.
- Györy, A.Z. (1971). Reexamination of the split oil droplet method as applied to kidney tubules. *Pflugers Arch.* **324**:328—343.
- Györy, A.Z., Lingard, J.M. and Young, J.A. (1974). Relation between active sodium transport and distance along the proximal convolutions of rat nephrons: evidence for homogeneity of sodium transport. *Pflugers Arch.* **348**:205—210.
- Harrap, S.B. and Doyle, A.E. (1985). Total body sodium in immature spontaneously hypertensive and Wistar Kyoto rats. *Clin. Exp. Pharmacol. Physiol.* **12**:315—318.
- Harris, P.J. and Young, J.A. (1977). Dose-dependent stimulation and inhibition of proximal tubular sodium reabsorption by angiotensin II in the rat kidney. *Pflugers Arch.* **367**:295—297.
- Harris, P.J., Cullinan, M., Thomas, D. and Morgan, T.O. (1987). Digital image capture and analysis for split-drop micropuncture. *Pflugers Arch.* **408**:615—618.
- Harris, P.J., Hiranyachattada, S., Antoine, A.M., Walker, L, Reilly, A.M. and Eitle, E. (1996). Regulation of renal tubular sodium transport by angiotensin II and atrial natriuretic factor. *Clin. Exp. Pharmacol. and Physiol.* Suppl 3:S112-S118.

- Herlitz, H., Ricksten, S.E., Lundin, S., Thoern, P., Aurell, M. and Berglund, G. (1983). Renal denervation and sodium balance in young spontaneously hypertensive rats. *Renal Physiol.* **6**:145–150.
- Hiranyachattada, S. and Harris, P.J. (1996). Modulation by locally produced luminal angiotensin II of proximal tubular sodium reabsorption via an AT₁ receptor. *British J. Pharmacol.* **119**:617-618.
- Liu, F-Y. and Cogan, M.G. (1986). Axial heterogeneity of bicarbonate, chloride and water transport in the rat proximal convoluted tubule. Effects of changes in luminal flow rate and of alkalemia. *J. Clin. Invest.* **78**:1547–1557.
- Sato, K. (1975). Reevaluation of micropuncture techniques: some of the factors which affect the rate of fluid absorption by the proximal tubule. In: *Biochemical Aspects of Renal Function: Current Problems in Clinical Biochemistry*, edited by Angielski, S. and Dubach, U.C. Hans Huber Publishers, Bern Stuttgart Vienna. **4**:175–187.
- Schuster, V.L., Kokko, J.P. and Jacobson, H.R. (1984). Angiotensin II directly stimulated sodium transport in rabbit proximal convoluted tubules. *J. Clin. Invest.* **73**:507–515.
- Seikaly, M.G., Arant, B.S., Jr. and Seney, F.D., Jr. (1990). Endogenous angiotensin concentration in specific intrarenal fluid compartment of the rat. *J. Clin. Invest.* **86**:1352–1357.
- Shibouta, Y., Inada, Y., Ojima, M., Wada, T., Noda, M., Sanada, T., Kubo, K., Kohara, Y., Naka, T. and Nishikawa, K. (1993). Pharmacological profile of a highly potent and long-acting angiotensin II receptor antagonist, 2-ethoxy-1-[[2'-(1H-tetrazol-5-yl) biphenyl-4-yl]-1H-benzimidazol-7-carboxylic acid (CV11974), and its prodrug, (±)-1-(cyclohexyloxycarbonyloxy)-ethyl 2-ethoxy-1-[[2'-(1H-tetrazol-5-yl)biphenyl-4-yl] methyl]-1-H-benzimidazol-7-carboxylate (TCV-116). *J. Pharm. Exp. Ther.* **266**(1):114-120.
- Smart, M.L., Hiranyachattada, S. and Harris, P.J. (1999). Effects of angiotensin II receptor blockade on proximal fluid uptake in the rat kidney. *British J. Pharmacol.* **126**:697-700.

- Spinelli, F. and Walther, A. (1979). Modulation by prostaglandins of angiotensin II-induced stimulation of sodium transport in the proximal tubule of the rat. In: *Hormonal Control of Epithelial Transport*, Vol 85, Paris, INSERM, pp. 273–278.
- Spitzer, A. and Windhager, E.E. (1972). Continuous *in vivo* perfusion of the postglomerular capillary network in superficial rat kidney cortex. *Yale J. Biol. Med.* **45**:307–311.
- Thomas, D., Harris, P.J. and Morgan, T.O. (1990). Altered responsiveness of proximal tubule fluid reabsorption to peritubular angiotensin II in spontaneously hypertensive rats. *J. Hypertens.* **8**:407–410.
- Wang, T. and Chan, Y.L. (1987). Evidence for functional angiotensin receptors on the apical side of the rat proximal convoluted tubule. *Kidney Int.* **31**:291–298.
- Wang, T. and Chan, Y.L. (1991). The role of phosphoinositide turnover in mediating the biphasic effect of angiotensin II on renal tubular transport. *J. Pharmacol. Exp. Ther.* **256**:309–317.
- Wiederholt, M., Langer, K.H., Thosenes, W. and Hierholzer, K. (1968). Funktionelle und morphologische untersuchungen am proximalen und distalen konvolut der rattenniere zur methode der gespalten ousaule (split droplet method). *Arch. Ges. Physiol.* **302**:166–191.
- Wong, P.C., Barnes, T.B. Chui, A.T. and Timmermans, P.B.M.W.M. (1991). Losartan (DuP753), an orally-active nonpeptide angiotensin II receptor antagonist. *Cardiovasc. Drug Rev.* **9**:317–339.
- Xie, M-H., Liu, F-Y., Wong, P.C., Timmermans, P.B.M.W.M. and Cogan, M.G. (1990). Proximal nephron and renal effects of DuP753, a nonpeptide angiotensin II receptor antagonist. *Kidney Int.* **38**:473–479.
- Zhuo, J., Song, K., Abdelrahman, A. and Mendelsohn, F.A.O. (1994). Blockade by intravenous losartan of AT₁ angiotensin II receptors in rat brain, kidney and adrenals demonstrated by *in vitro* autoradiography. *Clin. Exp. Pharmacol. Physiol.* **21**:557–567.