

## เอกสารอ้างอิง

- Akahane, Y. and Shimisu, Y. 1989. Effects of pH and sodium chloride on the water holding capacity surimi and its gel. *Nippon Suisan Gakkaishi*. 55: 1827-1832.
- Almas, K.A. 1981. Chemistry and microbiology of fish processing. Dept. of Biochemistry, Norwegian Inst. of Technol., Univ. of Trondheim. Norway.
- Alvarez, C., Couso, I., and Tejada, M. 1995. Sardine surimi gel as affected by salt concentration, blending, heat treatment and moisture. *J. Food Sci.* 60: 622-626.
- Aguilar, R.P., Sanchez, M., and Burgueno, M.r. 2000. Postmortem biochemical and functional characteristic of monterey sardine muscle stored at 0° C. *J. Food Sci.* 65:40-47.
- Ang, J.F., and Haltin, H.O. 1989. Denaturation of cod myosin during freezing after modification with formaldehyde. *J. Food Sci.* 54:814-818.
- A.O.A.C. 1990. Official Method of Analysis of the Association of Official Analytical chemists 14<sup>th</sup> ed. Virginia: The Association of Official Analytical Chemists,
- Asghar, A., Samejima, K., and Yasui, T. 1985. Functionality of muscle proteins in gelation mechanisms of structured meat products. *CRC Crit. Rev. Food Sci. Nutr.* 22:27-28.

- Babbitt, J.K. and Reppond, K.D. 1988. Factors affecting the gel properties of surimi. *J. Food Sci.* 53:965-966.
- Beas, V.E., Wagner, J.R., and Anon, M.C. 1991. Thermal denaturation in fish muscle proteins during gelling : effect of spawning condition. *J. Food Sci.* 53:281-284.
- Bechtel, P.J. 1986. Muscle development and contractile proteins. In *Muscle as Food*. P.J. Bechtel(Ed.), p.2-31. Academic Press, Inc., Orlando, FL.
- Benjakul, S., and Bauer, F. 2000. Physicochemical and enzymatic changes of cod muscle proteins subjected to different freeze-thaw cycled. *J. Sci Food Agric.* 80:1143-1150.
- Benjakul, S., and Bauer, F. 2001. Biochemical and physicochemical changes in catfish (*Silurus glanis Linne*) muscle an influenced by different freeze-thaw cycles. *Food Chem.* 72:207-217.
- Bligh, E.G. and Dyer, W.J. 1959. A rapid method of total lipid extraction and purification. *Can. J.Biochem. Physiol.* 37:911-917.
- Careche, M. and Li-Chan, E.C.V. 1997. Structural changes in cod myosin after modification with formaldehyde on frozen storage. *J. Food Sci.* 62:717-723.

- Casserly, U., Mooney, M., and Troy, D. 2000 Standardisation and application of semi-quantitative SDS-PAGE method for measurement of myofibrillar protein fragments in bovine *longissimus* muscle. *Food Chem.* 69:379-385.
- Chang, C.C. and Regenstein, J.M. 1997. Water uptake protein solubility and Protein changes of cod mince stored on ice as affected by polyphosphates. *J. Food Sci.* 62:305-309.
- Chan, J.K., Gill, T.A., Thompson, J.W., and Singer, D.S. 1995. Herring surimi during low temperature setting physicochemical and textural properties. *J. Food Sci.* 60:1248-1253.
- Chang-Lee, M.V., Lampila, L.E., and Crawford, D.L. 1990. Yield and composition of surimi from Pacific whiting (*Merluccius productus*) and the effect of various protein additive on gel strength. *J. Food Sci.* 55:83-86.
- Cheng, C.S., Hamann, D.D., and Webb, N.B. 1979. Effect of thermal processing on minced fish gel texture. *J. Food Sci.* 44:1080-1086.
- Chen, H.H. 1995. Thermal stability and gel-forming ability of shark muscle as related to ionic strength. *J. Food Sci.* 60:1237-1240.
- Connell, J.J. 1975. The role of formaldehyde as a protein crosslinking agent acting during the frozen storage of cod. *J. Sci Food Agric.* 26:1925-1929.
- Davies, J.R., Bardsley, R.G., and Ledward, D.A. 1988. Myosin thermal stability in fish muscle. *J. Sci Food Agric.* 45:69-78.

- Douglas-Swarz, M. and Lee, C.M. 1988. Comparison of the thermostability of red hake and Alaska pollock surimi during processing. *J. Food Sci.* 53: 1347-1351.
- Dziezak, J.D. 1990. Phosphates improve many foods. *Food Technol.* 44: 80-92.
- Foegeding, E.A., Allen, C.E., and Dayton, W.R. 1986. Effect of heating rate on thermally formed myosin fibrinogen and albumin gels. *J. Food Sci.* 51:104-108.
- Gill, T.A., Keith, R.A., and Smith, B. 1979. Textural deterioration of red hake and haddock muscle in frozen storage as related to chemical parameters and changes in the myofibrillar protein. *J. Food Sci.* 44:666-667.
- Gomez-Guillen, C., Borderias A.J., and Montero, P. 1997. Thermal gelation properties of two different composition sardine (*Sardina pilchardus*) muscles with addition of non-muscle proteins and hydrocolloids. *Food Chem.* 58:81-87.
- Gomez-Guillen, C., Sola, S. T., and Montero, P. 1997. Influence of added salt and non-muscle proteins on the rheology and ultrastructure of gels made from minced flesh of sardine (*Sardina pilchardus*). *Food Chem.* 58:193-202.
- Hashimoto, A., Kobayashi, A., and Arai, K. 1982. Thermostability of fish myofibrillar Ca-ATPase and adaptation to environmental temperature. *Nippon Suisan Gakkaishi.* 57:747

- Hashimoto, K., Watabe, S., Kono, M., and Shiro, K. 1979. Muscle protein composition of sardine and mackerel. *Bull. Jpn. Soc. Sci. Fish.* 45:1435-1441.
- Hastings, R.J., Rodger, G.W., Park, R., Matthews, A.D., and Anderson, E.M. 1985. Differential scanning calorimetry of fish muscle: the effect of processing and species variation. *J. Food Sci.* 50:503-510.
- Hastings, R.J., Keay, J.N., and Young, K.W. 1990. The properties of surimi and kamaboko gels from nine British species of fish. *Int. J. Food Sci. Technol.* 25:281-283.
- Henson, L.S. and Kowalewski, K.M. 1992. Use of phosphates in seafood. *INFOFISH International.* 5:52-54.
- Huidobro, A. and Tejada, M. 1993. Emulsifying capacity of fish mince from several species during frozen storage. *J. Sci. Food Agric.* 61: 333-338.
- Jiang, S.T., Hsieh, J.e., Ho, M.L., and Chung, Y.C. 2000. Microbial transglutaminase affects gel properties of golden threadfin bream and pollack surimi. *J. Food Sci.* 65:694-699.
- Jiang, S.T., Hwang, D.C. and Chen, C.S. 1988. Effect of storage temperatures on the formation of disulfides and denaturation of milk fish actomyosin (*Chanos chanos*). *J. Food Sci.* 53:1333-1335.

- Jiang, S.T. and Lee, T.C. 1985. Changes in free amino acids and denaturation of fish muscle during frozen storage. *J. Agric Food Chem.* 33: 839-844.
- Kim, B.Y. Hamann, D.D. Lanier, T.C. and Wu, M.C. 1986. Effects of freeze-thaw abuse on the viscosity and gel-forming properties of surimi from two species. *J. Food Sci.* 51:951-956.
- Kim, J.M. and Lee, C.M. 1987. Effect of starch on textural properties of surimi gel. *J. Food Sci.* 52:722-725.
- Koning, A.J. and Mol, T.H. 1991. Quantitative quality tests for frozen fish : soluble protein and free fatty acids content as quality criteria for hake (*Merluccius capensis*) stored at  $-18^{\circ}\text{C}$ . *J. Sci Food Agric.* 54:449-458.
- Konno, K. 1992. Suppression of thermal denaturation of myosin subfragment of Alaska pollack (*Theragra chacogramma*) by sorbitol and accelerated inactivation by pyrophosphate. *J. Food Sci.* 57:261-264.
- Konno, K. Yamadodera, K. and Kiuchi, H. 1997. Solubilization of fish muscle myosin by sorbitol. *J. Food Sci.* 62: 980-984.
- Krueger, D.J. and Fennema, O.R. 1989. Effect of chemical additives on toughening of fillets of frozen Alaska pollack. *J. Food Sci.* 54:1101-1106.
- Kumazawa, Y., Numazawa, T., Seuro, K., and Motoki, M. 1995. Suppression of surimi gel setting by transglutaminase inhibitors. *J. Food Sci.* 60:715-717.

- Lablanc, E.L. and Lablanc, R. 1992. Determination of hydrophobicity and reactive groups in proteins of cod (*Gadus morhua*) muscle during frozen storage. *Food Chem.* 43: 3-11.
- Laemmli, U.K. 1970. Cleavage of structural protein during the assembly of head bacteriophage T4. *Nature.* 277: 680-685.
- Lanier, T.C., Lin, T.S., Liu, Y.M., and Hamann, D.D. 1982. Heat gelation properties of actomyosin and surimi prepared from Atlantic croaker. *J. Food Sci.* 47: 1921-1925.
- Lanier, T.C 1986. Functional properties of surimi. *Food Technol.* 40 (3):107-108.
- Lee, C.M. 1984. Surimi process technology. *Food Technol.* 38(11):69-70.
- Lou, X., Wang, C., Xion, Y.L., Wang, B., and Mims, S.D. 2000 . Gelation characteristics of peddlefish (*polyodan spathula*) surimi under different heating condition. *J. Food Sci.* 65: 394-398.
- Love, R. M. 1992. Biochemical dynamics and the quality of fresh and frozen fish. In *Fish processing Tecnology*. G. M. Hall (Ed) pp. 1-30. Blackie Academic and processional. New York.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. 1951. Protein measurement with the folin phenol reagent. *J. Biol Chem.* 193:265-275.

- MacDonald, G.A., and Lanier, T.C. 1991. Carbohydrated as cryoprotectants for meat and surimi. *Food Technol.* 45(4):150-155.
- MacDonald ,G.A., and Lanier, T.C. 1994. Actomyosin stabilization to freeze-thaw and heat denaturation by lactate salts. *J. Food Sci.* 59: 101-105.
- MacDonald, G.A., Sterven, J., and Lanier, T. C. 1994. Characterization of New Zealand hoki and southern blue whiting surimi compared to Alaska pollock surimi. *J. Aqua. Food prod. Technol.* 3(1):19-38.
- Mackie, J.M. 1994. Fish protein. In *New and developing sources of Food proteins.* B.J.F. Hudson(Ed.) 95-114. Chamman and Hall, London.
- Matsumoto, J.J. 1980. Chemical deterioration of muscle proteins during frozen storage. In *Chemical Deterioration of Proteins* (Ed. by J.R. Whitaker, and M. Fujimaki), p.95 *Advances in Chemistry Series.* No.123. American Chemical Society Washington.DC.
- Min, T.S., Chung, N.M., Fujiwara, T.,Kuang, H.K., and Hasegawa, H. 1987. *Handbook on the processing of frozen surimi and fish jelly product in Southeast Asia.* Koon Wah Printing Pte. Ltd. Singapore. 30 pp
- Nielsen, R.G., and Pigott, G.M. 1994. Gel strength increased in low-grade heat-set surimi with blended phosphates. *J. Food Sci.* 59:246-250.



- Nilsson, K. and Ekstrand, B. 1995. Frozen storage and thawing methods affect biochemical and sensory attributes of rainbow trout. *J. Food Sci.* 60:627-630.
- Nishimoto, S., Hashimoto, A., Seki, N., Kimura, I., Toyada, K., Fujita, T. and Arai, K. 1987. Influencing factors on changes in myosin heavy chain and jelly strength of salted meat paste from alaska pollack during setting. *Nippon Suisan Gakkaishi* 53:2011-2020.
- Noguchi, S.F. 1986. Dynamic viscoelastic changes of surimi (minced meat) during thermal gelation. *Bull Japan. Soc. Fish.* 52:1261.
- Okada, M. 1985. Ingredient on gel texture. In. *Engineered Seafood Including Surimi*. R.E. Martin and R.L. Collette. (Eds) pp. 507-521. International Symposium On Engineered Seafood Including Surimi . Nov. 19-21, Sea. Ttle, Washington.
- Okada, T., Ohta, F., Inoue, N., and Akiba, M. 1985. Denaturation of carp myosin B in KCl solution during frozen storage. *Bull. Jap. Soc. Sci. Fish* 51:1887-1892.
- Park, J.W. and Lanier, T.C. 1987. Combined effects of phosphates and sugar or polyol on protein stabilization of fish myofibrills. *J. Food Sci.* 52:1509-1513.

- Park, J.W., Lanier, T.C., and Green, D.P. 1988. Cryoprotective effects of sugar polyols and/or phosphates on alaska pollack surimi. *J. Food Sci.* 53: 1-3.
- Pigott, G.M. and Tucker, B.W. 1990. *Seafood: Effect of Technology in Nutrition.* Marcel Dekker, Inc. USA. 362 pp.
- Pipatsattayanuwong, S., Park, J.W., and Marrissey, M.T. 1995. Functional properties and shelf life of fresh surimi from pacific whiting. *J. Food Sci.* 60: 1241-1244.
- Ramirez, J.A., Martin-PoLo, M.o., and Bandman, E. 2000. Fish myosin aggregation as Affected by freezing and initial physical state. *J.Food Sci.* 65:556-560.
- Rodgers, M.E., Karr, T., Biedermann, K. Ueno, H., and Harrington, W.F. 1987. Thermal stability of myosin rod from various species. *Biochemistry*, 26: 8703-8708.
- Rosinski, M.J., Barmore, C.R., Bridges, W.C., Dick, R.L., and Acton, J.C. 1989. Phosphate type and salt concentration effects on shear strength of packaging film adhesion to processed meat from a cook-in packaging system. *J. Food Sci.* 54 :1422-1425.
- Roura, S.I., Saavedra, J. P., Truco, R.e., and Crupkin, M. 1992. Conformational change in actomyosin from post-spawned hake stored on ice. *J. Food Sci.* 57:1109-1111.

- Scott, D.N., Porter, R.W., and Kourg, B. 1988. Effect of freezing and frozen storage of Alaska pollock on chemical and gel-forming properties of surimi. *J. Food Sci.* 53: 353-357.
- Shaban. O., Ochiai, Y., Watabe, S., and Hashimoto, K. 1985. Quality changes in alaska pollack meat paste (surimi) during frozen storage. *Nippon suisan Gakkaishi.* 51:1853-1858.
- Shenouda, S.Y.K. 1980. Theories of protein denaturation during frozen stored of fish flehs. *Adv. Food Res.* 26:275-280.
- Shimizu, Y., Machida R., and Takenami, S. 1981. Species variation in the gelforming characteristics of fish meat paste. *Bull Japan. Soc. Sci. Fish.*47: 95-97.
- Sikorski, Z.E. and Kotakowska, A. 1992. Changers in protein frozen stored fish. pp. 99-112
- Simpson, R., Kolbe, E., MacDonald, G. A., Lanier, T.C., and Morrissey, M.T. 1994. Surimi production from partially processed and frozen pacific whiting (*Merluccius productus*). *J. Food Sci.* 59: 272-276.
- Simpson, R., Morriseey, M.T., Kolbe, E., Lanier, T.C., and MacDonald, G.A. 1994. Effect of varying sucrose concentrations in Pacific whiting (*Merluccius productus*) *J. Aqua. Food Technol.* 3:41-52.

- Sirkorski, Z.E. 1990. Seafood : Resources, Nutritional composition and preservation CRC Press, Florida. pp 30-90.
- Sikorski, Z.E. and Kotakowoska, A.1992. Change in proteins is frozen stored fish. pp. 99-112 In.
- Smith, D.M. 1991. Factor influencing heat induce gelation of muscle proteins. In Interactions of food protein. pp. 243-256. Washington DC: American Chemical Society.
- Srinivasan, S., Xiong, Y.L., and Blanchard, S. 1997. Effects of freezing and thawing methods and storage time on thermal properties of freshwater prawns (*Macrobrachium rosenbergii*) . J. Sci Food Agric . 75: 37-44.
- Stabursvik ,E. and Martens, H. 1980. Thermal denaturation of protein in post rigor muscle tissue as studied by differential scanning calorimetry. J. Sci. Food Agric. 31:1034-1042.
- Suvanich, V., Jahncke, M.L., and Marshall, D.L. 2000. Changes in selected chemical quality characteristics of channel catfish frame mince during chill and frozen storage. J. Food Sci. 65:24-29.
- Suzuki, T. 1981. Fish and krill protein : proceesing technology. Applier science Publishers London.

- Sych, J., Lacroix, C., Adambounou, L.T., and Castaigne, F. 1990. Cryoprotective of some materials on cod-surimi proteins during frozen storage. *J. Food Sci.* 55:1222-1227.
- Wagner, J.R. and Anon, M.C. Effect of frozen storage on protein denaturation in bovine muscle. II. Influence on solubility, viscosity and electrophoretic behaviour of myofibrillar proteins. *J. Food Sci.* 21:547-558.
- Watabe, S., Ochiai, Y., and Hashimoto, K. 1982. Identification of 5,5-dithio-bis-z-nitrobenzoic acid (DTNB) and alkali light chains of piscine myosin. *Bull. Jpn. Soc. Sci. Fish.* 48:827-832.
- Watabe, S., Ushino, H., Iwamoto, M., Yamanaka, H., and Hashimoto, K. 1989. Temperature dependency of rigor-mortis of fish muscle: myofibrillar  $Mg^{2+}$ -ATPase activity and  $Ca^{2+}$  uptake by sarcoplasmic reticulum. *J. Food Sci.* 54: 1107-1115.
- Wright, D.J. and Wilding, P. 1983. Differential scanning calorimetry study of muscle and its protein : myosin and its subfragments. *J. Sci. Food Agric.* 35: 357-372.
- Wu, M.C., Lanier, T.C. and Hamann, D.D. 1985. Rigidity and viscosity changes of croaker actomyosin during thermal gelation. *J. Food Sci.* 50:14-15.
- Wu, M.C., Atallah, M.T., and Hultin, H.O. 1992. The proteins of washed minced fish muscle have significant solubility in water. *J. Food Biochem.* 15:209-218.

- Xiong, Y.L., Lou, X., Wang, C., Moody, W.G., and Harmon, R.J. 2000 Protein extraction from chicken myofibrils irrigated with various polyphosphate and NaCl solution. *J. Food Sci.* 65:96-100.
- Yean, Y.S. 1993. The quality of surimi made from Theadfin bream stored on ice for different periods. *Inter. J. Food Sci.* 28:343-346.
- Yongsawatdigul, J and Park, J.W. 1996. Linear heating rate affects gelation of Alaska pollock and Pacific whiting surimi. *J. Food Sci.* 61:149-153.
- Yongsawatdigul, J., Park, J.W., Virulhakul, P., and Viratchakul, S. 2000. Proteolytic degradation of tropical tilapia surimi *J. Food Sci.* 65:129-133.