

Appendix VI

Reprints

Index of molt staging in the black tiger shrimp (*Penaeus monodon*)

Waraporn Promwikorn¹, Pornpimol Kirirat²
and Pinij Thaweethamseewee²

Abstract

Promwikorn, W., Kirirat, P. and Thaweethamseewee, P.
Index of molt staging in the black tiger shrimp (*Penaeus monodon*)
Songklanakarin J. Sci. Technol., 2004, 26(5) : 765-772

Molting is a phenomenon of shedding the old cuticle and re-generating the new one found in crustaceans including shrimps and many other species of invertebrates. The molting cycle is a dynamic process, composed of pre-ecdysis (premolt, D stage), ecdysis (E stage), postecdysis (postmolt, A-B stages), and intermolting (C stage) stages. In healthy shrimp, molting cycles are repeated several times through shrimp life in order to increase body size and mass (growth). In this paper we gather the knowledges and important findings related to the molting cycle of crustaceans from the past until present, and highlight the physical evidence of cuticular tissue that we used for molt staging in the black tiger shrimp, *Penaeus monodon*. At the end of this paper we summarize the easily observed criteria used for molt staging in the black tiger shrimp.

Key words : molting stages, *Penaeus monodon*, setae, setal cone

¹Ph.D.(Cell and Molecular Biology), ²M.Sc.(Anatomy), Department of Anatomy, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla, 90112 Thailand.

Corresponding e-mail: waraporn.p@psu.ac.th

Received, 9 January 2004 Accepted, 8 April 2004

บทคัดย่อ

วารสารพรหมวิกร พรพิมล คีรีรัตน์ และ พินิจ ทวีธรรมเสวี
ดัชนีบ่งชี้ระยะการลอกคราบของกุ้งกุลาดำ (*Penaeus monodon*)

ว. สงขลานครินทร์ วทท. 2547 26(5) : 765-772

การลอกคราบคือกิจกรรมที่กุ้งสลัดเอาเปลือกเก่าที่ห่อหุ้มตัวออกและสร้างเปลือกใหม่ขึ้นมาแทนที่ การลอกคราบเป็นกระบวนการที่มีการเปลี่ยนแปลงของร่างกายเป็นวงจรอย่างต่อเนื่อง ซึ่งสามารถแบ่งย่อยเป็นระยะต่าง ๆ 4 ระยะ ได้แก่ ระยะก่อนการลอกคราบ ระยะลอกคราบ ระยะหลังการลอกคราบ และระยะระหว่างวงจรการลอกคราบ ทุก ๆ วงจรการลอกคราบของกุ้งที่อยู่ในสภาวะปกติ กุ้งจะมีขนาดเพิ่มขึ้นทุกครั้ง ดังนั้นจึงกล่าวได้ว่าการลอกคราบมีผลโดยตรงต่อการเจริญเติบโตของกุ้งกุลาดำ การศึกษาครั้งนี้ได้ประมวลความรู้และงานวิจัยที่สำคัญที่เกี่ยวข้องกับการลอกคราบของสัตว์ในตระกูลปูและกุ้งตั้งแต่อดีตจนถึงปัจจุบัน โดยเน้นที่การอธิบายลักษณะทางกายภาพของเปลือกและเนื้อเยื่อผิวหนังที่เปลี่ยนแปลงตลอดวงจรการลอกคราบของกุ้งกุลาดำ และในตอนท้ายได้สรุปเกณฑ์สำคัญที่ใช้เป็นดัชนีบ่งชี้ระยะการลอกคราบของกุ้งกุลาดำ

ภาควิชาวิทยาศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยสงขลานครินทร์ อำเภอหาดใหญ่ จังหวัดสงขลา 90112

Molting, a phenomenon of shedding the old cuticle and re-generating the new one, occurs in many species of invertebrates, including crustaceans. It is an essential process for post-larval growth of crustaceans. In healthy animals, molting cycles are repeated several times in order to allow growth throughout crustacean life. Through the molting cycle, the animals show dramatic changes in many aspects, including body structure, biology and behavior. The regulatory mechanism of the molting process has long been studied in various species of crustaceans, mainly in crabs. The studies related to the molting cycle have so far paid attention to molt-inhibiting hormone from the X-organs located at the eyestalks (Hubschman and Armstrong, 1972; Rao, 1973; Freeman and Costlow, 1979; Keller and Schmid, 1979; Bellon-Humbert and Van Herp, 1988; Sithigorngul *et al.*, 1999; Watson, 2001), and molt-stimulating ecdysteroid hormones secreted by the Y-organs (Carlisle, 1957; Hubschman and Armstrong, 1972; Charmantier and Trilles, 1973; 1979; Keller and Willig, 1976; Chaix *et al.*, 1976; Spaziani, 1999; Chang, 2001), although changes in the structures and biochemistry of hepatopancreas, haemolymph and integument related to molting cycle have also been reported in many

species (for example see Watanabe and Kono, 1997; Watanabe *et al.*, 2000; Sousa *et al.*, 2001; Fernandez *et al.*, 2001; Pratoomchart *et al.*, 2002a; b). These findings accumulate more knowledge about the regulatory mechanism of the molting. However, the direct evidence in the black tiger shrimps (*Penaeus monodon*), which is well known as an important agricultural export product of Thailand, has not been yet elucidated. The objective of this study is to gather background knowledge about molting stages and criteria used to determine molting stages in crustaceans, focusing especially on physical evidence used to determine molting stages in the black tiger shrimp. We hope that this paper will be useful for further researches which aim to promote shrimp growth, which is at the heart of shrimp farming.

Molting stages and criteria used to determine molting stages in crustaceans.


Each molting cycle of crustaceans is composed of different stages including preecdysis (pre-molt, D stage), ecdysis (E stage), postecdysis (postmolt, A-B stages), and intermolting (C stage) stages (Drach, 1939; Skinner, 1962). Various criteria are in use for staging the crustacean molt cycle. These include the degree of hardness of

the exoskeleton, changes in matrices of the setae (Drach, 1939), the growth of re-generating limb buds (Skinner, 1962; Stevenson *et al.*, 1968; Stevenson and Henry, 1971; Hopskin, 1982), the progressive development of gastroliths in the digestive organs (McWhinnie, 1962; Hopkins, 1977; Rao *et al.*, 1977), histological changes of organs and tissues (Travis, 1957; Skinner 1962; Stevenson, 1968; Stevenson *et al.*, 1968; Benhalima *et al.*, 1998), development of setae (Lyle and MacDonald, 1983; Moriyasu and Mallet, 1986), and a combination of the above methods (O' Halloran and O'Dor, 1988; Musgrove, 2000). The premolt stage is a period of biological preparation of organs and tissues all over the animal body for shedding its cuticle (exoskeleton). It is divided into four sub-stages; D1, D2, D3 and D4 depending on the degree of retraction of the epidermal tissue from the cuticle (Drach, 1939). After the D4 stage, the animals will spend a few seconds for shedding their old cuticle during ecdysis (E stage). The animals will subsequently enter postmolt stage where they will spend some time for building up and hardening the new cuticle. Postmolt is divided into an A and a B stage. The A stage follows immediately after the ecdysis. The main task of the A stage is the retraction of the contents of the new setae. The A stage is divided into A1 and A2 sub-stages. The B stage is the period for mineralisation of the cuticle, and is divided into the B1 and B2 sub-stages, the first is defined by the retraction of the contents of the setae to the region where cones will form inside, the latter is defined by the formation of setal cones. At the end of the B2 stage the formation of the setal cones and chemical change in the preecdysial layer of the newly-synthesized cuticle are completed (Drach, 1939). The animals subsequently enter the C stage, in which four sub-stages are distinguished; C1, C2, C3 and C4. Stages C1 and C2 are defined by an increase of integument rigidity, but there are no marked criteria to divide these. At stage C3 the integument reaches its final stage of rigidity, after which in stage C4 the synthesis of the membranous layer of the procuticle is completed. Thus C3-C4 stages are so-called intermolting stages, as the

formation of the new cuticle is finished, and the animals await the signals from ecdysteroid hormones from the Y-organs located at the mouth region to trigger re-entering the premolt of the next cycle (Carlisle, 1957; Hubschman and Armstrong, 1972; Charmantier and Trilles, 1973; 1979; Chaix *et al.*, 1976; Keller and Willig, 1976). While the animal body is not under limitation of the cuticle during the period from late premolt to postmolt, they are increasing up body size and mass, hence growing.

Molt staging in the black tiger shrimp.

Molt staging in penaeid shrimps was previously described by numbers of researchers based on various criteria (Schafer 1968; Longmuir 1983; Smith and Dall 1985; Yashiro, 1991). In this paper we reveal physical characteristics that we used to determine molting stages in the black tiger shrimp, *Penaeus monodon*, based on the criteria described by Drach, 1963 and a number of other researchers (for review see Skinner, 1985; Stevenson, 1985). From our direct observations, we found that in the intermolt stage (C stage, Figure a), setal cones (SC) were fully developed, and each was arranged in a neat line at the base of the setae. These two structures form a mature cuticle. The epidermal tissue (E), which is located under (interior to) the cuticle, was spreading throughout the area under the cuticle and between pairs of setal cones. When the premolt stage began, the process of continuous retraction of the epidermal tissue away from the cuticle started. The degree of retraction of the epidermal tissue determined the sub-stages of the premolt. Once the retraction begins, the shrimp is designated as stage D0. At the end of D0 stage, a clear straight margin of the epidermal tissue was observed at the base of the setal cones (Figure b). The D1 stage (Figure c) was defined by the presence of a narrow clear zone between the setal cones and the epidermal tissue. The clear zone implies the formation of a new cuticle. In the D2 stage (Figure d), the width of the clear zone between the setal cone and the epidermal tissue increased, and the edge of the epidermal tissue now had a wave-like pattern. This implies the continuous



Determination of molting stage. Uropods of the black tiger shrimps were examined and photographed under a light microscope (Olympus BX51) connected to a digital camera (Olympus DP11). The criteria used for molt staging followed Drach's staging. The images of physical characteristics indicate intermolt stage (a), premolt stage (b-f) and postmolt stage (g,h). Bar = 50 μ m. EE = epidermal edge, I = indent pattern of the epidermis, L = white layer at the edge of the epidermis, S = setae, SC = seta cone, SCn = newly-formed setal cones, Sn = newly-formed seta, \curvearrowright = wavy edge of epidermis, * = clear zone between cuticle and epidermis.

synthesis of new cuticle. Small and quite thin fiber-like projections between each setal cone and epidermal tissue were observed. These projections would later become new setae. In the next premolt stage, the D3 stage (Figure e), the clear zone between the setal cones and the epidermal tissue was clearly widened. A thin white layer at the edge of the epidermal tissue together with a highly wavy edge of the epidermal tissue could then be observed. The fiber-like projections connecting each setal cone with the epidermal tissue were still seen. At the final, the D4 stage of premolt (Figure f), the clear zone between the setal cones and the edge of the epidermal tissue was extremely clear and dominant. The typical characteristics of the epidermal tissue were clearly marked. As the white layer at its edge was reflecting the light, and formed sharp and serrate notches, young setae were clearly seen projecting from the impressions between the sharp notches. The pigments in the epidermal tissue were obviously indent and arranged in a paralleled-band fashion. The setal cones then started to deform. After ecdysis the shrimps immediately entered the post molt A stage (Figure g), in which the new cuticle and the setae were very soft and delicate. The setal cone could then not yet be seen. Only a few hours after ecdysis, the shrimps entered post molt B stage (Figure h) where the cuticle was hardening by mineralisation of inorganic compounds such as calcium and phosphorus (Roer and Dillaman, 1984; Cameron, 1985; Machado *et al.*, 1990; Compere *et al.*, 1993; Ziegler, 1997; Pratoomchart *et al.*, 2002a). In the B stage, young setal cones were developing. Different sizes of setal cones were thus observed at the base of the setae. When the development of the setal cones was completed, the shrimps entered C stage (Figure a). In late C stage (C 3-4) when the synthesis of all layers of new cuticle was complete, the shrimps were designated as in the intermolt stage.

Concluding Remarks

The criteria used for molt staging in the black tiger shrimp included:

- 1) The degree of retraction of the epidermal tissue from the cuticle.
- 2) The width of the clear zone between the setal cones and the epidermis.
- 3) The characteristics of the epidermis.
- 4) The presence of newly-formed setae.
- 5) The formation of setal cones

All of above criteria make up easily observed physical characteristics of each stage as follows:

- 1) Intermolt stage (C stage): mature setal cones and full-spread epidermis
- 2) Premolt stage:
 - D0- a clear margin of the epidermal tissue at the base of setal cones
 - D1- a clear narrow zone between the setal cones and the epidermis
 - D2- a wider clear zone and a wavy edge of the epidermis
 - D3- a wider clear zone, highly wavy edge of the epidermis and a white thin layer at the edge of the epidermis
 - D4- an obvious wider clear zone, serrated edge of the epidermis, light-reflecting white layer at the edge of the epidermis, and paralleled-band fashion of the epidermis.
- 3) Postmolt stage:
 - A- soft and delicate setae, absence of setal cones
 - B- presence of the young setal cones

Further application.

The above conclusions can serve as standard criteria for upcoming experiments. For example, examination of physical and biological changes of organs and tissues through the molting cycle, effect of diseases on molting cycle progression, role of environment on molting cycle, etc. The forthcoming results will lead to a clearer understanding of the regulatory mechanism of the molting cycle, specially of *Penaeus monodon*.

Acknowledgements

We specially thank Professor Boonsirm Withyachumnarnkul (Dept. of Anatomy and Centex

shrimp, Mahidol University) and Dr. AUSA Chandumpai (Dept. of Biochemistry, Faculty of Science, PSU) for critical suggestions. This work was funded by the Ministry of University Affairs, Thailand Research Fund (TRF), and the Dept. of Anatomy, Faculty of Science, PSU.

References

- Bellon-Humbert, C. and Van Herp, F. (1988). Localisation of serotonin-like immunoreactivity in the eyestalk of the prawn *Palaemon serratus* (Crustacea, Decapoda, Natantia). *J. Morphol.* 196: 307-320.
- Benhalima, K., Moriyasu, M. and Hebert, M. (1998). A technique for identifying the early-premolt stage in the male snow crab *Chionoecetes opilio* (brachyura: Majidae) in Baie des Chaleurs, southern gulf of St. Lawrence. *Can. J. Zool.* 76: 609-617.
- Cameron, J.N. (1985). Post-moult calcification in the blue crab (*Callinectes sapidus*): Relationships between apparent net H⁺ excretion, calcium and bicarbonate. *J. Exp. Biol.* 119: 275-285.
- Carlisle, D.B. (1957) On the hormonal inhibition of moulting in Decapod Crustacea II. The terminal anecdyosis in crabs. *J. Mar. Biol. Assoc. U.K.* 36: 291-307.
- Chaix, J.-C., Trilles, J.-P. and Vernet, G. (1976). Degenerescence de l'organe Y chez les males puberes d' *Acanthonyx lunulatus* (Risso) (Crustacea, Decapoda, Oxyrhyncha). *C.R. Acad. Sc. Paris* 283: 523-525.
- Charmantier, G. and Trilles, J.-P. (1973). Degenerescence de la glande de mue chez les males puberes de *Sphaeroma serratum* (Crustace, Isopode). *C.R. Acad. Sc. Paris* 276: 581-583.
- Charmantier, G. and Trilles, J.-P. (1979). La degenerescence de l'organe y chez *Sphaeroma serratum* (Fabricius, 1787) (Isopoda, Flabllifera): Etude ultrastructurale. *Crustaceana* 36: 29-38.
- Chang, E.S., Chang, S.A., Mulder, E.P. (2001). Hormones in the lives of crustaceans: An overview. *Amer. Zool.* 41: 1090-1097.
- Compere, P., Morgan, J.A. and Goffinet, G. (1993). Ultrastructure location of calcium and magnesium during mineralization of the cuticle of the shore crab, as determined by the K-pyromantimonate method and X-ray microanalysis. *Cell Tissue Res.* 274: 567-577.
- Drach, P. (1939). Mue et cycle d'intermue chez les crustaces Decapodes. *Annls. Inst. Oceanogr.* 19: 103-391.
- Fernandez-Gimenez, A.V., Garcia-Carren, F.L., Navarrete del Toro, M.A. and Fenucci, J.L. (2001). Digestive proteinases of *Artemesia longinans* (Decapoda, Penaeidae) and relationship with molting. *Comp. Biochem. Physiol.* 130: 331-338.
- Freeman, J.A. and Costlow, J.D. (1979). Hormonal control of apolysis in barnacle tissue epidermis *in vitro*. *J. Exp. Zool.* 210: 333-346.
- Hopkins, P.M. (1977). Control of gastrolith deposition in crayfish. *Amer. Zool.* 17: 900.
- Hopskin, P.M. (1982). Growth and regeneration patterns in the fiddler crab, *Uca pugilator*. *Biol. Bull.* 163: 301-319.
- Hubschman, J.H. and Armstrong, P.W. (1972). Influence of ecdysterone on molting in *Palaemonetes*. *Gen. Comp. Endocrinol.* 18: 345-438.
- Keller, R. and Schmid, E. (1979). *In vitro* secretion of ecdysteroids by Y-organs and lack of secretion by mandibular organs of the crayfish following molt induction. *J. Comp. Physiol.* 130: 347-353.
- Keller, R. and Willig, A. (1976). Experimental evidence of the molt controlling function of the Y-organ of a macruran decapod, *Orconectes limosus*. *J. Comp. Physiol.* 108: 271-278.
- Longmuir, E. (1983). Setal development, moult-staging and ecdysis in the banana prawn *Penaeus merguensis*. *Mar. Biol.* 77: 183-190.
- Lyle, W.G. and MacDonald, C.D. (1983). Molt stage determination in the Hawaiian spiny lobster *Panulirus marginatus*. *J. Crust. Biol.* 3: 208-216.
- Machado, J. Ferreira, K.G., Ferreira, H.G. and Fernandes, P. L. (1990). The acid-base balance mantle epithelium *Anodonta cygnea*. *J. Exp. Zool.* 150: 159-169.

- McWhinnie, M.A. (1962). Gastrolith growth and calcium shifts in the freshwater crayfish, *Orconectes virilis*. *Comp. Biochem. Physiol.* 7: 1-4.
- Moriyasu, M. and Mallet, P. (1986). Molt stages of the snow crab *Chionoecetes opilio* by observation of morphogenesis of setae on the maxilla. *J. Crustacean Biol.* 6: 709-718.
- Musgrove, R.J. (2000). Molt staging in the southern rock lobster *Jasus edwardsii*. *J. Crustacean Biol.* 20: 44-53.
- O'Halloran, M.J. and O'Dor, R.K. (1988). Molt cycle of male snow crabs, *Chionoecetes opilio*, from observations of external features, setal changes, and feeding behavior. *J. Crustacean Biol.* 8: 164-176.
- Pratoomchart, B., Sawangwong, P., Guedes, R., De lurdres Reis, M. and Machado, J. (2002b). Cuticle ultrastructures changes in the crab *Scylla serrata* over the molt cycle. *J. Exp. Zool.* 293: 414-426.
- Pratoomchart, B., Sawangwong, P., Pakkong, P. and Machado, J. (2002a). Organic and inorganic compound variations in haemolymph, epidermal tissue and cuticle over the molt cycle in *Scylla serrata* (Decapoda). *Comp. Biochem. Physiol.* 131A: 243-255.
- Rao, K.R., Fingerman, S.W. and Fingerman, M. (1973). Effects of exogenous ecdysones on the molt cycles of fourth and fifth stage American lobsters, *Homarus americanus*. *Comp. Biochem. Physiol.* 44A: 1105-1120.
- Rao, K.R., Mohrherr, C.J., Reinschmidt, D., and Fingerman, M. (1977). Gastrolith growth during proecdysis in the crayfish *Faxonella clypeata* (hay, 1899) (Decapoda astacoida). *Crustaceana.* 32: 256-264.
- Roer, R. and Dillaman, R. (1984). The structure and calcification of the crustacean cuticle. *Amer. Zool.* 24: 893-909.
- Schafer, H.J. (1968). The determination of some stages of the molting cycle of *Penaeus duorarum*, by microscopic examination of the setae of the endopodites of pleopods. *FAO Fish. Rep.* 57: 381-391.
- Skinner, D.M. (1962). The structure and metabolism of a crustacean integumentary tissue during a molt cycle. *Biol. Bull.* 123: 635-647.
- Skinner, D.M. (1985). Molting and regeneration. In: Bliss, D.E. and Mantel, L.H. *The biology of crustacea*, Academic Press, New York, Vol. 9: 43-128.
- Sithigorngul, W., Jaideechoey, S., Saraithongkum, W., Longyant, S. and Sithigorngul, P. (1999). Purification and characterization of an isoform of crustacean hyperglycemic hormone from the eyestalk of *Macrobrachium rosenbergii*. *J. Exp. Zool.* 284: 217-224.
- Smith, D.M. and Dall, W. (1985). Moulting staging the tiger prawn *Penaeus esculentus* Haswell. In: Rothlisberg P.C., Hill, B.J. and Staples, D.J. eds., *Second Australian national Prawn Seminar*, Cleveland, Queensland, Australia. pp 85-93.
- Sousa, L.G. and Petriella, A.M. (2001). Changes in the hepatopancreas histology of *Palaemonetes argentimus* (Crustacea, Caridea) during moult. *Biocell.* 25: 275-281.
- Spaziani, E., Mattson, M. and Wang, W.L. (1999). Signaling pathways for ecdysteroid hormone synthesis in crustacean Y-organs. *Amer. Zool.* 39: 496-512.
- Stevenson, J.R. (1968). Metecdysial molt staging and changes in the cuticle in the crayfish *Orconectes sariborni* (Faxon). *Crustaceana* 14: 169-177.
- Stevenson, J.R. (1985). Dynamics of the integument. In: Bliss, D.E. and Mantel, L.H. *The biology of crustacea*. Academic Press, New York, Vol. 9: 1-41.
- Stevenson, J.R. and Henry, B.A. (1971). Correlation between the molt cycle and limb regeneration in the crayfish *Orconectes sariborni* (Hagen) (decapoda, Astacidea). *Crustaceana.* 20: 301-307.
- Stevenson, J.R., Guckert, R.H. and Cohen, J.D. (1968). Lack of correlation of some proecdysial growth and developmental processes in the crayfish. *Biol. Bull. (Woods Hole, Mass.)* 134: 160-175.

- Travis, D.F. (1957). The molting cycle of the spiny lobster, *Panulirus argus* Latreille. II. Preecdysial histological and histochemical changes in the hepatopancreas and integumental tissue. Biol. Bull. (Woods Hole, Mass). 108: 88-112.
- Watanabe, T. and Kono, M. (1997). Isolation of a DNA encoding a chitinase family protein from cuticular tissues of the Kuruma prawn *Penaeus japonicus*. Zoolog. Sci. 14: 65-68.
- Watanabe, T., Persson, P. End, H. and Kono, M. (2000). Molecular analysis of two genes, DD9A and B, which are expressed during the postmolt stage in the decapod crustacean *Penaeus japonicus*. Comp. Biochem. Physiol. 125: 127-136.
- Watson, R.D., Lee, K.J., Qiu, S., Luo, M., Umphrey, H.R., Roer, R.D. and Spaziani, E. (2001). Molecular cloning, expression, and tissue distribution of Crustacean molt-inhibiting hormone. Amer. Zool. 41: 407-417.
- Yasiro, R. (1991). Methods for molt and ovary staging in marine shrimp. Thai Fisheries Gazette. 43(6): 451-454. (in Thai).
- Ziegler, A. (1997). Ultrastructural changes of the anterior and posterior sternal integument of the terrestrial isopod *Porcellio scaber* Latr. (Crustacea) during the moult cycle. Tiss. Cell. 29: 63-76.

Histological characterization of cuticular depositions throughout the molting cycle of the black tiger shrimp (*Penaeus monodon*)

Waraporn Promwikorn¹, Piyakorn Boonyoung², and Pornpimol Kirirat³

Abstract

Promwikorn, W., Boonyoung, P., and Kirirat, P.

Histological characterization of cuticular depositions throughout the molting cycle of the black tiger shrimp (*Penaeus monodon*)

Songklanakarin J. Sci. Technol., 2005, 27(3) : 499-509

The aim of this study was to investigate changes of cuticular deposition throughout the molting cycle of the black tiger shrimp (*Penaeus monodon*). The molting stages were first determined using physical criteria. Then the deposition of collagen fibers, carbohydrate, lipid and calcium salt in the cuticle were examined histologically throughout the molting cycle. The results show that the mature cuticle of the intermolt stage-shrimp is composed of four sub-layers from exterior to interior including an epicuticle, an exocuticle, an endocuticle, and a membranous layer. The differences of cuticular depositions reflect the different functions of each cuticular sub-layer. We also investigated the changes of these depositions that were related to the molting cycle. We found that the dynamic process of cuticular degradation and regeneration occurred constantly throughout the molting cycle. During the premolt period, the new epicuticle and exocuticle (pre-ecdysial cuticle) were synthesized on the inside of the old ones. After exuviations (post-ecdysis), the syntheses of the endocuticle and membranous layer of the new cuticle progressed chronologically. When the synthesis of the membranous layer is complete, the shrimps then enter the intermolt stage. At the end of this paper, we also summarize the histological criteria to use for determining the stage of the molt in the black tiger shrimp.

Key words : cuticular sub-layers, molting stages, *Penaeus monodon*

¹Ph.D. (Cell and Molecular Biology), ²M.Sc. (Zoology), ³M.Sc. (Anatomy), Department of Anatomy, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand.

Corresponding e-mail: waraporn.p@psu.ac.th

Received, 13 September 2004

Accepted, 26 November 2004

บทคัดย่อ

วารกรณ์ พรหมวิกิร ปิยากร บุญยัง และ พรพิมล คีรีรัตน์
การเปลี่ยนแปลงสัณฐานทางเนื้อเยื่อวิทยาของเปลือกกุ้งกุลาค่าตลอดวงจรการลอกคราบ
ว.สงขลานครินทร์ วทท. 2548 27(3) : 499-509

ระยะของการลอกคราบในกุ้งกุลาค่าถูกตรวจสอบโดยใช้เกณฑ์ทางกายภาพก่อนที่เส้นใยคอลลาเจนคาร์โบไฮเดรตไขมัน และ เกลือแคลเซียม ในเปลือกจะถูกศึกษาด้วยวิธีการทางเนื้อเยื่อวิทยา ผลการศึกษาพบว่า กุ้งที่อยู่ในระยะระหว่างวงจรการลอกคราบจะมีเปลือกที่สมบูรณ์เต็มที่ซึ่งประกอบด้วย ชั้นย่อย ๆ 4 ชั้นเรียงจากด้านนอกเข้าด้านใน ได้แก่ epicuticle exocuticle endocuticle และ membranous layer โดยที่ในแต่ละชั้นมีการสะสมของเส้นใยคอลลาเจน คาร์โบไฮเดรต ไขมัน และ เกลือแคลเซียมที่แตกต่างกัน ซึ่งความแตกต่างนี้สะท้อนให้เห็นถึงหน้าที่ที่แตกต่างกันของแต่ละชั้นย่อย นอกจากนี้ยังพบว่าลำดับการย่อยสลายและเกิดใหม่ของเปลือกกุ้งกุลาค่าเกิดขึ้นอย่างเป็นแบบแผนที่แน่นอนตลอดวงจรการลอกคราบ และในคอนท้ายได้สรุปลักษณะทางเนื้อเยื่อวิทยาของเปลือกที่สามารถใช้เป็นเกณฑ์บ่งชี้ระยะของการลอกคราบของกุ้งกุลาค่าเพิ่มเติมจากลักษณะทางกายภาพที่เราเคยรายงานไว้ก่อนหน้านี้

ภาควิชากายวิภาคศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยสงขลานครินทร์ อำเภอหาดใหญ่ จังหวัดสงขลา 90112

Crustacea molting cycles are controlled by neurosecretory hormones from the X- and Y-organs located in the eye stalks and near mouth region, respectively (Carlisle, 1957; Keller and Schmid, 1979; Soumoff and O'Connor, 1982; Skinner, 1985a&b; Lee et al., 1998; Spaziani et al., 1999; Watson et al., 2001; Gilbert et al., 2002; Nakatsuji and Sonobe, 2004). Ecdysteroid hormones are secreted from the Y-organ to stimulate "apolysis" and "ecdysis". Apolysis, corresponding to the D0 stage, is a process of separation of the old cuticle from the epidermis thus producing a space between the old cuticle and the epidermis underneath. This space is later replaced with newly synthesized cuticle to replace the old one. Ecdysis is the process at the brief moment of shedding the cuticle from the animal's body. Apolysis and ecdysis are followed by the process of cuticular re-building and increasing body size. These biological events are in a continuous cycle throughout the animal's life. Studies about the molting cycle have been investigated for seventy years (for example see Adams et al., 1982; Skinner, 1985a&b; Lee et al., 1998; Spaziani et al., 1999; Musgrove, 2000; Watson

et al., 2001; Gilbert et al., 2002; Gorokhova, 2002; Pratoomchat et al., 2002a; b; Nakatsuji and Sonobe, 2004), however, the mechanisms that regulate the molting cycle are not yet fully understood, especially that of the molting cycle of the black tiger shrimp (*Penaeus monodon*), which is an important agricultural export product of Thailand. We are therefore attempting to understand the regulatory mechanism of ecdysis and the molting cycle in the black tiger shrimp. We hope that the results will be beneficial for promoting the growth the black tiger shrimp and other species of crustaceans. We have recently reported on the physical characteristics of the cuticle that can be used to determine the stage of the molt in the black tiger shrimp (Promwikorn et al., 2004). In this paper we report further on the cuticular changes throughout the molting cycle as revealed by histological characterization of the collagen fibers, carbohydrate, lipid and calcium salt. We also summarize the histological criteria used as an alternative method for determining the stage of the molt in the black tiger shrimp.

Materials and Methods

Animals

Healthy black tiger shrimps (*Penaeus monodon*) were brought from commercial farms. During transportation to our laboratory in PSU, the shrimps were continuously oxygenated. The age of the shrimps was estimated at 90 days, and their body weight was between 10-20 g.

Shrimp culture

Natural sea water used in all experiments was stored in a tank for at least 2 weeks before use. Some days before the experiments, the sea water in the aquarium was continuously aerated. The salinity was adjusted to be similar to the sea water used in the farms (10-20 ppt. depending on the salinity of each farm). Once the shrimps had arrived, the molting stage of each shrimp was determined prior to its cultivation with continuous aeration. The shrimps were fed three times a day with commercial pellet. Natural day-light and atmospheric temperatures were used throughout the experiment.

Determination of molting stages

The molting stages were examined daily. Briefly, each shrimp was gently picked up by hand. They were then quickly examined with a light microscope. The physical criteria used for determining the stage of the molt were those used in our previous report (Promwikorn *et al.*, 2004). After 1-2 min examination, the shrimps were either placed back into the aquarium, or executed.

Histological study

The cuticular tissues at the carapace and trunk (first abdominal somite) of at least 10 shrimps at each molting stage were dissected and immediately fixed in Davidson's fixative to eliminate any calcium salt in the cuticle. The fixation stage was continued for 72 h at room temperature with a daily change of fresh fixative. The cuticular tissues were subsequently dehydrated in increasing concentrations of ethanol that ranged from 50 to 100%,

and prepared for routine histological embedding in paraffin blocks. Paraffin sections of 0.5 μ m thickness were routinely de-paraffinized and stained for collagen fibers with Masson's trichrome (Bancroft and Gamble modification, 2002a), and carbohydrate with periodic acid Schiff's reagent, PAS (Bancroft and Gamble, 2002b). The stained tissue sections were examined with a light microscope (Olympus BX 51) and photographed with a digital camera (Olympus DP11) connected to the microscope. For calcium salt staining, the cuticular tissues were fixed with 10% formalin instead of Davidson's fixative to preserve calcium salt in the tissue (Bancroft and Gamble, 2002c). For lipid staining, fresh cuticular tissues were sectioned with a cryostat (Leica CM1850) at -20 °C, and stained with Oil Red O, before mounting in glycerine jelly (Sheehan and Hrapchak, 1980).

Results and Discussion

The crustacean cuticle is composed of 2 main layers from exterior to interior which are the epicuticle and procuticle, respectively. The procuticle is composed of three sub-layers including an exocuticle, an endocuticle and a membranous layer. The epicuticle and exocuticle are synthesized before ecdysis and form the so-called pre-ecdysial layer. The endocuticle and the membranous layer are synthesized after ecdysis, and form the so-called post-ecdysial layer (Skinner, 1985b). The main components of the cuticle include both inorganic and organic materials in a 60:40 ratio. The inorganic materials found in the cuticle include calcium, chloride, copper, magnesium, manganese, phosphorus, potassium, and sulfur (Roer and Dillaman, 1984; Mangum, 1992; Compere *et al.*, 1993; Pierce, *et al.*, 2001; Pratoomchat *et al.*, 2002a; b; Wang *et al.*, 2003). The organic materials include lipid, glycoprotein, protein, glycosaminoglycans, mucopolysaccharides, carbohydrate, and chitin (Glynn, 1968; Vigh and Dendinger, 1982; Roer and Dillaman, 1984; Wheeler and Sikes, 1984; Marlowe *et al.*, 1994; Andersen, 1999; Roer *et al.*, 2001; Pratoomchat

et al., 2002b). Although these components are found in many species of crustaceans, mainly in crabs, the evidence of their location in the cuticle is limited. Using histochemical methods we have localised both organic and inorganic components including collagen fibers, carbohydrate, lipid and calcium salt in the mature cuticle of the black tiger shrimp. We have also investigated how these components are deposited in the cuticle with respect to the molting cycle.

1. Organic and inorganic contents are deposited differently in cuticular sub-layers.

Histochemical methods were used to locate the collagen fibers, carbohydrate, lipid and calcium salt in the cuticle of the black tiger shrimp. The collagen fibers, that give strength to the cuticle, were stained blue with Masson's trichrome (aniline blue, Ponceau S, and hematoxylin), carbohydrate was stained pink with periodic acid Schiff's (PAS) reagent, lipid was stained red-pink with oil red O (ORO), and calcium salt was stained orange-red with alizarin red S (ARS). It was shown that the mature cuticle of the intermolt shrimp consisted of four layers each with a distinctive texture and characteristic (Figure 1).

The outermost layer, an epicuticle, was a thin layer, which stained with PAS (deep magenta), trichrome (red), ORO (red-pink), and ARS (orange-red) methods. This indicates that the epicuticle is rich in carbohydrate, lipid, calcium salt and protein, but lacks collagen fibers. The red colour obtained after staining with Ponceau S in trichrome indicated that the epicuticle is rich in protein. Also the deep magenta colour obtained with PAS staining, gave a strong indication that glycoprotein is abundant in the epicuticle.

The procuticle could be divided into three distinctive layers from the exterior to the interior; an exocuticle, an endocuticle and a membranous layer. The exocuticle close to the epicuticle has typical alternating light and dark lamellae. It stained blue with trichrome, dark blue with PAS (counterstained with hematoxylin), and orange-red with ARS, but was not stained with ORO. This

indicates that the exocuticle consists of bundles of collagen fibers arranged in a lamella-fashion with an intervening interlamella-matrix composed of carbohydrate and protein (the latter may be a precursor material for collagen fibers). Calcium salt is deposited throughout, but lipid is not detected in this layer.

The endocuticle lies inside the exocuticle. This layer stained light blue with trichrome, light pink with PAS and deep orange-red with ARS. Again the endocuticle did not stain with ORO. The characteristic of the outer part of the endocuticle is similar to that of the exocuticle, but has less content of collagen fiber, and carbohydrate. The inner part of the endocuticle has more delicate and compact lamellae of collagen fibers, carbohydrate and calcium salt, and closely resembles the innermost membranous layer. The membranous layer, found between the endocuticle and epidermis (sometimes referred as hypodermis, or epithelium), was clearly distinguishable from the endocuticle after staining by PAS. The membranous layer stained a darker blue with PAS, a stronger blue with trichrome, and orange with ARS when compared with the endocuticle, but did not stain with ORO. This indicates that the endocuticle and membranous layer are also rich in collagen fibers, protein, carbohydrate, and calcium salt, but lack lipid.

Our results indicate that collagen fibers, carbohydrate, calcium salt and lipid are the main components in the cuticle of the black tiger shrimp. However, they are deposited differently in each cuticular sub-layer. This presumably is a reflection of their functions. The lipid is detected only in the epicuticle where it could act as a water-shield or in body-temperature regulation. The collagen fibers, calcium salt, and carbohydrate are abundant in all sub-layers of the procuticle. The collagen fibers play a major role in cuticular architecture, and give strength to the cuticle. Calcium salt increases the hardness of the cuticle, possibly to protect the shrimp from external mechanical forces. In the epicuticle, the carbohydrate is in the form of a glycoprotein. The carbohydrate may be a




Figure 1. Localisation of carbohydrate (stained magenta, a), collagen fibers (stained blue, b), calcium salt (stained orange-red, c), and lipid (stained pink, d) in the mature cuticle of intermolt shrimp. Four cuticular sub-layers are shown; epicuticle (Ep), exocuticle (Ex), endocuticle (Ed), and membranous layer (Ml). Bar = 10 µm.

precursor for chitin, a polymer consisting of 80-90% N-acetylglucosamine and 10-20% glucosamine. The presence of protein and carbohydrate in the exocuticle and membranous layer may indicate the sites for chitin deposition. Our study provides additional information to previous studies on the location of collagen fibers, carbohydrate, calcium salt and lipid in the cuticular sub-layers of the black tiger shrimp.

2. The epicuticle and exocuticle are synthesized before ecdysis.

It would be of interest to learn how the degradation and regeneration of the cuticular sub-layers are related to the molting cycle. During the molting cycle, the cuticle changes dynamically in two aspects; i) the thickness of the cuticle, this depends on the number of cuticular sub-layers

(Mykes, 1980; Stevenson, 1985), and ii) the deposition of the organic and inorganic components in the cuticle. We have shown that collagen fibers, carbohydrate, calcium salt and lipid are deposited in different cuticular sub-layers. We therefore examined in more detail of changes in the depositions of these components during the molting cycle. We found that during the early premolt period (D0-2), Figure 2, the membranous layer stained light blue with trichrome, light pink with PAS, and light orange-red with ARS, or often could not be clearly identified. These staining reactions were decreased if compared to the staining reactions found at the intermolt stage. This indicates that the membranous layer is being degraded during the premolt stages. This may allow the protein, carbohydrate, and calcium salt present in these two layers to be re-absorbed by




Figure 2. Localisation of carbohydrate (a,b), collagen fibers (c,d), calcium salt (e,f), and lipid (g,h) in the cuticle during the early premolt period (left column) and late premolt period (right column). New epicuticle (nEp) and exocuticle (nEx) is synthesized on the inside of the old exocuticle (oEx), and thickening in a later stage. Bar = 10 µm.




Figure 3. The cuticle of the black tiger shrimp immediately after ecdysis. Only the new epicuticle (nEp) and exocuticle (nEx) are present. The cuticle was stained with PAS for carbohydrate (a), and alizarin red S for the calcium salt (b). Bar = 10 μ m.

the underlying epidermis. In the late premolt period (D3-4 stage), newly secreted epicuticle (nEp) and exocuticle (nEx) were sequentially produced inside the old cuticle. These data indicate that during the premolt period the endocuticle and membranous layer of the old cuticle are re-absorbed, prior to the formation of the new epicuticle and exocuticle and the shedding of the exuviae.

3. The endocuticle and membranous layers are synthesised after ecdysis.

Immediately after ecdysis, during A1 - 2 stages, only an epicuticle and an exocuticle were observed in the intact cuticle (Figure 3). Ten hours after ecdysis, a small layer of endocuticle was seen at the inner border of the exocuticle. This layer stained blue with trichrome, pink with PAS, and light orange-red with ARS, but did not stain with ORO. Two days after ecdysis, in the B2 stage, the thickness of the newly synthesized endocuticle increased. The stain for the calcium salt was now clearly visible (Figure 4f). Three to four days after ecdysis, all layers of the cuticle including the membranous layer were visible (see Figure 1). These findings indicate that the syntheses of the new epicuticle and exocuticle have finished before ecdysis. However, another two layers are synthesised after ecdysis. The synthesis

of the endocuticle starts immediately after ecdysis, followed by the synthesis of the membranous layer. It is noted that the late post molt stage C1-2 is difficult to distinguish from late B stage by these methods. When all layers of the cuticle have been produced in the cuticle, the shrimp is then designated as being in the intermolt stage (C3-4). At this time the thickness of the post-ecdysial layer (endocuticle and membranous layer), approximately 24 mm, was approximately twice that of the pre-ecdysial layer (epicuticle and exocuticle). Determination of the intermolt stage by physical examination is difficult, and can easily be misdiagnosed. Using the histological characteristics of the cuticle is a better way to determine the intermolt stage of the shrimp. We would like to emphasise here that observation of the membranous layer in the cuticle could indicate the level of maturity of the cuticle, hence the existence of the intermolt stage.

Taken together the results indicate that the biological processes of degradation and regeneration of the cuticle occur throughout the molting cycle. These processes include the syntheses and degradations of collagen fibers, protein, carbohydrate, calcium salt and lipid. Synthesis of collagen fibers, and carbohydrate occur constantly throughout the molting cycle, while glycoprotein and




Figure 4. Localisation of carbohydrate (a,b), collagen fibers (c,d), calcium salt (e,f), and lipid (g,h) in the cuticle during the early postmolt period (left column) and late postmolt period (right column). New epicuticle (nEp) and exocuticle (nEx) are now mature. The new endocuticle (nEd) is being synthesized in early postmolt period. It is noted that the stain for calcium salt in the endocuticle became more intense in the late postmolt stage (compare nEd in 4e and 4f). Bar = 10 μm.

lipid syntheses occur only at the end of the premolt stage. Deposition of calcium salt increases from the late premolt until the intermolt stage. This information may be useful for managing the feeding regimes to ensure that the shrimp's demand for a particular nutrient at a particular stage in the molting cycle is met. This could support maximum shrimp growth and health after each round of the molting cycle. We are continuing our investigations to uncover potential links between cuticular changes and other organs. This knowledge will help us to understand the processes that regulate the molting process in the black tiger shrimp.

Conclusions

1. The mature cuticle of the black tiger shrimp is composed of four sub-layers from the exterior to the interior: an epicuticle, an exocuticle, an endocuticle, and a membranous layer. Each sub-layer contains different organic and in-organic components. Carbohydrate and calcium salt is found in all sub-layers. Collagen fibers deposit in most sub-layers, except the epicuticle. Lipid deposits only in the epicuticle.

2. These cuticular sub-layers change chronologically throughout the molting cycle. Before ecdysis, the endocuticle and membranous layer of the old cuticle are degraded while the epicuticle and exocuticle are synthesized on the inside of the old cuticle. After the old cuticle is shed the new endocuticle and membranous layer are sequentially synthesized together with the depositions of carbohydrate, collagen fibers, calcium salt, and lipid. In the intermolt stage, syntheses of all the cuticular sub-layers are completed. The ratio of the thickness of the pre-ecdysial and post-ecdysial layers should now be approximately 1:2. (see Figure 5 for the summary).

3. The histological properties used to determine the stage of the molt are given below

- i) D0-2 (early premolt), lightly stained membranous layer

- ii) D3-4 (late premolt stage), the existence of new epicuticle and exocuticle on the inside of the old cuticle
- iii) A stage (early postmolt stage), the existence of an epicuticle and an exocuticle
- iv) B stage (late post molt) the existence of the developing endocuticle and
- v) C3-4 stage (intermolt stage), the existence of a membranous layer in the cuticle. The ratio of the thickness the pre-ecdysial : post-ecdysial cuticle is approximately 1:2.

Acknowledgements

We specially thank Professor Boonsirm Withyachumnarnkul (Centex, Mahidol University), and Pinij Thaweethamseewee (Department of Anatomy, PSU) for critical comments and support, and Dr. Brian Hodgson for his comprehensive comments on this manuscript. This work is financially supported by the Ministry of University Affairs, Thailand Research Fund (TRF), and the Department of Anatomy, Faculty of Science, Prince of Songkla University.

References

- Adams, E., Simkiss, K., and Taylor, M. 1982. Metal ion metabolism in the moulting crayfish (*Austropotamobius pallipes*). *Comp. Biochem. Physiol.* 72A : 73-76.
- Andersen, S.O. 1999. Exoskeletal proteins from the crab, *Cancer pagurus*. *Comp. Biochem. Physiol.* 123A : 203-211.
- Bancroft, J.D. and Gamble, M. 2002a. *Theory and Practice of Histological Techniques*. 5th ed. Harcourt publisher limited, London, p.153.
- Bancroft, J.D. and Gamble, M. 2002b. *Theory and Practice of Histological Techniques*. 5th ed. Harcourt publisher limited, London, p.175.
- Bancroft, J.D. and Gamble, M. 2002c. *Theory and Practice of Histological Techniques*. 5th ed. Harcourt publisher limited, London, p.258.




Figure 5. Summary of cuticular-degradation and -regeneration throughout the molting cycle. Early premolt (a), the endocuticle and membranous layer are being degenerated from the old cuticle. The new epicuticle and exocuticle are next synthesized (b). Immediately after ecdysis, both new epicuticle and exocuticle are present, while the old cuticle was shed (c). Early postmolt, the endocuticle is newly synthesized (d). End of postmolt, the newly secreted endocuticle becomes thicker (e). When the synthesis of the membranous layer is begun, the shrimp is said to be in the intermolt stage (f). Ed = endocuticle, Ep = epicuticle, Ex = exocuticle, MI = membranous layer, nEd = new endocuticle, nEp = new epicuticle, nEx = exocuticle, oEd = old endocuticle, and oEx = old exocuticle.

Carlisle, D.B. 1957. On the hormonal inhibition of moulting in Decapod Crustacea II. The terminal anecdyosis in crabs. J. Mar. Biol. Assoc. U.K. 36 : 291-307.

Compere, P., Morgan, J.A. and Goffinet, G. 1993. Ultrastructure location of calcium and magnesium during mineralization of the cuticle of the shore crab, as determined by the K-pyroantimonate method and X-ray microanalysis. Cell Tissue Res. 274 : 567-577.

Gilbert, L.I., Rybczynski, R., and Warren, J.T. 2002. Control and biochemical nature of the ecdysteroidogenic pathway. Ann. Rev. Entomol. 47 : 883-916.

Glynn, J.P. 1968. Studies on the ionic, protein and phosphate changes associated with the moult cycle of *Homarus vulgaris*. Comp. Biochem. Physiol. 26 : 937-946.

Gorokhova, E. 2002. Moulting cycle and its chronology in *Mysis mixta* and *Neomysis integer* (Crustacea,

- Mysidacea): implications for growth assessment. *J. Exp. Marine Biol. Ecol.* 278 : 179-194.
- Keller, R. and Schmid, E. 1979. *In vitro* secretion of ecdysteroids by Y-organs and lack of secretion by mandibular organs of the crayfish following molt induction. *J. Comp. Physiol.* 130 : 347-353.
- Lee, K.J., Watson, R.D., and Roer, R.D. 1998. Molt-inhibiting hormone mRNA levels and ecdysteroid titer during a molt cycle of the blue crab, *Callinectes sapidus*. *Biochem. Biophys. Res. Comm.* 249 : 624-627.
- Mangum, C.P. 1992. Physiological aspects of molting in the blue crab, *Callinectes sapidus*. *Amer. Zool.* 32 : 459-469.
- Marlowe, R.L., Dillaman, R.M. and Roer, R.D. 1994. Lectin binding by crustacean cuticle: the cuticle of *Callinectes sapidus* throughout the molt cycle, and the intermolt cuticle of *Procambarus clarkii* and *Ocypode quadrata*. *J. Crustacean Biol.* 14 : 231-246.
- Musgrove, R.J. 2000. Molt staging in the southern rock lobster *Jasus edwardsii*. *J. Crustacean Biol.* 20 : 44-53.
- Mykles, D.L. 1980. The mechanism of fluid absorption at ecdysis in the American lobster, *Homarus americanus*. *J. Exp. Biol.* 84 : 89-101.
- Nakatsuji, T. and Sonobe, H. 2004. Regulation of ecdysteroid secretion from the Y-organ by molt-inhibiting hormone in the American crayfish, *Procambarus clarkii*. *Gen. Comp. Endocrinol.* 135 : 358-364.
- Pierce, D.C., Butler, K.D., and Roer, R.D. 2001. Effects of exogenous *N*-acetylhexosaminidase on the structure and mineralization of the post-ecdysial exoskeleton of the blue crab, *Callinectes sapidus*. *Comp. Biochem. Physiol.* 128B : 691-700.
- Pratoomchat, B., Sawangwong, P., Guedes, R., De Iurdes Reis, M. and Machado, J. 2002b. Cuticle ultra-structures changes in the crab *Scylla serrata* over the molt cycle. *J. Exp. Zool.* 293 : 414-426.
- Pratoomchat, B., Sawangwong, P., Pakkong, P. and Machado, J. 2002a. Organic and inorganic compound variations in haemolymph, epidermal tissue and cuticle over the molt cycle in *Scylla serrata* (Decapoda). *Comp. Biochem. Physiol.* 131A : 243-255.
- Promwikorn, W., Kirirat, P. and Thaweethamseewee, P. 2004. Index of molting cycle in the black tiger shrimp (*Penaeus monodon*). *Songklanakarin J. Sci. Technol.* 26 : 765-772.
- Roer, R. and Dillaman, R. 1984. The structure and calcification of the crustacean cuticle. *Amer. Zool.* 24 : 893-909.
- Roer, R.D., Halbrook, K.E. and Shafer, T.H. 2001. Glycosidase activity in the post-ecdysial cuticle of the blue crab, *Callinectes sapidus*. *Comp. Biochem. Physiol.* 128B : 683-690.
- Skinner, D.M. 1985a. Interacting factors in the control of the crustacean molt cycle. *Amer. Zool.* 25 : 275-284.
- Skinner, D.M., 1985b. Molting and regeneration. In: Bliss, D.E. and Mantel, L.H. *The Biology of Crustacea*. Academic Press, New York, 9 : 43-128.
- Sheehan, D.C. and Hrapchak, B.B. 1980. *Theory and Practice of Histotechnology*. 2nd ed. The C.V. Mosby company, St. Louis, 205.
- Soumoff, C. and O'Connor, J.D. 1982. Repression of Y-organ secretory activity by molt inhibiting hormone in the crab *Pachygrapsus crassipes*. *Gen. Comp. Endocrinol.* 48 : 432-439.
- Spaziani, E., Mattson, M. and Wang, W.L. 1999. Signaling pathways for ecdysteroid hormone synthesis in crustacean Y-organs. *Amer. Zool.* 39 : 496-512.
- Stevenson, J.R. 1985. Dynamics of the integument. In: Bliss, D.E. and Mantel, L.H. *The biology of crustacea*. Academic Press, New York, 9 : 1-41.
- Vigh, D.A. and Dendinger, J.E. 1982. Temporal relationships of postmolt deposition of calcium, magnesium, chitin and protein in the cuticle of the Atlantic blue crab, *Callinectes sapidus* Rathbun. *Comp. Biochem. Physiol.* 72A : 365-369.
- Wang, W., Wang, A., Wang, D., Wang, L., Liu, Y. and Sun, R. 2003. Calcium, phosphorus and adenylate levels and Na⁺-K⁺-ATPase activities of prawn, *Macrobrachium nipponense*, during the moult cycle. *Comp. Biochem. Physiol.* 134A : 297-305.
- Watson, R.D., Lee, K.J., Qiu, S., Luo, M., Umphrey, H.R., Roer, R.D. and Spaziani, E. 2001. Molecular cloning, expression, and tissue distribution of Crustacean molt-inhibiting hormone. *Amer. Zool.* 41 : 407-417.
- Wheeler, A.P. and Sikes, C.S. 1984. Regulation of carbonate calcification by organic matrix. *Amer. Zool.* 24 : 933-944.