

References

- Ababouch, L., Afilal, M.N., Benadelijelil, H. and Busta, F.F. 1991. Quantitative changes in bacteria, amino acids and biogenic amines in sardine (*Sardina pilchardus*) stored at ambient temperature (25-28°C) and in ice. Int. J. Food Sci. Technol. 26: 297-306.
- Abe, H., Doson, G.P., Hoeger, H. and Parkhouse, W.S. 1985. Role of histidine-related compounds to intracellular buffering in fish skeletal muscle. Am. J. Physiol. Regul. Intergr. Comp. Physiol. 249: 449-454.
- Adegoke, G.O. and Odesola, B.A. 1996. Storage of maize and cowpea and inhibitory of microbial agents of biodeterioration using the powder and essential oil of lemon grass (*Cymbopogon citrates*). Int. Biodeter. Biodegra. 81-84.
- Ahmad, S. and Branen, A.L. 1981. Inhibition of mold growth by butylated hydroanisole. J. Food Sci. 46:1059-1063.
- Aksnes, A., Brekken, B. 1988. Tissue degradation, amino acid liberation and bacterial decomposition of bulk stored capelin. J. Sci. Food Agric. 45: 53-60.
- Al-Dagal, M.M. and Bazaraa, W.A. 1999. Extension of shelf-life of whole peeled shrimp with organic salts and bifidobacteria. J. Food Prot. 62: 51-56.
- Alemela L., Nieto-Snadoval, J.M. and Lopez, F.J.A. 2002. Microbial inactivation of paprika by a high temperature short-x time treatment. Influence on color properties. J. Agric. Food Chem. 50: 1435-1440.
- Alzamora, S.M., Tapia, M.S. and Lopez-Molaa, A. 2000. Minimally Processed Fruits and Vegetables. Fundamental Aspects and Applications. An Aspen Publication. Gaitherburg, Maryland.
- Anderson, U.B. Thomassen, M.S. and RØra, B.A.M. 1997. Texture properties of farmed rainbow trout (*Oncorhynchus mykiess*): effect of diet, muscle fat content and time of storage on ice. J. Sci. Food Agric. 74: 347-353.
- Ando, M., Nishiyabu, A., Tsukamasa, Y. and Makinodan, Y. 1999. Post-mortem softening of fish muscle during chilled storage as affected by bleeding. J. Food Sci. 64: 423-428.
- Ando, M., Toyahara, H. and Sakaguchi, M. 1992. Post-mortem tenderization of rainbow trout muscle caused by the disintegration of collagen fiber in the pericellular connective tissue. Bull. Jpn. Soc. Sci. Fish. 58: 567-570.

- Ando, M., Toyahara, H., Shimizu, Y. and Sakaguchi, M. 1991. Post-mortem tenderization of rainbow trout (*Oncorhynchus mykiss*) muscle cause by gradual disintegration of the extracellular matrix structure. *J. Sci Food Agric.* 55: 589-597.
- Ankri, S. and Mirelman, D. 1999. Antimicrobial properties of allicin from garlic. *A Rev. Microbes Infec.* 1: 125-129.
- Annon. 2003. <http://www.eximword.com/thaicookingcenter/matching.asp?item=1>
- Ashie, I.N.A. and Simpson, J.P. 1998. Effects of pressurization and refrigerated storage on microstructure of fish muscle tissue. *J. Muscle Foods.* 9: 193-199.
- Ashie, I.N.A., Smith, J.P. and Ramaswamy, H.S. 1997. Changes in texture during storage. *J. Muscle Foods.* 8:13-32.
- Ashie, I.N.A., Smith, J.P. and Simpson, B.K. 1996. Spoilage and shelf-life extension of fresh fish and shellfish. *Crit. Rev. Food Sci. Technol.* 36: 67-121.
- Aureli, P., Constantini, A. and Zolea, S. 1992. Antimicrobial activity of some plant essential oils against *Listeria monocytogenes*. *J. Food Sci.* 55: 344-348.
- Bailey, A.J. and Light, N.D. 1989. Connective Tissue. *In* Meat and Meat Products. London. Elsevier Applied Science.
- Bak, L.S., Andersen, A.B., E.M. and Bertlsen, G. 1999. Effect of modified atmosphere packaging on oxidative changes in frozen stored coldwater shrimp (*Pandalus borealis*). *Food Chem.* 64: 169-175.
- Banks, H., Nickelson, R. II. and Finne, G. 1980. Shelf-life studies on carbon dioxide packaged finfish from the gulf of Mexico. *J. Food Sci.* 45: 157-162.
- Barnett, H.J., Conrad, J.W. and Nelson, R.W. 1987. Use of laminated high and low polyethylene flexible packaging to store trout (*Salmo gairdneri*) in modified atmosphere. *J. Food Prot.* 50: 643-651.
- Barr, A. 1990. Consumer motivational forces affecting the sale of light dairy products. *Food Technol.* 44: 97-98.
- Belitz, H.D. and Grosch, W. 1999. Lipids. *In* Food Chemistry. Heidelberg, Germany. Springer-Verlag. Pp. 184-185.
- Ben-gigirey, B., De Sousa, V.B., Villa, T.G. and Barros-velazquez, J. 1999. Chemical changes and visual appearance of albacore tuna as related to frozen storage. *J. Food Sci.* 64: 20-24.

- Bhaskarachary, K., Roa, D.S.S., Deoshale, Y.G. and Reddy, V. 1995. Carotene content of some common and familiar foods of plant origin. *Food Chem.* 54: 189-193.
- Bito, M., Yamada, K., Nikumo, Y. and Amano, K. 1983. Studies on rigor among some varieties of fish by modified. Cutting's method. *Bull. Tokai Reg. Fish. Res. Lab.* 109: 89-96.
- Board, P.G. and Gould, G.W. 1991. Future Prospects. *In* Food Preservative (N.J. Russel and G.W. Gould eds.). Glasgow: Blackie and Son. 267-284.
- Boerema, J.A., Penney, N., Cummings, T.L. and Bell, G. 1993. Carbon dioxide controlled atmosphere packaging of slice ham. *Int. J. Food Sci. Technol.* 25: 435-442.
- Bone, Q. 1978. Locomotor muscle. *In* Fish Physiology. Vol. VII (W.S. Hoar and D.J. Randall eds.). Academic Press New York. London. 361-424.
- Boonyaratpalin, M. 1994. Asian Seabass, *Lates calcarifer*. *In* CRC, Handbook of Nutrient Requirement of Finfish. (R.P.Wilson ed.). CRC. Press. Boston London.
- Botta, J.R. 1995. Evaluation of Seafood Freshness Quality. VCH Publisher Inc.
- Botta, J.R., Laude, J.L. and Jewer, M.A. 1984. Effect of methodology on total volatile basic nitrogen (TVB-N) determination as an index of quality of fresh Atlantic cod (*Gadus morhua*). *J. Food Sci.* 49: 734-736, 750.
- Brandes, C.H. and Dietrich, R. 1970. The Chemical Biology of Fishes (R.M. Love ed.). Academic Press. London and New York.
- Branen, A.L., Davidson, P.M. and Katz, B. 1980. Antimicrobial properties of phenolic antioxidants and lipids. *Food Technol.* 34: 42-53.
- Breithaupt, D.E. and Bemedi, A. 2001. Carotenoid esters in vegetables and fruits. Screening with emphasis on β - cryptoxanthin esters. *J. Agric. Food Chem.* 49: 2064-2070.
- Bremner, H.A. 1992. Fish flesh structure and the role of collagen-its post-mortem aspects and implications for fish processing. *In* Quality Assurance in the Fish Industry (H.H. Huss., J.K. Jakosen and J. Liston eds.). Pp. 39-62. Elsevier Science Publishers Amsterdam.
- Bremner, H.A. and Hallett, I.C. 1985. Muscle connective tissue junctions in the fish blue grenadier (*Macruronus novaezelandiae*): a scanning electron microscope study. *J. Food Sci.* 50: 975-980.

- Bremner, H.A. and Statham, J.A. 1987. Packaging in CO₂ extends shelf-life of scallops. *Food Technol. Austral.* 39: 177-179.
- Buege, J.A. and Aust, S.D. 1978. Microsomal lipid peroxidation. *Method Enzymol.* 52: 302-304.
- Cann, D.C. 1988. Modified atmosphere packaging of fishery products. *Infofish Int.* 1: 37-39.
- Cann, D.C., Houston, N.C., Taylor, L.Y., Smith, G.L., Thomson, A.B. and Craig, A. 1984. *Studies of Salmonids Packed and Stored Under a Modified Atmosphere.* Aberdeen: Torry Research Station.
- Cann, D.C., Smith, G.L. and Houston, N.C. 1983. *Further Studies on Marine Fish Stored Under Modified Atmosphere Packaging.* Aberdeen: Torry Research Station.
- Capplein, G. and Jessen, F. 2002. ATP, IMP and glycogen in cod muscle at onset and during development of rigor mortis depend on the sampling location. *J. Food Sci.* 67: 991-995.
- Carpene, E., Veggetti, A. and Mascarello, F. 1982. Histochemical fibre types in the lateral muscle of fishes in fresh, brackish and salt water. *J. Fish Biol.* 20: 379-396.
- Cha, A.S. and Ho, C.T. 1988. Studies of the interaction between aspartame and flavor vanillin by high performance liquid chromatography. *J. Food Sci.* 53: 562-564.
- Church, N. 1994. Developments in modified atmosphere packaging and related technologies. *Trends Food Sci. Technol.* 5: 345-352.
- Church, I.J. and Parsons, A.L. 1995. Modified atmosphere packaging technology: a review. *J. Sci. Food Agric.* 67: 143-152.
- Colby, J.W., Enriquez-Ibarra, L. and Flick, G.J. Jr. 1993. Shelf-life of fish and shellfish. *In Shelf-Life Studies of Foods and Beverages-Chemical, Biological, Physical and Nutritional Aspects* (G. Charalambous ed.). Pp. 85-143. Amsterdam: Elsevier.
- Connell, J.J. 1975. *Control of Fish Quality.* London
- Connell, J.J. and Howgate, P.F. 1986. Fish and fish products. *In Quality Control in the Food Industry* (S.M. Herschdoerfer ed.). Vol. 2. Academic Press.
- Conway, E.T. 1950. *Microdiffusion and volumetric error.* Crosby Lack Wood and Son. Ltd., London.

- Cornforth, D.P. and West, E.M. 2002. Evaluation of the antioxidant effect of dried milk mineral in cooked beef, pork, and turkey. *J. Food Sci.* 67: 615-617.
- Cornell, D.G., De Vilbiss, E.D. and Pallansch, M.J. 1971. Binding of antioxidants by milk proteins. *J. Food Sci.* 54: 634-637.
- Coyne, F.P. 1933. The effect of carbon dioxide on bacterial growth with special reference to the preservation of fish. Part II. *J. Soc. Chem. Ind.* 52: 19T-24T.
- Dalgaard, P., Gram, L. and Huss, H.H. 1993. Spoilage and shelf-life of cod fillets packed in vacuum or modified atmospheres. *Int. J. Food Microbiol.* 19: 283-294.
- D' Amelio, F.S. 1999. *Botanicals: a Phytocosmetic Desk Reference*. CRC Press., Boca Raton, Washington, D.C.
- Deis, R. 1999. *Food product Design: The Secret of World of Spices*. <http://www.foodproductdesign.com/archive/1999/0899cs.html>.
- Deli, J. and Toth, G. 1997. Carotenoid composition of the fruits of *Capsicum annum* cv. Bovet 4 during ripening. Original paper. *Z Lebensm Unters Forsch A.* 205:388-391.
- Deli, J., Matus, Z. and Toth, G. 1996. Carotenoid composition in the fruit of *Capsicum annum* cv. Szentesi Kosszarvu during ripening. *J. Agric. Food Chem.* 44: 711-716.
- Deli, J., Matus, Z., Molnar, P. and Toth, G. 2001. Separation and identification of beta-carotenoids from different coloured paprika (*Capsicum annum*) by reversed-phase high performance liquid chromatography. Original paper. *Eur. Food Res. Technol.* 213: 301-305.
- Dhananjaya, S. and Stroud, G.D. 1994. Chemical and sensory changes in haddock and herring stored under modified atmosphere. *Int. J. Food Sci. Technol.* 29: 575-583.
- Division of Health Statistics. 1989. Office of the Permanent Secretary, Ministry of Public Health. *Public Health Statistic*. Thailand.
- Dore, I. 1991. *Fish and Shellfish Quality Assessment. A guide for retailers and restauranters*. An Osprey Book Published by Van Nostrand Reinhold. New York.

- Drosinos, E.H. and Board, G.R. 1995. A survey of minced lamb packaged in modified atmospheres. *Fleisch*. 75: 281-284.
- Dunajski, E. 1979. Texture of fish muscle. *J. Text. Stud.* 10: 301-308.
- Faergemand, J., Ronsholdt, B., Alsted, N., and Borresen. T. 1995. Fillet texture of rainbow trout as affect by feeding stategy, slaughtering procedure and storage post-mortem. *Water Sci. Technol.* 10: 225-231.
- Farid, M., Bal'A, M.F.A. and Marshall, D.L. 1998. Organic acid dipping of catfish fillets: effect on color, microbial load and *Listeria monocytogenes*. *J. Food Prot.* 61: 1470-1484.
- Fauconneau, B., Alami-Durante, H., Laroche, M., Marcel, J. and Vallot, D. 1995. Growth and meat quality relations in carp. *Aquacult.* 129: 265-267.
- Fauconneau, B., Chmaitilly, J., Andre, S., Cardinal, M., Cornet. J., Vallet, J.L., Dumont, J.P. and Laroche, M. 1993. Characteristics of rainbow trout flesh: I. Chemical composition and cellularity of muscle and adipose tissue. *Sciences des Aliments*. 13: 173-187.
- Fennema, O.R. 1996. *Food Chemistry*. 3th edition. Macel Dekker, Inc. New York.
- Ferdouse, F. 1995. Cultures seabass market. *Infofish Int.* 2: 20-22.
- Fletcher, G.C., Hallett, I.C., Jerrett, A.R. and Holland, A.J. 1997. Changes in the fine structure of the myocommata muscle fiber junction related to gaping in rested and exercised muscle from king salmon (*Oncorhynchus tshawytscha*). *Lebensm. – Wiss. Technol.* 30: 246-252.
- Frankel, E.N. 1998. *Lipid Oxidation*. The Oil Press.
- Fujii, T., Hirayama, M., Okuzumi, M., Yasuda, M., Nishino, H. and Yokoyama, M. 1989. Shelf-life studies on fresh sardine packaged with carbon dioxide-nitrogen gas mixture. *Nipp. Sui. Gakk.* 55: 1971-1975.
- Fung, D.Y.C. 1995. Rapid Methods and Automation in Food Microbiology and Update. 3 *In* *Microbiological Control for Food and Agricultural Products*. (C.M. Bourgeois, and Leveau eds.) VCH Publishers. U.K
- Fung, D.Y.C., Lin, C.C. and Gailani, M.B. 1985. Effect of phenolic: antioxidants on microbiology growth. *Crit. Rev. Microbiol.* 12: 153-183.
- German, J.B. and Kinsella, J.E. 1985. Lipid oxidation in fish tissue. Enzymatic initiation via lipoxygenase. *J. Agric. Food Chem.* 33: 680-683.

- German, J.B. and Kinsella, J.E. 1996. Hydroperoxide metabolism in trout gill tissue. Effect of glutathione on lipoxygenase products generated from arachidonic acid and docosahexaenoic acid. *Biochem Biophys Acta.* 879: 378.
- Gill, H.S., Weatherley, A.H. and Bhesania, T. 1982. Histochemical characterization of myotomal muscle in the bluntnose minnow, *Pimephales notatus*. *Rafinesque. J. Fish Biol.* 21: 205-214.
- Gill, H.S., Weatherley, A.H., Lee, R. and Legere, D. 1989. Histochemical characterization myotomal muscle of five teleost speices. *J. Fish Biol.* 34: 375-386.
- Gill, T. 2000. Nucleotide Degrading Enzymes. *In* Utilization and Influence of Postharvest Seafood Quality ((N.F. Haard and B.K. Simpson eds.). Macel Dekker, Inc. New York.
- Gram, L. and Huss, H.H. 1996. Microbiological spoilage of fish and fish products. *Int. J. Food Microbiol.* 33: 121-137.
- Granado, F., Olmedilla, B., Gil-Martinez E. and Blanco, I. 2001. A Fast, reliable and low-cost saponification protocol for analysis of carotenoids in vegetables. Original paper. *J. Food Comp. Anal.* 14: 479-489.
- Gray, J.I and Monahan, F.J. 1992. Measurement of lipid oxidation in meat and meat products. *Trends Food Sci. Technol.* 3: 315-319.
- Gross, K.L., Bollinger., R., Thawnghmung, P. and Coliings, G. F. 1994. Effect of 3 different preservative systems on stability of extrudated dog food subjected to ambient and high temperature storage. *J. Nutri.* 124 (12S): 2638S-2642S.
- Guldager, H.S., Bøknæs, N., østerberg, C., Nielsen, J. and Dalgaard, P. 1998. Thawed cod fillets spoil less rapidly than unfrozen fillets when stored under modified atmosphere at 2°C. *J. Food Prot.* 61: 1129-1136.
- Haard, N.F. 2000. Seafood enzymes. The Role of Adaptation and Other Intraspecific Factors. *In* Seafood Enzymes Utilization and Influence of Postharvest Seafood Quality (N.F. Haard and B.K. Simpson eds.). Marcel Dekker, Inc. New York. Basel.
- Handumrongkul, C. and Silva, J.L. 1994. Aerobic counts, color and adenine nucleotide changes in CO₂ packed refrigerated striped bass strips. *J. Food Sci.* 59: 67-69.

- Hart, D.J. and Scott, K.J. 1995. Development and evaluation of an HPLC method for the analysis of carotenoid content of vegetables and fruits commonly consumed in UK. *Food Chem.* 54: 101-111.
- Hatae, K., Tobimatsu, A., Takeyama, M. and Matsumoto, J.J. 1986. Contribution of connective tissue on the texture difference of various fish species. *Bull. Jpn. Soc. Sci. Fish* 52: 2001-2007.
- Hatae, K., Yoshimatsu, F. and Matsumoto, J.J. 1990. Role of muscle fibers in contributing firmness of cooked fish. *J. Food Sci.* 55: 693-696.
- Heath, A.G. 1995. *Water Pollution and Fish Physiology*. CRC. Lewis Publishers.
- Heinonen, M.L., Ollilainen, V., Linkola, E. K., Varo, P.T. and Koivistoinen, P.E. 1998. Carotenoids in finish foods: vegetables, fruits and berries. *J. Agric. Food Chem.* 37: 655-659.
- Heisler, N. 1986. Acids-base regulation in fishes. *In* *Acid-Base Regulation in Animals* (N. Heisler ed.). Elsevier, Amsterdam. Pp. 309-356.
- Hildebrand, M. 1988. *Analysis of Vertebrate Structure*. John Wiley and Sons, Inc.
- Hoffman, L.C. and Prinsloo, J.E. 1990. A comparison of the dressout percentage of the red and normal colored strains of the African Sharptooth catfish, *Clarias gariepirus* (Bruchell). *The SA J. Food Sci. Nutr.* 2: 35-38.
- Hoffman, L.C., Prinsloo, J.E., Casey, N.H. and Theron, J. 1994. The anatomical heterogeneity in the proximate compositions of muscle of the African Sharptooth catfish, *Clarias gariepirus* (Bruchell). *The SA. J. Food Sci. Nutr.* 6: 30-35.
- Hoffman, L.C., Prinsloo, J.E., Casey, N.H. and Theron, J. 1995. The intrinsic variation in the chemical composition of the African Sharptooth catfish, *Clarias gariepirus* (Bruchell). I. Lipid, amino acid and mineral composition of dark and light muscle. *The SA. J. Food Sci. Nutr.* 7: 13-17.
- Hong, L.C., Leblanc, E.L., Hawrysh, Z.J. and Hardin, R.T. 1996. Quality of Atlantic mackerel (*Scomber scombrus* L.) fillets during modified atmosphere storage. *J. Food Sci.* 61: 646-651.
- Hornero-Moendez D. and Minguez-Mosquera M.I. 2000. Xanthophyll esterification accompanying carotenoid over accumulation in chromopast of *Capsicum annuum* ripening fruits is constitutive process and useful for ripeness index. *J. Agric. Food Chem.* 48: 1617-1622.

- Hornero-Mendez, D., de Guevara, R. G-L. and Minguéz-Mosquera M.I. 2000. Carotenoid biosynthesis changes in five red pepper (*Capsicum annuum*) cultivars during ripening, cultivar selection for breeding. *J. Agric. Food Chem.* 48: 3857-3864.
- Howard, L.R., Talcott S.T., Brenes, C.H. and Villalon B. 2000. Changes in phytochemical and antioxidant activity of selected pepper cultivars (*Capsicum species*) as influenced by maturity. *J. Agric. Food Chem.* 48: 1713-1720.
- Howgate, P. 1977. *Sensory Properties of Foods.* (G.G. Birch., J.G. Brennan. and K.J. Parker eds.) Applied Science Publishers, London.
- Howgate, P. 1986. Approaches to the definition and measurement of the storage life of chilled and frozen fish. a brief review. *Refriger. Sci. Technol.* 60: 45-53.
- Hsieh, R.J. and Kinsella, J.E. 1989. Lipoxygenase generation of specific volatile carbonyl compounds in fish tissue. *J. Agric. Food Chem.* 37: 279-286.
- Huis in't Veld, J.H.J. 1996 Microbial and biochemical spoilage of foods: an overview. *Int. J. Food Microbiol.* 33: 1-18.
- Hultin, H.O. 1985. Characteristics of muscle tissue. *In Food Chemistry* (O.R. Fennema ed.). New York. Marcel Dekker Inc. Pp. 725-789
- Hultin, H.O. 1988. Potential lipid oxidation problem in fatty fish processing. *In Fatty utilization: upgrading from feed to food, proceeding of a national technical conference* (N. Davis ed.). Raleigh, NC: UNC Sea grant college program. Pp. 185-233.
- Hurling, R., Rodell, J.B. and Hunt, H.D. 1996. Fiber diameter and fish texture. *J. Text. Stud.* 27: 679-685.
- Huss, H.H., 1995. Quality and Quality Changes in Fresh Fish. Fisheries Technical Paper no. 348. Rome, Italy: FAO.
- Huss, H.H., Dalgaard, P. and Gram, L. 1997. Microbiology of fish and fish products. *In: Seafood from Producer to Consumer: Integrated Approach to Quality* (J.B. Luten, T. Børresen and J. Oehlenschläger eds.). Pp. 413-430. Amsterdam: Elsevier.
- Hussein, A., Odumeru, J.A., Ayanbadejo, T., Faulkner., H., McNab, W.B., Hager, H. and Szijarto, L. 2000. Effects of processing and packaging on vitamin C and β -carotene content of ready - to - use (RTU) vegetables. *Food Res. Int.* 33: 131-136.

- Hussein, M.M., D'Amelia, R.P., Manz, A.L and Jacin, H. 1984. Determination of reactivity of aspartame with flavor aldehydes by gas chromatography, HPLC and GC. *J. Food Sci.* 49: 520-524.
- Hutabarat, L.S., Mulholland, M. and Greenfield, H. 1998. Development and validation of an isocratic high-performance liquid chromatographic method for quantitative determination of phytoestrogens in soya bean. Short communication. *J. Chromat. A.* 795: 377-384. *In* The properties of water in foods. ISOPOW 6. (D.S. Reid ed.). Backie Academic and Professional.
- Inoue, K., Morioka, K., Shioya, I., Mitsuboshi, T., Itoh, Y., Obtake, A. and Satake, T. 1998a. Short-term rearing of red sea bream in deep seawater and its effects on rigor mortis. *Fish. Sci.* 64: 789-803.
- Inoue, K., Morioka, K., Shioya, I., Mitsuboshi, T., Itoh, Y., Obtake, A. and Satake, T. 1998b. Effect of deep seawater acclimation on components and texture of the muscle red sea bream. *Fish. Sci.* 64: 804-807.
- Iwamoto, M., Yamanaka, H., Watanabe, S. and Hashimoto, K. 1987. Effect of storage temperature on rigor-mortis and ATP degradation in plaice *Paralichthys olivaceous* muscle. *J. Food Sci.* 52: 1514-1517.
- Iwamoto, M., Yamanaka, H., Watanabe, S. and Hashimoto, K. 1991. Changes in ATP and related breakdown compound in the adductor muscle of Itayagai scallop *Pecten albicans* during storage at various temperatures. *Nipp. Sui. Gakk.* 57: 153-156.
- Izquierdo-Pulido, M.L., Hatae, K. and Hoard, N.F. 1992. Nucleotide catabolism and changes in texture indices during ice storage of culture sturgeon, *Acipenser transmontranus*. *J. Food Biochem.* 16: 173-192.
- Jabarsyah, A., Tsuchimoto, M., Kozuru, Y., Misina, T., Yada, O. and Tachibana, K. 1999. The influence of pink muscle fiber in ordinary muscle of fishes on the rigor mortis progress. *Fish. Sci.* 65: 472-477.
- Jamora, J.J. and Rhee, K.S. 2002. Storage stability of extruded products from blends of meat and non meat ingredients: evaluation methods and antioxidative effects of onion, carrot, and oat ingredients. *J. Food Sci.* 67: 1645-1654.
- Jensen, M.H., Petersen, A., Røge, E.H. and Jepsen, A. 1980. Storage of chilled cod under vacuum and at various concentrations of carbon dioxide. *In* Advances in Fish Science and Technology (J.J. Connell ed). Pp. 294-297. Farnham, Surrey: Fish News Books.

- Jerret, A.J., Stevens, J. and Holland, A.J. 1996a. Tensile properties of white muscle in rested and exhausted king salmon (*Oncorhynchus mykiss*) (Walbaum). *Aquacult. Res.* 29: 89-101.
- Jerret, A.J., Stevens, J. and Holland, A.J. 1996b. Tensile properties of white muscle in rested and exhausted king salmon (*Oncorhynchus tshawytscha*). *J. Food Sci.* 61: 527-532.
- Johnston, I.A. 1999. Muscle development and growth : potential implications for fresh quality in fish. *Aquacult.* 177: 99-115.
- Johnston, I.A., Alderson, R., Sandham, C., Dingwall, A., Mitchell, D. Selkirk, C., Nickell, D., Baker, R., Robertson, B., Whyte, D. and Springate, J. 2000. Muscle fibre density in relation to the colour and texture of smoked Atlantic salmon (*Salmon salar* L.). *Aquacult.* 189: 335-349.
- Johnston, I.A. and Maitland, B. 1980. Temperature acclimation in crucian carp, *Carassius. L.*, morphometric analysis of muscle fibre ultrastructure. *J. Fish Biol.* 17: 113-125.
- Johnston, L.A. 1982. Physiology of muscle in hatchery raised fish. *Comp. Biochem. Physiol* 73B. 105-124.
- Johnston, L.A., Ward, P.S. and Goldspink, G. 1975. Studies on the swimming musculature of rainbow trout I. Fibre types. *J. Fish Biol.* 7: 451-458.
- Jones N.L., Shabib, S. and Sherman, P.M. 1997. Capsaicin as an inhibitory the growth of gastric pathogen *Helicobacter pylori*. *FEMS Microbiol. Let.* 146: 227-233.
- Juven, B.J., Kanner, J., Schved, F. and Weisslowicz, H. 1994. Factors that interact with the antibacterial action of thyme essential oil and its active constituents. *J. App. Bacteriol.* 76: 626-631.
- Kabara, J.J. Eklund, T. 1991. Organic acids and esters. *In* Food Preservative (N.J. Russel and G.W. Gould eds.). Glasgow: Blackie and Son Ltd. Pp. 44-71.
- Kakouri, A. and Nychas, G.J.E. 1994. Storage of poultry meat under modified atmosphere or vacuum packs: possible role of microbial metabolites as indicator of spoilage. *J. App. Bacteriol.* 76: 163-172.
- Kiessling, A., Johansson, L. and Storebakken, T. 1989. Effect of reduced feed ratio levels on fat content and fatty acid composition in white and red muscle from rainbow trout. *Aquacult.* 79: 169-175.

- Kolakowski, E. 1986. Changes of non protein nitrogen fractions in Antarctic krill (*Euphausia superba* Dana) during storage at 3°C and 20°C. *Z. Lebensm Unters Forsch.* 183: 421-425.
- Konosu, S. and Yamaguchi, K. 1982. The flavor components in fish and shellfish. *In* Chemistry and Biochemistry of Marine Food Products (R.E. Martin ed.). Pp. 367-404. The AVI Publishing Company, Inc. USA.
- Konosu, S., Watanabe, K. and Shimizu, T. 1974. Distribution of nitrogenous constituents in the muscle extracts of eight species of fish. *Nipp. Sui. Gakk.* 40: 909-915.
- Kosugi, H., Kato, T. and Kikugawa, K. 1987. Formulation of yellow, orange, and red pigments in the reaction of alk-2-enals with 2-thiobarbituric acid. *Anal. Biochem.* 165: 454-464.
- Kuo, J-D., Hultin, H.O., Peleg, M. and Atallah, M.T. 1991. *J. Food Process. Preserv.* 15: 125-133.
- Labuza, T.P. 1996. An introduction to active packaging for foods. *Food Technol.* 50: 68-71.
- Lannelongue, M., Finne, G., Hanna, M.O. Nickelson, R.II and Vanderzant, C. 1982. Microbiological and chemical changes during storage of swordfish (*Xiphias gladius*) steaks in retail packages containing CO₂-enriched atmospheres. *J. Food Prot.* 45: 1197-1203.
- Layrisse, M.E. and Matches, J.R. 1984. Microbiological and chemical changes of spotted shrimp (*Pandalus platyceros*) stored under modified atmosphere. *J. Food Prot.* 47: 453-457.
- Lee, B. J.Hendricks, D.G. and Cornforth, D.P. 1999. A comparison of carnosine and ascorbic acid on color and lipid stability in ground beef patti model system. *Meat Sci.* 51: 245-253.
- Lees, R. 1975. Food Analysis and quality control methods for the food manufacture and buyer. London Hill Books.
- Lie, O. 2001. Quality-Flesh the role of nutrition. *Aquacul. Res.* 32(suppl. 1). 341-348.
- Limroongreungrat, K. 2001. Development of Pasta Production using Sweetpotato Flour. PhD. Dissertation. University of Georgia. Athens, GA.
- Lin, C-C. and Liang, J-H. 2002. Effect of antioxidants on the oxidative stability of chicken breast meat in a dispersion system. *J. Food Sci.* 67: 530-533.

- Liston, J. 1980a. Fish and Shellfish and their Products. *In* Microbial. International Commission on the Microbiological Specifications of Food Academic Pres. New York
- Liston, J. 1980b. Microbiology in fisheries science. *In* Advances in Fish Science and Technology (J.J. Connell ed.). Pp. 138-157. Farham, Surrey: Fishing News Books.
- Lopez-Malo, A and Guerrero, S. 2000. Natural Antimicrobials from Plants *In* Minimally Processed Fruits and Vegetables: Fundamental Aspects and Application. (S.M. Alzamora, M.S. Tapia and A. Lopez-Molo eds.). An Aspen Publication. Aspen Publishers, Inc. Gaithersburg, Maryland.
- Love, R.M. 1970. The Chemical Biology of Fishes. Vol. 1. Academic Press New York.
- Love, R.M. 1988. The Food Fishes: Their Intrinsic Variation and Practical Implications. London. Farrand. Press.
- Love, R.M., Robertson, J., Smith, G.L. and Whittle, K. 1974. The texture of cod muscle. *J. Text. Stud.* 5: 201-212.
- Lyhs, U., Lahtinen, J., Fredriksson-Ahomaa, M., Hyytia-Trees, E., Elfin, K. and Korkeala, H. 2001. Microbiological quality and shelf-life of vacuum-packaged "gravad" rainbow trout stored at 3 and 8°C. *Int. J. Food Microbiol.* 70: 221-230.
- MacDonald, R.E. and Hultin, H.O. 1987. Some characteristics of the enzymic peroxidation system in the microsomal fraction of flounder skeletal muscle. *J. Food Sci.* 52: 15-21,27
- Mackeen, M.M., Ali, A.M., El-Sharkawy, S.H., Mnap, M.Y., Salleh, K.M., Lajis, N.H. and Kawazu, K. 1997. Antimicrobial and cytotoxic properties of some Malaysian traditional vegetable (Ulam). *Int. J. Phar.* 35: 174-178.
- Mai, J and Kinsella, J.E. 1979. Changes in lipid composition of cooked mince carp (*Cyprinus carpio*) during frozen storage. *J. Food Sci.* 44: 1619-1624.
- Martinez-Valverde, I., Periago, M. and Chesson, A. 2002. Phenolic compounds, lycopene and antioxidant activity in commercial varieties of tomato (*Lycopersicum esculentum*). Research Article. *J. Sci. Food Agric.* 82: 323-330.

- Matches, J.R. 1982. Effect of temperature on the composition of Pacific coast shrimp (*Pandalus jordani*). J. Food Sci. 47: 1044-1069.
- Matches, J.R. and Layrisse, M.E. 1985. Controlled atmosphere storage of spotted shrimp (*Pandalus platyceros*). J. Food Prot. 48: 709-711.
- Miller, A., Scanlan, R.A., Lee, J.S. and Libbey, L.M. 1973a. Identification of the volatile compounds produced in sterile fish muscle (*Sebastes melanops*) by *Pseudomonas fragi*. App. Microbiol. 25: 952-955.
- Miller, A., Scanlan, R.A., Lee, J.S. and Libbey, L.M. 1973b. Identification of the volatile compounds produced in sterile fish muscle (*Sebastes melanops*) by *Pseudomonas putrefaciens*, *Pseudomonas fluorescens* and *Achromobacter* species. App. Microbiol. 26: 18-21.
- Minguez-Mosquera, M.I. and Hornero-Mendez, D. 1994. Comparative study of the effect of paprika processing on the carotenoids in peppers (*Capsicum annum*) of the Bola and Agridulce varieties. J. Agric. Food Chem. 42: 1555-1560.
- Miron, T., Rabinkov, A., Mirelman, D., Wilchek, M. and Weiner, L. 2000. The mode of action of allicin : its ready permeability through phospholipid membranes may contribute to its biological activity. Biochem. Biophys. Acta. 1463: 20-30.
- Mishra, A.K. and Dubey, N.K. 1994. Evaluation of some essential oils for their toxicity against fungicausing deterioration of stored food commodities. App. Microbiol. 60: 1101-1105.
- Miyagawa, M., Tabuchi, Y., Yamane, K., Matsuda, H., Watabe, S., Hashimoto, K., Katakai, R. and Otsuka, Y. 1990. Changes in the free amino acid profile of snow crab (*Chionoetes opilio*) muscle during storage in ice. Agric. Biol. Chem. 54: 359-364.
- Mohri, S., Cho, S-Y., Endo, Y. and Fujimoto, K. 1992. Linoleate 13 s)-lipoxygenase in sardine skin. J. Agric. Food Chem. 40: 573-576.
- Montero, P. and Borderias, J. 1990. Influence of age on muscle connective-tissue in trout (*Salmo irideus*) J. Sci. Food Agric. 51: 261-269.
- Morkore, T., Hansen, A.A. Unander, E. and Einen, O. 2002. Composition, liquid leakage and mechanical properties of firming rainbow trout: variation between fillet sections and the impact of ice and frozen storage. J. Food Sci. 67: 1933-1938.

- Murakami, A., Kondo, A., Nakamura, Y., Ohigashi, H. and Koshimizu, K. 1993. Possible anti-tumor promoting properties of edible plants from Thailand and identification of an active constituent, cardomonin (2',4' dihydroxy-6'methoxychalcone), of *Boesenbergia pandurata*. Biosci. Biotechnol. Biochem. 1971- 1973.
- Murata, M. and Sakaguchi, M. 1986. Changes in contents of free amino acids, trimethylamine and nonprotein nitrogen of oyster during ice storage. Nipp. Sui. Gakk. 52: 1975-1980.
- Nakayama, T., Ooguchi, N and Ooi, A. 1999. Changes in rigor mortis of res seabream dependent on season and killing method. Fish. Sci. 65: 284-290.
- Ng, P.K.W., Hoehm, E. and Bushuk, W. 1989a. Binding of vanillin by fababeans proteins. J. Food Sci. 54: 105-107.
- Ng, P.K.W., Hoehm, E. and Bushuk, W. 1989b. Sensory evaluation of binding of fababeans proteins. J. Food Sci. 54: 324-325,346.
- Nassos, P.S., King Jr., A.D. and Stafford, A.E. 1983. Lactic acid concentration and bacterial spoilage in ground beef. App. Envir. Microbiol. 49: 894-900.
- National Committee for Clinical Laboratory Stands. 2000. Performance stands for antimicrobial disk susceptibility. Approved stand 7th ed. NCCLS document M2-A-7. Wayne, PA: NCCLS.
- Nawar, W.W. 1996. Lipids. *In* Food Chemistry. (O.W. Fennema ed.). Macel Dekker, Inc. New York.
- Newsome, R.L., Langlois, B.E., Moody, W.G., Gay, N. and Fox, J.D. 1984. Effect of time and aging on the microbiological quality of beef loins and corresponding steaks. J. Food Prot. 47: 122-125.
- Nishimoto, M., Mizuta, A. and Yoshinaka, R. 2003. Charaterization of molecular species of collagen in muscles of Japanese amberjack, *Seriola quinqueradiata*. Food Chem. 1-6.
- Nishimura, H., Takahashi, T., Wijaya, C.H., Satoh, A. and Ariga, T. 2000. Thermochemical transformation of sulfur compounds in Japanese domestic *Allium*, *Allium victoralis* L. Bio. Factors. 13: 257-263.
- Nychas, G.J.E. 1995. Natural Antimicrobials from Microorganisms. *In* New Methods of Food Preservation. (G.W. Gould ed.). Blackie Academic and Professional. Glasgow.

- Oberlender, V., Hanna, M.O., Miget, R., Vanderzant, C. and Finne, G. 1983. Storage characteristics of fresh swordfish seaks stored in carbon dioxide-enriched controlled (flow-through) atmosphere. *J. Food Prot.* 46: 434-440.
- Ofstad, R., Egelanddal, B., Kidman, S., Myklebust, R., Olsen, R.L. and Hermansson, A-M. 1996. Liquid loss as affected by post-mortem ultrastructural changes in fish muscle; cod (*Gadus morhua* L) and salmon (*Salmo salar*). *J. Sci. Food Agric.* 71: 301-312.
- Oliver, J., Palou, A. and Pons A. 1998. Semi-quantification of carotenoids by high – performance liquid chromatography: saponification-induced losses in fatty foods. Short communication. *J. Chromat A.* 829: 393-399.
- Olsson, G.B., Olsen, R.L and Ofstad, R. 2003. Postmortem structural characteristics and water- holding capacity in Atlantic halibut muscle. *LWT.* 36: 125-133.
- Ordóñez, J.A. Pablo, B., Castro, B. F., Asensio, M.A. and Sanz, B. 1991. Selected chemical and microbiological change in refrigerated pork stored in carbon dioxide and oxygen enriched atmosphere. *J. Agric. Food Chem.*
- Osuna-García, J.A., Wall, M.M. and Widdell, C.A. 1998. Endogenous levels of tocopherols and ascorbic acid during fruit ripening of new Mexican-type chile (*Capsicum annum* L.). *J. Agric. Food Chem.* 46: 5093-5096.
- Parish, M.E. and Davidson, P.M. 1993. Methods for testing the efficacy of food antimicrobials. *Food Technol.* 43: 148-155.
- Parkin, K.L., Wells, M.J. and Brown, W.D. 1981. Modified atmosphere storage of rockfish fillets. *J. Food Sci.* 47: 181-184.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, M.L. 1996. Effect of carbon dioxide atmosphere on microbial growth and quality of salmon slices. *J. Sci. Food Agric.* 72: 348-352.
- Penney, N., Bell, R.G. and Cummings, T.L. 1994. Extension of the chilled storage life of smoked blue cod (*Paraper ciscolias*) by carbon dioxide packaging. *Int. J. Technol.* 29: 167-178.
- Perez-Villarreal, B. and Howgate, P.F. 1987. Spoilage of European hake (*Merluccius merluccius*) in ice. *J. Food Agric.* 41: 335-350.
- Post, L.S., Lee, D.A., Solberg, M., Furgang, D., Specchio, J. and Graham, C. 1985. Development of botulinal toxin and sensory deterioration during storage of vacuum and modified atmosphere packaged fish fillets. *J. Food Sci.* 50: 990-996.

- Prasad, M.M. and Rao, C.C.P. 1995. Storage studies as dry salts cured fish with special reference to red discoloration. *Fish Technol.* 31: 162-166.
- Prestwich, K.N. 2003. Fiber Type of Animal Muscle. Department of Biology, College of the Holy Cross. Worcester, M.A. 01610 USA.
- Raccach, M. 1984. The antimicrobial activity of phenolic antioxidants in foods: A review. *J. Food Saf.* 6: 141-170.
- Randell, K., Ahvenainen, R. and Hattula, T. 1995. Effect of the gas/product ratio and CO₂ concentration on the shelf-life of MA packed fish. *Pack. Technol. Sci.* 8: 205-218.
- Randell, K., Hattula, T., Skyttä, E., Sivertsvik, M. and Berslien, H. 1999. Quality of filleted salmon in various retail packages. *J. Food Qual.* 22: 483-497.
- Rankin, J.C and Jensen, F.B. 1993. *Fish Ecophysiology*. Fish and Fisheries Series 9. Chapman and Hull.
- Reddy, N.R., Roman, M.G., Villanueva, R.M., Solomon, H.M., Kautter, D.A. and Rhodehamel, E.J. 1997a. Shelf-life and *Clostridium botulinum* toxin development during storage of modified atmosphere-packaged fresh catfish fillets. *J. Food Sci.* 62: 878-884.
- Reddy, N.R., Solomon, H.M., Yep, H., Roman, M.G. and Rhodehamel, E.J. 1997b. Shelf-life and toxin development during storage of modified-atmosphere-packaged fresh aquacultured salmon fillets. *J. Food Prot.* 60: 1055-1063.
- Reddy, N.R., Solomon, H.M. and Rhodehamel, E.J. 1999. Comparison of margin of safety between sensory spoilage and onset of *Clostridium botulinum* toxin development during storage of modified atmosphere (MA)-packaged fresh marine cod fillets with MA-packaged aquacultured fish fillets. *J. Food Saf.* 19: 171-183.
- Reddy, N.R., Villanueva, M. and Kautter, D.A. 1995. Shelf-life of modified atmosphere packaged fresh tilapia fillets stored under refrigeration and temperature-abuse condition. *J. Food Prot.* 58: 909-914.
- Redmayne, P.C. 1989. World aquaculture developments. Quality and Safety Attributes of Aquacultured Product. *Food Technol.* 80-81.
- Rehbein, H. and Schreiber, W. 1984. TMAOase activity in tissues of fish species from the Northeast Atlantic. *Comp. Biochem. Physiol.* 79B: 447-452.
- Rico-Munoz, E. and Davidson, P.M. 1983. Effect of corn oil and casein on the antimicrobial activity of phenolic antioxidants. *J. Food Sci.* 48: 1284-1288.

- Ringo, E. and Storm, E. 1994. Microflora of Arctic charr, *Salvenus alpinus* L, gastrointestinal microflora of free-living fish, effect of diet and salinity on the intestinal microflora. *Aquacult. Fish Manage.* 25: 623-629.
- Robach, M.C., Smoot, L.A. and Pierson, M.D. 1977. Inhibition of *Vibrio parahaemolyticus* O₄: K₁₁ by butylated hydroxyanisole. *J. Food Prot.* 40: 549-551.
- Robards, K., Kerr, A.F. and Patsalides, E. 1988. Rancidity and its measurement in edible oils and snack foods. *Analyst.* 113-224.
- Robb, D.H.F., Kestin, P.D. and Warries, P.D. 2000. Muscle activity at slaughter : I. Changes in flesh colour and gaping in rainbow trout. *Aquacult.* 182: 261-269.
- Roura, S.I., Saavedra, J.P., Truco, R.E. and Crupkin, M. 1992. Conformational changes in actomyosin from post-spawned hake stored on ice. *J. Food Sci.* 57: 1109-1111.
- Ruiz-Barba, J.L., Rios-Schez, R.M., Ferdriani-Triso, C., Olias, J.M. 1990. Bactericidal effect of phenolic compounds from green olives against *L. plantarum* system. *App. Microbiol.* 13: 199-205.
- Ruiz-Capillas, C. and Moral, A. 2001a. Changes in free amino acids during chilled storages of hake (*Merluccius merluccius* L) in controlled atmospheres and their use as quality control index. *Eur. Food Res. Technol.* 212: 302-307.
- Ruiz-Capillas, C. and Moral, A. 2001b. Chilled bulk storage gutted hake (*Merluccius merluccius* L.) in CO₂ and O₂ enriched controlled atmospheres. *Food Chem.* 74: 317-325.
- Ruiz-Capillas, C. and Moral, A. 2001c. Residual effect of CO₂ on hake (*Merluccius merluccius* L.) stored in modified (MA) and controlled (CA) atmospheres. *Eur. Food Res. Technol.* 212: 413-420.
- Ruiz-Capillas, C., Moral, A. Morales, J. and Montero, P. 2002. Preservation of shelf-life of pota and octopus in chilled storage under controlled atmospheres. *J. Food Prot.* 65: 140-145.
- Sadowska, M. and Kotlowski, R. 1999. Physicochemical Properties of Collagen FormFish, Pig and Beef. *In* Gelatine-Properties, Technology and Application. (A. Rutkowski ed.). Polska Izba Dodtkow do Zywnosci.
- Sakai, T., Habiro, A. and Kawahara, S. 1999. High-performance liquid chromatographic determination of 1,3-diethyl-2-thiobarbituric acid, malonaldehyde adduct in fish meat. *J. Chromat. B.* 726: 313-316.

- Sakaguchi, M., Murata, M. and Kawai, A. 1982. Changes in free amino acids and creatine contents in yellowtail (*Seriola quinqueradiata*) muscle during ice storage. *J. Food Sci.* 47: 1662-1666.
- Santamaria, R.I., Reyes-Duarte, M.D., Barzana, E., Fernando, D., Mota, M. and Lopez-Munguia, A. 2000. Selective enzyme-mediated extraction of capsaicinoids and carotenoids from chili Guajiillo Puya (*Capsicum annum* L) using ethanol as solvent. *J. Agric. Food Chem.* 48: 3063-3067.
- Sato, K., Ando, M., Kubota, S., Orisaga, K., Kawase, H., Toyohara, H., Sakaguchi, Nakagawa, T., Makinodan, Y., Ohtsuki, K. and Kawabata, M. 1997. Involvement of type V collagen in softening of fish muscle during short-term chilled storage. *J. Agric. Food Chem.* 45: 343-348.
- Sato, K., Ohashi, C., Ohtsuki, K. and Kawabata, K. 1991. Type V collagen in trout muscle and solubility change during chilled stored of muscle. *J. Agric. Food Chem.* 39: 1222-1225.
- Sato, K., Yoshinaka, R. Sato, M. and Shimizu, Y. 1986. Collagen content in the muscle of fishes in association with their swimming movement and meat texture. *Bull. Jnp. Soc. Sci. Fish.* 52: 1595-1600.
- Sato, K., Yoshinaka, R. Sato, M., Itoh, Y. and Shimizu, Y. 1988. Isolation of types I and V collagens from carp muscle. *Comp. Biochem. Physiol.* 90B: 155-158.
- Schaneberg, B.T. and Khan, I.A. Comparison of extraction method for marker compounds in essential oil of lemon grass by GC. *J. Agric. Food Chem.* 50: 1345-1349.
- Schipp, G. 1996. Barramudi farming in the North Territory. Department of Primary Industry and Fisheries, Darwin, N.T.
- Scott, D.N., Fletcher, G.C. and Summer, G. 1984. Modified atmosphere and vacuum packaging of snapper fillets. *Food Technol. Aust.* 36: 330-334.
- Schmmedes, A. and Holmer, G. 1987. A new thiobarbutiric acid (TBA) method for determining free malondialdehyde (MDA) and hydroperoxides selectivity as a measure of lipid peroxidation. *J.A.O.C.S.* 66: 813-816.
- Shelef, L.A. 1983. Antimicrobial effects of spices. *J. Food Saf.* 6: 29-44.
- Shewfelt, R.L. 1981. Fish Muscle Lipolysis. A review. *J. Food Biochem.* 5: 79-100.
- Shewfelt, R.L., MacDonald, R.E. and Hultin. 1981. Effect of phospholipid hydrolysis on lipid oxidation in flounder muscle microsomes. *J. Food Sci.* 46: 1297-1301.

- Sheu, J-Y., Chen, P-H., Tseng, W-C., Tsai, L-Y. and Zhuang, Y-L. 2003. Spectrophotometric determination of a thiobarbituric acid-reaction substance in human hair. *Anal. Sci.* 19: 957-960.
- Sikoski, Z.E. 2001. The role of nitrogenous compounds in food quality. *In* Chemical and Functional Properties of Food Co-Protein. Chapman and Hall.
- Sikorski, Z.E. and Kostuch, S. 1982. Trimethylamine N-oxide demethylase: its occurrence, properties and role in technological changes in frozen food. *Food. Chem.* 9: 213-222
- Sikorski, Z.E. and Borderias, J.A. 1994. Collagen in the Muscle and Skin of Marine Animals. *In* Seafood proteins (Z.E Sikorski, B.S. Pan and F. Shahidi eds.) New York London. Chapman and Hall.
- Sikorski, Z.E. and Sun, P. 1994. Preservation of seafood quality. *In* Seafood: Chemistry, Processing, Technology and Quality (F. Shahidi and J.R. Botta eds.). Pp. 168-195. Glasgow: Blackie Academic and Professional.
- Silva, J.L. and White, T.D. 1994. Bacteriological and color changes in modified atmosphere-packaged refrigerated channel catfish. *J. Food Prot.* 57: 715-719.
- Simpson, J.A., Narita, S., Gieseg, S., Gebicki, S., Gebicki., J. and Dean, R.T. 1992. Long-lived reaction species on free radical damaged proteins. *Biochem. J.* 282: 621-624.
- Sinnhuber, R.O., and Yu, T.C. 1958. 2-thiobarbituric acid methods for the measurement of rancidity in fishery products 2. The qualitative determination of malonaldehyde. *Food Technol.* 12: 9-12.
- Sivakumar, P., Arichandran, R., Sugana, L., Mariappan, M. and Chandrakasan, G. 2000. The composition and characteristics of skin and muscle collagens from freshwater catfish grown in biologically treated tannery effluent water. *J. Fish Biol.* 56: 999-1012.
- Sivertsvik, M., Jeksrud, W.K. and Rosnes, J.T. 2002. A review of modified atmosphere packaging of fish and fishery products-significance of microbial growth, activities and safety. *Int. J. Food Sci. Technol.* 37: 107-127.
- Skura, B.J. 1991. Modified atmosphere packaging of fish and products. *In* Modified Atmosphere Packaging of Food (B. Ooraikul and M.E. Stiles eds.). Pp. 148-168. London: Ellis Horwood.

- Sotelo, C.G. and Rehbein, H. 2000. TMAO-degrading enzymes. *In* Seafood Enzymes: Utilization and Influence on Postharvest Seafood Quality (N.F. Haard and B.K. Simpson eds.). Marcel Dekker, Inc. New-York.
- Spencer, C.M., Cai, Y., Martin, R. and Gaffney, S.H. 1988. Polyphenol complexation: Some thoughts and observations. *Phytochem.* 27: 2397-2409.
- Stammen, K., Gerdes, D. and Caporaso, F. 1990. Modified atmosphere packaging of seafood. *Crit. Rev. Food Sci. Technol.* 29: 301-331.
- Stansby, M.E. and Griffiths, F.P. 1935. Carbon dioxide in handling fresh fish-Haddock. *Ind. Eng. Chem.* 27: 1452-1458.
- Stansby, M.A. and Olcott, H.S. 1963. Composition of fish. *In* Industry Fishery Technology (E. Robert ed.). Krieger Publishing Co., Huntington. New York.
- Statistics of Marine Fish Farms Survey. 2000. Fisheries Statistic and Farmsfish Technology Sub-Division Fishery. Economic Division Department of Fishery Ministry of Agricultural Cooperation Thailand.
- Stier, R.F., Bell, L. and Ito, K.A. 1981. Effect of modified atmosphere storage on *C. botulinum* toxigenesis and the spoilage microflora of salmon fillets. *J. Food Sci.* 46: 1639-1642
- Stiles, M.E. 1991. Scientific principles of controlled/modified atmosphere packaging. *In* Modified Atmosphere Packaging of Food (B.Ooraikul and M.E. Stiles eds.). Pp. 18-25. London: Ellis Horwood.
- Storey, R.M., Davis, H.K., Owen, D. and Moore, L. 1984. Rapid approximate estimation of volatile amines in fish. *J. Food Technol.* 19: 1-10.
- Suyama, M., Kobayashi, H. 1980. Free amino acids and quaternary ammonium base in mantle muscle of squid. *Nipp. Sui. Gakk.* 46: 1261-1264.
- Szebedinszky, C. and Gilmour, K.M. 2002. The buffering power of plasma in brown bullhead (*Ameiurus nebulosus*). *Comp. Biochem. Physiol.* 131B: 171-183.
- Szczesniak, A.S. 1998. Effect of Storage on Texture. *In* Food Storage Stability (I.A. Taub and R.P. Singh eds.) CRC. Press. LLC. United State of America.
- Tarladgis, B.G., Pearson, A.M. and Dugan, L.R. Jr. 1964. Chemistry of the 2-thiobarbituric acid test for determination of oxidative rancidity in foods. II. Formation of the TBA-malonaldehyde complex without acid-heat treatments. *J. Sci. Food Agric.* 15: 602-607.
- Tassou, C.C. and Nychas, G.J.E. 1994. Inhibition of *Staphylococcus aureus* by olive phenolics in broth and in a model food system. *J. Food Prot.* 57: 120-124.

- Tateo, F., Triangeli, L., Panna, E. and Berte, F. 1988. Stability and reactivity of aspartame in cola-type drinks. *In* *Frontiers of Flavor* (G. Charambous ed.). Elsevier Science.
- Tironi, V.A., Tomas, M.C. and Anon, M.C. 2002. Structural and functional changes in myofibrillar proteins of sea salmon (*Pseudoperca semifasciata*) by interaction with molonaldehyde (RI). *J. Food Sci.* 67: 930-935.
- Uno, K., Morishita., T. and Takahashi, T. 1987. Variation with growth in the fatty acid compositions of lipids from cultured sea bream. *Nipp. Sui. Gakk.* 53: 1609-1615.
- Valente, L.M.P., Rocha, E., Gomes, E.F.S., Silva, M.W., Oliveira, M.H., Monteiro, R.A.F. and Fauconneau, B. 1999. Growth dynamics of white and red muscles in fast and slow growing strains of rainbow trout. *J. Fish Biol.* 55: 675-691.
- Villemure, G., Simaro, R.E. and Picard, G. 1986. Bulk storage of cod fillets and gutted cod (*Gadus morhua*) under carbon dioxide atmosphere. *J. Food Sci.* 51: 317-320.
- Wang, M.Y. and Brown, W.O. 1983. Effects of elevated CO₂ atmosphere on storage of freshwater crayfish (*Pacifastacus Leniusculcs*). *J. Food Sci.* 48: 158-162.
- Wassef, E.A., Masry, E.L. and Mikhail, E.R. 2001. Growth enhancement and muscle structure of striped mullet, *Mugil cephalus* L, fingerlings by feeding algal meal-based diets. *Aquaculture Research*.32 (suppl.1): 315-322.
- Watanabe, H., Yamanaka, H. and Yamakawa, H. 1992. Post mortem biochemical changes in the muscle of disk abalone during storage. *Nipp. Sui. Gakk.* 58: 2081-2088.
- Wilknison GR. 1997. The effect of diet, aging and disease states on presystemic elimination and oral drug bioavailability in humans. *Adv. Drug Deli. Rev.* 25: 129-159.
- Woyewoda, A.D., Bligh., E.G. and Shaw, S.J. 1984. Controlled and modified atmosphere storage of cod fillets. *Can. Inst. Food Sci. Technol. J.* 17: 24-27.
- Xiao, H. and Parkin, K. 2000. Antioxidant functions of selected *Allium* thiosulfates and S-Alk (en)yl- L-cystein sulfoxides. *J. Agric. Food Chem.* 50: 2488-2493.
- Yamaguchi, S., 1979. The Umami taste. *In* *Food Taste Chemistry*. (J.C. Boudreau ed.). ACS. Symp. Ser., 115, American Chemical Society Washington, D.C. Pp. 85-97.

- Yamauchi, R., Tatsumu, Y., asano, M., Kato, K. and Ueno, Y. 1988. Effect of method salts and fructose on the oxidation of methyl linoleate in emulsion. *Agric. Biol. Chem.* 52: 849-850.
- Zaika, L.L. 1988. Spices and herbs: Their antimicrobial activity and its determination. *J. Food Saf.* 9: 97-118.
- Zhang, C.H., Hirano, T., Suzuki. and Shirai, T. 1992a. Enzymatically generated specific volatile compounds in ayu tissue. *Nipp. Sui. Gakk.* 58: 559-565
- Zhang, C.H., Shirai, T., Suzuki, T. and Hiramono, T. 1992b. Lipoxygenase-like activity and formation of characteristic aroma compounds from wild and cultured ayu. *Nipp. Sui. Gakk.* 58: 959-964.
- [http:// ataspice.org](http://ataspice.org)
- [http:// www. 10Thaidish.com/herb/Herb Spice. htm](http://www.10Thaidish.com/herb/Herb Spice. htm)