

## Contents

	Page
บทคัดย่อ	iii
Abstract	v
Acknowledgements	vii
Contents	viii
List of Tables	xi
List of Figures	xii
List of Appendices	xiv
Chapter	
1. Introduction	1
1.1 Background and rationale	1
1.2 Definition of terms	2
1.3 Review of literature	3
1.3.1 Imposex in gastropods	3
1.3.2 Macrobenthic fauna variation	5
1.3.3 The Gulf of Thailand	7
1.3.4 The Songkhla Lake	9
1.4 Objective	11
1.5 Publications of the thesis	11

## Contents (cont.)

	Page
1.6 Expected advantages of the thesis	12
1.7 Structure of the thesis	13
2. Methodology	14
2.1 Prevalence of imposex	14
2.1.1 Data source	14
2.1.2 Path diagram and research questions	15
2.2 Macrobenthic fauna abundance	16
2.2.1 Data source	16
2.2.2 Steps for model-development	19
2.3 Data managements	20
2.4 Statistical methods	20
2.4.1 Logistic regression	20
2.4.2 Multivariate multiple regression	22
2.4.3 Factor analysis	24
3. Using a logistic regression to model the prevalence of imposex	25
3.1 Description of variables	25
3.2 Characteristics of the data	26
3.3 Logistic regression model to imposex outcome	28

## Contents (cont.)

	Page
4. A multivariate multiple regression model for macrobenthic fauna density	34
4.1 Description of the variables	34
4.2 Characteristics of the data	35
4.2.1 Occurrence and abundance of macrobenthic fauna	35
4.2.2 Environmental variables	37
4.3 Factor analysis of environmental variables	40
4.4 Multivariate multiple regression model	41
5. Discussion and conclusions	47
5.1 Prevalence of imposex in gastropods in the Gulf of Thailand	47
5.2 Macrobenthic fauna density in the Middle Songkhla Lake	48
References	53
Vitae	101

## List of Tables

Table	Page
3.1 Variables and their role, data type, and categories	26
3.2 Lists of families and species of the gastropods, with a summary of total number of sample (nTotal), number of females (nFem), percentage of females (%Fem), and percentage of females showing imposex (%Imp) in 2006	27
3.3 Results of the logistic model with outcome “Imposex”	30
3.4 Prevalence of imposex (%) by area before and after adjusting for species	31
3.5 Prevalence of imposex (%) by species before and after adjusting for area	33
4.1 Variables, their roles and data type	34
4.2 The taxonomy, percentages of occurrence (%occ) and densities (ind m <sup>-2</sup> ) of the 24 macrobenthic fauna families in the Middle Songkhla Lake from April 1998 to February 1999. [* note the unidentified species in Bivalvia]	36
4.3 Factors and factor loadings for water quality parameters	40
4.4 Regression coefficients and standard errors (in parenthesis) from fitting MMR model with the four environmental predictors; coefficients with <i>p</i> -values < 0.05 are shown; those adjudged not honestly statistically significant are shown in italics and those with <i>p</i> -values < 0.01 are shown in bold; the coefficients that were not statistically significant are labeled “ns”. [* note the unidentified species in Bivalvia]	43

## List of Figures

Figure	Page
1.1 The Gulf of Thailand	8
1.2 The Songkhla Lake and three zones of the lake	10
2.1 Path diagram showing roles of variables	16
2.2 Dataset of the study of macrobenthic fauna variation	17
2.3 The Songkhla Lake and nine sampling stations (1: Ban Laem Chak, 2: Ban Koh Nang Khum, 3: Ban Nak Ka Rat, 4: Ban Koh Khob, 5: Ban Ta Kura, 6: Ban Laem Kruad, 7: Ban Hat Kai Tao, 8: Ban Ta Wa, and 9: Ban Bang Tan) within the Middle Songkhla Lake	18
2.4 Steps for development of model to assess macrobenthic fauna variations	19
3.1 The study areas and sampling sites: (A) is Overview; (B) is northern part of the Gulf with study areas A-F; (C) is Pattani Bay and surroundings with study areas H-L; (D) is Narathiwat province with study area M. Dots with their number corresponds with the sampling sites in the detailed of Appendix 4. Value between brackets gives the mean of imposex frequency in the area	32
4.1 The water characteristics: water depth (wDep), water temperature (wTemp), salinity (Sal), water pH (wpH), dissolved oxygen (DO), total suspended solids (TSS) in the Middle Songkhla Lake from April 1998 to February 1999 by station (left panel) and month (right panel)	38

## List of Figures (cont.)

Figure	Page
4.2 The sediment characteristics: sediment pH (spH), total nitrogen contents (TN), organic carbon contents (OC), percentage of clay, silt, and sand in the Middle Songkhla Lake from April 1998 to February 1999 by station (left panel) and month (right panel)	39
4.3 Residuals plotted against corresponding normal quantiles for the model	42
4.4 Plots of the observed densities (dots) and the predicted densities (dots with model fitted line) for the 24 families by the periods of observation	45
4.5 Plots of the observed densities (dots) and the predicted densities (dots with model fitted line) for the 24 families by sampling stations	46

## List of Appendices

Appendix		Page
1	Publication 1: TBT-pollution in the Gulf of Thailand: a re-inspection of imposex incidence after 10 years	65
2	Publication 2 (Manuscript): Regression-based modeling of macrobenthic fauna density in the Middle Songkhla Lake, Thailand	73
3	Publication 3: Confidence intervals for adjusted proportions using logistic regression	88
4	Names of the study areas indicated by a letter and the name of a town or village on the adjacent coast, the positional coordinates, average depth of the sampling sites, the names of the species, number of males, number of females, and number of imposex in 2006	95